



PARA-SEV

TOBACCO DUST



POISON

Contains Sevin and Parathion



POISON

DIRECTIONS

TOBACCO - Against Aphids (plant lice), cutworms, hornworms and flea beetle, use at the rate of 20 pounds per acre. Begin dusting when insects or damage is first noticed or when plants are 12 to 14 inches high. Repeat at weekly intervals or as needed. Do not use on mature tobacco plants.

CAUTION: Do not apply to tobacco within 5 days of priming and 15 days of cutting. When cutting the crop do not allow juices of treated crops to come in contact with skin or other parts of the body of those engaged in cutting the crop.

This product is toxic to fish and wildlife.

Keep out of any body of water.

Do not apply when weather conditions favor drift of dust from area treated.

Apply this product only as specified on this label.

- ANTIDOTE FOR PARATHION -
CALL A PHYSICIAN AT ONCE! (NOTE:
For physician only: Therapeutic doses of atropine appear to be effective as an antidote.)

ACTIVE INGREDIENTS by Wt.

Parathion 1.00%
(O, O-Diethyl O-P-Nitrophenyl
thiophosphate)

Carbaryl "(1-Naphthyl
N-Methylcarbamate)" 10.00%

INERT INGREDIENTS 89.00%

Total 100.00%

DO NOT RE-USE CONTAINERS—
Paper Containers should be burned
or buried. Stay well away from
smoke.

DANGER: KEEP OUT OF REACH OF CHILDREN.

May be fatal if swallowed, inhaled, or absorbed through skin! Rapidly absorbed through skin! Do not get in eyes or on skin or on clothing. Wear natural rubber gloves, protective clothing and goggles. In case of contact wash immediately with soap and water. Wear a mask or respirator of a type passed by the U. S. Department of Agriculture for parathion protection. Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. Do not contaminate feed and foodstuffs. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use.

INTERNAL: Give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have patient lie down and keep warm.

EXTERNAL: If the skin is contaminated, wash thoroughly with soap and water.

U.S.D.A. Reg. No. 1812-61