




10% SEVIN*



ACTIVE INGREDIENTS:

Carbaryl (1-Naphthyl methylcarbamate)	10%
INERT INGREDIENTS	90%
TOTAL	100%

*Trademark of Union Carbide Co. for 1-Naphthyl N-Methylcarbamate



Corn: Apply 20 lbs. per acre or 3 oz. per 50 ft. row. Corn earworm, European corn borer, fall armyworm, flea beetles, leafhopper, corn rootworms damage and foliage feeders, apply to entire plant. For damage to silks and ears, apply at 2 or 3 days after silking when silks first appear and continuing until silks begin to dry. Three or more applications may be required. DO NOT APPLY WITHIN 7 DAYS OF HARVEST OR FORAGE.

Potato, Tomato, Pepper, Eggplant: Colorado potato beetle, flea beetle, leafhopper—10-15 lbs. per acre. European corn borer, fall armyworm, lace bug, tomato fruit worms, tomato hornworm, tarnished plant bug, stink bugs—20-30 lbs. per acre. **MAY BE APPLIED TO DATE OF HARVEST.**

Tobacco: Use 10-15 lbs. per acre for flea beetle, leafhopper, fall armyworms, and stink bugs. Avoid excessive applications. If late season applications are necessary, treat immediately after priming rather than before priming.

Cucumber, Squash: Cucumber beetle, flea beetle, leafhopper, squash bug, corn earworm, stink bug—10 lbs. per acre. Avoid excessive applications. Do not apply when tender, foliage is wet or when there is prolonged high humidity.

Dogs and Cats: Ticks and fleas—Dust entire animal. Rub thoroughly into fur. Dust around sleeping quarters and wherever fleas and ticks occur. Repeat weekly if necessary. Do not treat cats or dogs less than one month old.

Sevin will not control mites. Sevin will not control insects if used with lime or highly alkaline materials.

Ornamentals: Bagworm, leaf miner, blister beetles, elm leaf aphid, flea beetle Japanese beetle, lace bug, leaf roller, plant bugs. Thoroughly dust the infected plants at the first sign of damage. Repeat weekly or as necessary for good control.

Scale control—apply in spring and early summer when scale crawlers are present.

Caution: KEEP OUT OF REACH OF CHILDREN.

Harmful if swallowed or inhaled. Do not take internally. Avoid breathing dust. AVOID prolonged or repeated contact with skin. Wash thoroughly after handling. Change contaminated clothing daily. AVOID contamination with feed and foodstuffs. Wear regular long sleeved work clothing. Wash hands & face before eating.

Do not reuse container. Destroy when empty.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.