

DIRECTIONS -- PINE TREES -- Black Turpentine Beetle.
Mix 1 gallon BHC in 14 gallons water or No. 2 fuel oil.

Do not use oil where injury to grass and foliage may result.

First, remove loose bark and trash near the ground level so that thorough coverage may be obtained. Next, apply to the bark as a coarse spray to the point of run-off. Spray from the ground up to about 18 inches on the tree. Do not soak the tree. Do not spray higher than 18 inches except in spots where there are signs of beetles. If an oil solution is used, do not spray above 5 feet on the tree as injury to limbs and needles may result.

On small trees (no larger than 4 inches in diameter), do not spray higher than 18 inches from ground.

Ips Beetle and Southern Pine Beetle. Mix 1 gallon of BHC in 55 gallons of water and spray all bark to the point of run-off.

NOTE - DO NOT SPRAY AFTER RAIN WHEN BARK IS WET.

B H C

EMULSIFIABLE CONCENTRATE

ACTIVE INGREDIENTS:	BY WT.
Gamma Isomer of Benzene	
Hexachloride	11.65%
Related Isomers of Benzene	
Hexachloride	17.10%
Xylene	67.25%
INERT INGREDIENTS:	4.00%
TOTAL	100.00%

PARRAMORE & GRIFFIN
VALDOSTA, GEORGIA

LAWN INSECTS: Fall Armyworm, Sod Webworm, and Mole Crickets. Mix ½ pint of BHC in 10 gallons of water to cover 1000 square feet of lawn. Do not increase the rates as injury to grass may occur.

CAUTION: KEEP OUT OF REACH OF CHILDREN.

Harmful if swallowed or absorbed through skin. May produce irritation of skin or eyes. Do not breathe spray mist.

Avoid contact with skin or eyes. In case of contact, wash immediately with soap and water; for eyes, flush with water and get medical attention.

Avoid contamination of feed and foodstuffs.

Do not use on household pets or humans.

Do not use on citrus, cucurbits, Irish potatoes, or on crops to be followed by potatoes or peanuts; or on forage for dairy cattle or animals being finished for slaughter.

U. S. D. A. Reg. No. 1812-X

1812-X
THE FEDERAL INSECTICIDE ACT
MAY 1947