

THIODAN® 3 DUST

DIRECTIONS FOR USE

Apply when insects first appear and repeat as required to maintain effective control. Coverage of both upper and lower leaf surfaces is essential for good control. Observe use limitations. If insect control is required beyond the following use patterns, supplement the control program with other suitable pesticides. Observe days interval between last application and harvest indicated by number in () following crop.

SUCCULENT BEANS: Mexican Bean Beetle - 15 to 20 lbs. per acre. Do not apply after pods begin to form. Do not feed treated threshings to livestock.

AULIFLOWER: Imported Cabbage Worm, Cabbage Loop, Diamondback Moth Larvae, Cross-striped Cabbage Worm, Flea Beetles, Harlequin Bugs, Stink Bugs - 25 to 33 lbs. per acre. Do not apply after edible parts begin to form. Close trim other leaves prior to marketing.

BROCCOLI (7), CABBAGE (7): Imported Cabbage Worm, Cabbage Looper, Diamondback Moth Larvae, Cross-striped Cabbageworm, Flea Beetles, Harlequin Bugs, Stink Bugs - 25 to 33 lbs. per acre.

CUCUMBERS (0): Aphids, Cucumber Beetles, Striped Cabbage Flea Beetle - 25 to 33 lbs. per acre.

EGGPLANTS (1): Blister Beetle, Colorado Potato Beetle, Flea Beetle, Green Peach Aphid, Green Stink Bug, White fly - 17 lbs. per acre.

LETTUCE, HEAD (14): Cabbage Looper, Diamondback Moth Larvae, Green Peach Aphid, Imported Cabbage Worm - 25 to 33 lbs. per acre. Do not make more than 3 applications after thinning. Remove wrapper leaves at harvest. Do not feed crop refuse to livestock.

MELONS (0): Aphids, Cucumber Beetle, Striped Cabbage, voracious Leaf Roller - 33 lbs. per acre.

ACTIVE INGREDIENT

Endosulfan Hexachlorocyclohexadiene
2, 4, 3 benzo-hexahydro-4H-pyran oxide

INERT INGREDIENTS

Thiodan U.S. Pat. No. 2,799,685

Thiodan is a registered trademark of
Farbwerke Hoechst A. G.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing of dust. Avoid contact with skin, eye, or clothing. In case of contact with skin or eyes, flush with plenty of water. For eyes, get medical attention. Wash with soap and water after handling and before eating or smoking, wear clean clothing. Avoid contamination of food and feedstuffs. Do not apply or allow drift to areas occupied by unprotected humans or bees.

Use only as directed. Do not apply to any body of water or to any water which may be used for drinking or for irrigation.

ORNAMENTALS: Aphids, Whitefly, etc. - 17 lbs. per acre. On Chrysanthemum, do not use if applied before plants flower. Do not use on flowers.

PEPPER (1): Flea Beetle, Green Peach Aphid - 33 lbs. per acre. Pepper Maggot - 17 lbs. per acre.

POTATOES (0): Aphids, Colorado Potato Beetle, Flea Beetle, including Shield Shaped Bugs, Potato Flea Beetle, Leafhopper, Potato Tuberworm, etc. - 33 lbs. per acre to 35 lbs. per acre; European Corn Borer - 25 lbs. per acre; False Chinch Bug - 17 lbs. per acre; Ringed Potato Beetle - 17 lbs. per acre.

SOYBEANS: Green Stink Bug, Mexican Bean Beetle - 25 to 33 lbs. per acre. Do not apply after pods begin to form. Do not feed treated threshings to livestock.

SQUASH (0): Aphids, Cucumber Beetles, Striped Cabbage Flea Beetle, Squash Vine Borer - 25 to 33 lbs. per acre. For Vine Borer control, apply weekly to flower buds, beginning when moths first appear.

STRAWBERRY (4): Meadow Spittlebug, Strawberry Aphid - 33 lbs. per acre. Do not apply at intervals of less than 15 days or more than twice during a season when fruit is present.

TOBACCO (5): Flea Beetles, Hornworm, Leafworm, etc. - 25 to 30 lbs. per acre. Thorough coverage is essential for good control. SEED BED - Use 2 lbs. per square yards.

TOMATOES (FIELD AND GREENHOUSE): Colorado Potato Beetle, Flea Beetle, Green Peach Aphid, Hornworms, Whitefly - 17 lbs. per acre; Tomato Fruitworm, Yellow Necked Armyworm - 25 lbs. per acre; Stink Bugs - 20 to 33 lbs. per acre; Leafminer - 25 lbs. per acre.

Peaches: (30) Green Leafhopper, etc. - 30 lbs. per acre. Do not make more than 2 applications during the fruiting period.

ACCEPTED

Feb 19, 1968

1812-100

Comments

UNDER THE FEDERAL INSECTICIDE

AND RODENTICIDE ACT

FOR ECONOMIC PEST CONTROL

ED UNDER NO. 1812-100

Potato p

re-use container.
Turn bags.
Keep from moisture.

ACCEPTED WITH COMMENTS