

50 W SEVIN*

A WETTABLE POWDER FOR SPRAYING PEACHES

ACTIVE INGREDIENTS:

Carbaryl (I-Naphthyl N-Methylcarbamate) ----- 50%

INERT INGREDIENTS: ----- 50%

TOTAL ----- 100%

* Trademark of Union Carbide Corp. for I-Naphthyl N-Methyl Carbamate

ACCEPTED
 Jan-22-1968
 1812-101

DIRECTIONS: Apply according to chart below. For best results, keep material well-agitated. Do not apply within 3 days of harvest.

Insect	Lbs. 50 W Sevin To 100 Gals. Water	Where and When To Apply
Green June Beetle	2 Lbs.	When adults appear on fruit.
Japanese Beetle	2 Lbs.	When beetles appear. Repeat in 10-14 days if needed.
Leaf Rollers	2 Lbs.	When insects become active.
Oriental Fruit Moth	2 Lbs.	3 times at 10-day intervals beginning at shuck split and at six and 3 weeks before harvest.
Plum Curculio	2 Lbs.	When application is needed close to harvest
Scales, Forbes, San Jose, Walnut and White Peach	1 Lb.	When crawlers are active.
Scales, European fruit Licanium, Terrapin and Cottony Peach	2 Lbs.	At completion of hatching. A second application may be made if infestation persists.
Stink Bugs	2 Lbs.	Petal Fall and 2-3 times 10 days apart.

Danger: KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed or inhaled. Do not take internally. Avoid breathing dust or spray mist. Avoid prolonged or repeated contact with skin. Wash thoroughly after handling. Change contaminated clothing daily. Do not contaminate foodstuffs, feed, feeding troughs or receptacles.

U.S.D.A. Reg. No. 1812-X

1812-101