

THIODAN 3 DUST PARATHION 2

DANGER-POISON KEEP OUT OF REACH OF CHILDREN

ANTIDOTE: If swallowed, Give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep quiet. Call a Physician immediately.

If on skin. In case of contact, remove contaminated clothing and immediately wash skin with soap and water.

NOTE TO PHYSICIAN: Parathion is a cholinesterase inhibition and treatment of severe poisoning should include the use of atropine.

ATROPINE IS ANTIDOTAL

Keep on hand an emergency supply of 20 tablets, grains 1 100. In case of accidental poisoning take 3 tablets (grs. 1 100) and call physician at once.

Physician. Enough atropine must be given in the shortest practical time to produce dilation of pupils. Patient must be under constant supervision and additional atropine may be necessary without delay. Clear chest by postural drainage. Use artificial respiration and oxygen if patient becomes comatose. Use of airway may be necessary.

WARNING: Poisonous if swallowed, inhaled or absorbed through skin. Rapidly absorbed through skin. Do not get in eyes, on skin or on clothing. Wear natural rubber gloves, protective clothing and goggles. In case of contact, wash immediately with soap and water. Wear a mask or respirator of a type passed by the U.S. Department of Agriculture for Parathion-Thiodan mixtures. Keep all unprotected persons and beneficial animals out of operating area or vicinity where there may be danger or drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. Do not contaminate feed and foodstuffs. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse. Highly toxic to fish and wildlife. Do not contaminate any body of water nor apply to any area not specified on the label.

USDA REG. NO. 7072-A

ACTIVE INGREDIENTS	
*Parathion	2.0%
**Endosulfan Hexachlorohexahydro- methano-2, 4, 3 benzodioxathiepin oxide	3.0%
INERT INGREDIENTS:	
	95.0%
	100.0%
*O, O Diethyl O-p nitrophenyl thiophosphate	
**Thiodan, U.S. Patent No. 2,799,685	

DIRECTIONS FOR USE

Apply 25 lbs. of this material per acre unless otherwise noted to control the following insects on the crops shown below. Apply when the air is calm by ground machinery. Thorough coverage of the plant is necessary for effective control. Observe days interval between last application and harvest indicated by number in parenthesis following crop.

TOMATOES (10): For the control of Aphids, Colorado Potato Beetle, Flea Beetle and Whitefly.

HEAD LETTUCE (14): For the control of Cabbage Looper, Green Peach Aphid, Diamond back Moth, Larvae, and Imported Cabbage Worm. Do not make more than 3 applications after thinning. Remove wrapper leaves at harvest. Do not feed crop refuse to livestock.

MELONS (7): For the control of Aphids and Cucumber Beetles.

PEPPERS (15): For the control of Flea Beetle and Green Peach Aphid.

POTATOES (5): For the control of Flea Beetle, Colorado Potato Beetle, Three Lined Beetle, Leafhopper, Aphids and Plant Bugs, Potato Tuberworm, and Whitefly.

Thiodan is a registered trademark of
Farbwerke Hoechst A.G.

DIRECTIONS FOR USE (CONT.)

CABBAGE: For the control of Cabbage Looper, Imported Cabbage worm, Diamond back Moth Larvae, and Cross-striped Cabbageworm. Note: Do not apply after edible parts begin to form.

BROCCOLI (7), CABBAGE (10): For the control of Cabbage Aphids, Cabbage Looper, Imported Cabbage-worm, Diamond-back Moth Larvae, Cross-striped Cabbage-worm Flea Beetle, Harlequin Bug, Stink Bug.

SUCULENT BEANS: Mexican Bean Beetle, Leafhopper, Aphid, Leaf Roller, Armyworms, Leaf Miner - 15 to 20 lbs. per acre. Do not apply after pods begin to form. Do not feed treated thresings to livestock.

SOYBEANS: Green Stink Bug, Mexican Bean Beetle, Velvet Bean Caterpillar - 15 to 20 lbs. per acre. Do not apply after pods begin to form. Do not feed treated thresings to livestock.

MELONS (7), CUCUMBER (15), SQUASH (15), PUMPKINS (10): Aphids, Cucumber Beetles, Striped Cabbage Flea Beetle, Squash Vine Borer, Stink Bugs, Melonworms, Pickleworm, Serpentine Leaf Miner. Do not apply to cucurbits unless plants are dry. Make no application before plants start to vine.

TOBACCO (15): Flea Beetles, Hornworm, Budworm, Green Peach Aphid. Thorough coverage is essential for good control. Make no application within 5 days of pruning. Avoid contact with plant juices when cutting or pruning.

For the control of **FLYING INSECTS** (e.g., House Fly, Stable Fly, etc.) apply 25 lbs. per acre. **NOTE:** Do not apply to fields used for pasturing of livestock or meat animals in treated fields.

DISPOSAL WARNING: Do not reuse the container. Completely empty the contents and bury the unused chemical at least 18 inches deep in an isolated location away from water supplies. Burn the empty container completely. Exercise caution and stay well away from the smoke.

2/31/68
1312