

PEANUT DUST

10% SEVIN • COPPER - S L.P.

Active Ingredients:

Copper (1-Naphthyl) (as carbonate)	1.00
Copper (as Cuprous Oxide) expressed as metal	3.16
Sulphur	0.00

Preparation of dust: 10% Sevin, 10% Copper, 10% Sulphur, 70% Diatomaceous Earth, 10% Potassium carbonate
Sulphur Fineness: 40 SS Sieve

CAUTION: KEEP OUT OF REACH OF CHILDREN

10% Sevin-Copper-Sulphur is for use on peanuts for control of Leaf spot, the Velvet bean Caterpillar (Peanut Worm) & Thrips. Consult your County Agent or Experiment Station for complete details on the peanut dusting program best suited to your local conditions. For disease control, apply dust about 30 days after planting or whenever spots appear on the leaf.

General Suggestions: Apply 15 lbs. per acre, repeated as often as needed.

CAUTION: Harmful if swallowed or inhaled.

Avoid inhaling dust. Avoid skin contact. Use respirator and goggles. Wash with mild soap and water after exposure. Keep clothing free from residues. Avoid contamination of foodstuffs. Sulphur may burn foliage when temperature is high. Do not make sulphur applications at such times.

Poise

Antidote: Induce vomiting by tickling the throat with a feather or the finger and give a quart of strong mustard water, or if available, use syrup of ipecac, 15 to 20 tablespoons; then give 1/2 cup water, milk and honey, whey or any mucilaginous drinks in copious draughts. Call on a physician or take to a hospital immediately.

Caution: Forage crops treated with this dust should not be fed to dairy animals or to animals to be slaughtered.

FOR PROTECTION OF HONEY BEES: AVOID USE, IF POSSIBLE, DURING PERIODS WHEN HONEY BEES ARE VISITING CROPS. WHEN NECESSARY TO USE DURING THIS PERIOD WARN BEEKEEPERS WELL IN ADVANCE TO LOCATE HIVES AT A SAFE DISTANCE UNTIL 1 week after application.