



10% MALATHION DUST

ACTIVE INGREDIENTS:

Malathion ° 10.00%

° (O, O-Dimethyl dithiophosphate of
diethyl mercaptosuccinate)

INERT INGREDIENTS 90.00%

TOTAL 100.00%

CAUTION: KEEP OUT OF REACH OF CHILDREN.

Harmful if swallowed. Avoid prolonged breathing of dust. Avoid prolonged or repeated contact with skin; wash thoroughly after using. Avoid contamination of feed and foodstuffs.

DIRECTIONS To control the insects listed below, dust at these recommended rates, but not less than 12 pounds per acre.

BEANS: Mexican Bean Beetle, Leafhopper and Spider Mite Apply at a rate of 17 pounds per acre.

VEGETABLE CABBAGE, BROCCOLI, TURNIP, EGGPLANT, TOMATO and POTATO: Apply at a rate not to exceed 30 pounds per acre of product for control of Aphids, Imported Cabbage Worm, Cabbage Looper and Spider Mites.

CUCUMBERS, SQUASH and MELONS: Apply at a rate not to exceed 22 pounds of product per acre for control of Aphids, and Spider Mites.

PECANS Pecan Bud Moths and Pecan Nut Case-bearers. Dust at the rate of 40 lbs. per acre. Make full coverage application. For Pecan Bud Moths, apply at 2 to 3 week intervals, when insects appear. For pecan nut case bearers, apply when first-brood larvae begin to enter nuts and repeat after 7 days if necessary.

Do not apply to plants unless they are dry, or injury may result. Do not use on any of the above crops within seven days of harvest.

This product is toxic to fish.

Keep out of lakes, streams and ponds.

Apply this product only as specified on this label.

Do not re-use empty container. Destroy it by burning with waste or burying. Stay away from smoke or fumes.

U. S. D. A. Registration No. 1812-120