

POISONOUS IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN. RAPIDLY ABSORBED THROUGH SKIN. REPEATED INHALATION OR SKIN CONTACT MAY, WITHOUT SYMPTOMS, PROGRESSIVELY INCREASE SUSCEPTIBILITY TO METHYL PARATHION POISONING.

Do not swallow or get in eyes, on skin, or on clothing. Do not breathe vapors. Wear clean, natural rubber gloves; clean waterproof protective clothing, and goggles. Replace gloves frequently and destroy used gloves. Wear a mask or respirator of a type passed by the U.S. Department of Agriculture for Methyl Parathion protection. Wash thoroughly with soap and water after handling and before eating or smoking. If the material gets into the eyes, immediately flush with water for at least ten minutes and get medical attention. In case of contact, remove all contaminated clothing and wash skin with soap and water; wash clothing before re-use. In case of spillage, wash down with large amounts of water. Decontaminate areas and equipment by washing with a concentrated solution of alkali and detergent and rinse with water.

Do not apply within 5 days of hand picking. (Workers entering cotton fields within 24 hours after application should wear protective clothing.) Do not graze dairy animals or animals being finished for slaughter on treated fields.

This product is toxic to wildlife.

Keep out of Lakes, Streams and Ponds.

Apply this product only as specified on this label.

Do not apply when weather conditions favor drift of spray.

Birds feeding on treated areas may be killed.



# 4 lb. METHYL PARATHION

## An Emulsible Concentrated Insecticide For Spray

Contains 4 Lbs. Methyl Parathion Per Gallon

### ACTIVE INGREDIENTS:

O, O-Dimethyl O-p-nitrophenyl thiophosphate -----  
Xylene -----

### INERT INGREDIENTS: -----

TOTAL -----

MIX WITH WATER BY AGITATION; AGITATE THE MIXTURE WHILE ALLOWED TO STAND. RE-AGITATE BEFORE USING.

DIRECTIONS: Cotton. Aphids, Boll Weevil, Boll worm, Flea hopper, mite, thrip and cotton leaf perforator. Use 1 quart per acre in sufficient water for thorough coverage (Approximately 3 to 10 gallons). Beet Army Worm. Use one quart per acre in sufficient water for thorough coverage. Soybeans. Stink bug, Three-cornered Alfalfa hopper, Velvet bean bug. Use 1 pint per acre in sufficient water for thorough coverage. Do not apply 20 days of harvest or grazing. Do not apply by aircraft.

**DANGER:** KEEP OUT OF REACH OF CHILDREN

Red ↑

**PARRAMORE & GRIFFIN**

VALDOSTA, GEORGIA

