



DANGER • POISON



ANTIDOTE

Note to Physicians: Active ingredient is a central nervous system stimulant and may cause convulsions. There is no specific antidote. Barbiturate acid derivatives may be used in treatment.

Danger: Hazardous if swallowed, inhaled, or absorbed through skin. Do not breathe spray mist. Do not get in eyes, on skin or on clothing. In case of contact, immediately remove contaminated clothing and flush skin or eyes with plenty of water; for eyes, get medical attention. Wash thoroughly with soap and water after handling and before eating or smoking; wear clean clothing. During commercial or prolonged exposure in spray-mixing and loading operations, wear clean synthetic rubber gloves and a mask or respirator of a type passed by the U. S. Department of Agriculture for Endosulfan protection. Do not allow to drift to areas occupied by unprotected humans or beneficial animals.

Danger: Workers entering treated areas within 24 hours of application should wear protective clothing.

**DO NOT STORE AT TEMPERATURES BELOW 20° F.
DO NOT USE IN UNDILUTED FORM.
DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME.**

This product is toxic to fish and wildlife.
Keep out of any body of water.

Apply this product only as specified on this label. This product is toxic to bees and should not be applied when bees are actively visiting the area.

Reuse equipment and containers and **DISPOSE OF WASTES** by burying at least 18 inches deep in non-crop lands away from water supplies. Do not reuse container; smash or perforate and bury with wastes.

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

Birds feeding on treated areas may be killed. Do not apply when runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes.

WARRANTY: Pee Gee warrants that this product, as of the time of sale by Pee Gee (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. Pee Gee MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

Dealers Should Sell in Original Packages Only.

U.S.D.A Reg No 1812 142

5 GAL. NET

ACCEPTED
Jan-30-1972
1812-142
COMMENTS



THIODAN® MISCIBLE

INSECTICIDE

ACTIVE INGREDIENTS:

*Endosulfan (Hexachlorohexahydromethano-2,4,3-benzodioxathiepin oxide)	24.0%
Xylene base aromatic petroleum solvent	69.0%

INERT INGREDIENTS: 7.0%

*Thiodan U. S. Patent No. 2,799,685 100.0%

Thiodan Miscible contains 2 pounds Thiodan per gallon.

Thiodan is a registered trademark of Canadian Hoechst Limited

DANGER

KEEP OUT OF REACH OF CHILDREN.

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS.

Not For Use or Storage In or Around The Home

DIRECTIONS FOR USE

Apply the listed amount per acre when insects first appear and repeat as required to maintain effective control. Use in sufficient water for thorough coverage, unless otherwise noted, for listed crops. Coverage of upper and lower leaf surfaces is essential for good control. Observe use limitations. If insect control is required beyond the following use patterns, supplement the control program with other suitable pesticides. Observe days interval between last application and harvest indicated by number in () following the crop.

When applying this material by aircraft, mix the recommended amount with sufficient water to provide a minimum of 1 gallon finished spray per acre. Where more than 2 quarts of this material are recommended, mix with sufficient water to provide a minimum of finished spray equal to twice the amount of Thiodan Miscible used.

SWEET CORN, fresh vegetable use only (0): Corn Leaf Aphid - 2 qts. Corn Earworm - 3 qts. Apply at 2 to 3 day intervals beginning when silks first appear and continuing until they begin to dry. Do not exceed 5 applications. Do not apply to sweet corn to be processed. Do not feed treated forage or ensilage to livestock or allow to graze in treated fields.

TOBACCO (5): Flea Beetle, Hornworm, Budworm, Green Peach Aphid, Cabbage Looper - **FIELD:** 1 to 2 qts. - **SEED BED:** Use 1 qt. per 100 gals. of water and apply 6 gals. spray per 100 sq. yds. Green June Bug Larvae - Use ½ qt. per 100 gals. of water and drench plant bed with 1 gal. spray per sq. yd. Stink Bug - Use field application of 2 to 3 qts. per 100 gals.

TOMATOES (Field & Greenhouse) (1): Blister Beetle, Colorado Potato Beetle, Flea Beetle, Green Peach Aphid - 1 qt. Hornworms - 1 to 2 qts. Tomato Russet Mite, Tomato Fruitworm, Yellow-striped Armyworm - 2 qts. Stink Bugs - 1½ to 2 qts. Cabbage Looper - 1½ qts. Whitefly - 1 qt. per 100 gals. in 100 to 200 gals. water.

POTATOES (0): Aphids, Colorado Potato Beetle, Plant Bugs (including Shield Shaped Bugs), Potato Flea Beetle, Potato Leaf Hopper, Potato Tuberworm, Southern Armyworm, Whitefly - 1 to 2 qts. European Corn Borer, Potato Psyllid - 1½ to 2 qts. False Chinch Bug - 2 qts. Three-lined Potato Beetle - 1 qt. per 100 gals. of water.

PARRAMORE & GRIFFIN

Valdosta, Georgia

ACCEPTED WITH COMMENTS

