

ACTIVE INGREDIENTS:

*Endosulfan [Hexachlorohexahydromethano-2.4.3-benzodioxathiepin oxide]------3.0%

INERT INGREDIENTS: ----- 97.0%

*Thiodan U.S. Patent No. 2,799,685

Total----- 100.00%

ANTIDOTE

NOTE TO PHYSICIAN Active ingredient is a central nervous system stimulant and may cause convulsions. There is no specific antidote. Bar bituric acid derivatives may be used in treatment

CAUTION: Harzardous if swallowed, inhalled, or absorbed through skint Do not breathe spray mist. Do not get in eyes, on skin or on clothing. In case of contact, immediately remove contactinated clothing and flush. skin or eyes with plenty of water, for eyes, get medical attention. Wash thoroughly with soap and water after handling and before eating or smoking, wear clean, "other g. Divring commercial or protonged exposure in spray-mixing and loading aperations, what clean synthetic rubber gloves and a mask or respirator of a typi passed by the U.S. Department of Agriculture for Endosettar calibration. No not allow to drift to are as occupied by a protected number of the left rall animals. CAUTION Worker enter gitt aterfareact, the 24 hours of appear

should wear protective clothing. DO NOT STORE AT TEMPERATURES BELOW 20 F DO NOT USE IN UNDILUTED FORM

DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME This product is toxic to fish and wildlife. Keep out of any body of

This product is toxic to bees and should not be applied when bees are actively visiting the area.

Birds feeding on treated areas may be killed.

Do not apply when weather conditions favor drift from areas treated Do not contaminate water by cleaning of equipment or disposal of waste.

Rinse equipment and containers and DISPOSE OF WASTES by burying at least 18 inches deep in non-crop lands away form water supplies. Do not reuse container; smash or perforate and bury with wastes

DO NOT ALLOY, THE MATERIAL TO DRIFT ONTO MEIGHBOR ING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IS ACCOMPANCE A THILDREL .: RECTIONS PERMUSE WINDOWS PLATE IN CROP INJURY TELEGAL RESIDERS OF MORE MORE HARRIFURESHITS MORE OCCUR

Dealers Should Sell in Original Packages Only.

TERMS OF SALE OR USE. On purchase or use of this product buyer. and user agree to the following conditions:

WARRANTY Parramore & Griffin warrants that this product, as of the time of sale by Parramore & Griffin (1) conforms to the ingredient statement on the label, and [2] is reasonably fit for the purpose set forth. in the Directions for Use EXCEPT AS SO WARRAN!ED THE PRO-DUCT IS SOLD AS IS Parramore & Griffin MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED

DANGER: KEEP OUT OF REACH OF CHILDREN.

Apply This Product Only As Specified On This Label

All amounts of material listed below are for actual Thiodan per acre. Unless otherwise stated for a particular product.

DIRECTIONS FOR USE

Apply the Ested amount per acre when insects first appear and repeat as required to maintain effective control. Coverage of upper and lower heaf surfaces is essential for good control. Observe use limitations, If insect control is required beyond the following use patterns, supplement the control program by the other scutable pesticides,

When applying this material by aircraft, use the recommended amount of finished dust per acre. Use cire to insure uniform coverage.

SWEET CORN: Corn Earworm, apply 50 lbs. dust or 1.5 lbs. actual per acre. Apply at 2 or 3 day intervals beginning when silks first appear and continuing until they begin to dry. Do not exceed 5 applications. Do not apply to sweet corn to be processed. Do not feed treated forage or ensilage to fivestock or graze in treated fields.

TOBACCO: Flea Beetle, Hornworm, Budworm, Green Peach Aphid, Cabbage Looper, and Stink Bug use field applications of a minimum of 20 ths of dust, or 0.6 lb actual per acre

TOMATOES: (Field & Greenhouse) Blister Beetle 20 lbs dust or 0.6 lbs actual per acres Colorado Potato Beetle, Flea Beetle, Green Peach Aphid and Hornworm apply 17 lbs, dust or 0.5 lb, actual per scre-Tomato Russet Mite, Tomato Fruitworm, Yellow-stripe Armyworm apply 33 bs dust or 10 lb. actual per acre Stink Bugs, Cabbage Looper, and White Fly apply 25 lbs dust or 75 lb actual per acre Do Not Treat Within One Day of Harvest

POTATO: Plant Bogs, Shield Shaped Bogs, and Potato Flea Beetle, apply 17 lbs dust or 0.5 lb actual per acre. Southern Army vorm, Aphid, apply 33 lbs. dust or 1.0 lb. actual per acre

USDA Reg No 1812 145

PARRAMORE & GRIFFIN

- Vaidosta, Georgia

ACCEPTED WITH COMMENTS





ACTIVE INGREDIENTS:

*Endosultan : Hexachtorohexahydromethand 2.4,3 benzodioxathiepin oxide) -- - -

INFRT INGREDIENTS. ------

THE TABLE OF THE CONTRACT OF T

and the first transfer to the control of the contro

Dealer Chourt Settler, enques to a serie

The state of the s

.

.

the second

970

Total - 100 00

BANGER: NETTER COMMENTED OF

IIII.DRIT.

The Part of the Pa

all amounts of anitorials of a below one for actual Thiodan per agre. Unless otherwise it and that a particular product,

DIRECTIONS FOR USE

.

2-11-71

1812-145