

Not for use or storage in or around the home.



POISON



Be Sure To Read and Understand The Precautionary Statement Before Using.

ATROPINE IS AN ANTIDOTE: Consult Physician for Emergency Supply. **Never Give Anything By Mouth To An Unconscious Person.**

FIRST AID: Call a Physician at once in all cases of Suspected Parathion Poisoning. If symptoms or signs of poisoning include blurred vision, **abdominal cramps, and tightness in the chest do not wait for a doctor** but give two atropine tablets (each 1/100 grain or 0.65 milligrams) at once. (One Tablet to children under five years of age.) Remove Patient from area where Parathion is present, Remove contaminated clothing and wash the skin clean with plenty of soap and water. If swallowed, induce vomiting. Have victim lie down and keep quiet.

NOTICE TO PHYSICIAN: Large doses of Atropine are required. For adults, give 2.0 mg. to 4.0 mg. of Atropine. Preferably by intravenous injection, repeat every 10 to 25 minutes until pupils dilate. 2-PAM is also antidotal and may be administered in conjunction with Atropine.

DANGER: Do not breath vapor or spray mist. Wear a mask or respirator of a type passed by the U. S. Department of Agriculture for Parathion protection. Do not apply or allow drift to areas occupied by unprotected humans or benficial animals. Do not store near food or feed products. Bury spillage. Do not reuse container. Destroy empty container by rinsing with lye and detergent-Crush and bury container at least 18 inches deep away from water supplies.

Do not get in eyes, on skin or on clothing.
Wear natural rubber gloves, protective clothing and goggles.
Vacated areas should not be reentered until drifting insecticide and volatile residues have dissipated.
Wash hands, arms and face before eating or smoking. Wash all contaminated clothing with soap and water before reuse.

ACCEPTED WITH COMMENTS



* Equivalent to 6.0 lbs. of 100% Parathion per gallon.

Emulsifiable Concentrate

FOR CONTROLLING CERTAIN INSECTS ON



POISON

DANGER: Keep Out of

Poisonous by skin contact, inhalation through skin. Repeated exposure may be especially hazardous. See other precautions.

ACTIVE INGREDIENTS:

- *Parathion (0,0-diethyl O-p-nitrophenyl Phosphorothioate)
- *O,O-dimethyl O-p-nitrophenyl Phosphorothioate

INERT INGREDIENTS:

Use only in manner recommended on this label.

U.S.D.A. Reg. No.



* Equivalent to 6.0 lbs. of 100% Parathion and 3.0 lbs. of 100% Methyl Parathion per gallon.

Emulsifiable Concentrate Insecticide

FOR CONTROLLING CERTAIN INSECTS ON FIELD, VEGETABLE & FORAGE CROPS



POISON



DANGER: Keep Out of Reach of Children

Poisonous by skin contact, inhalation or swallowing. Rapidly absorbed through skin. Repeated exposure may, without symptoms, be increasingly hazardous. See other precautionary on side panel.

ACTIVE INGREDIENTS:

*Parathion (0,0-diethyl O-p-nitrophenyl Phosphorothioate)	57.94%
*O,O-dimethyl O-p-nitrophenyl Phosphorothioate	29.29%

INERT INGREDIENTS: 12.77%

100.00%

Use only in manner recommended on this label because this material is dangerous.

U.S.D.A. Reg. No. 1812-154

DIRECTIONS FOR USE

This product should be applied by trained personnel using approved protective equipment. Fill spray tank half full with water then pour in specified amount of BIG 9. Keep agitator running while filling tank and during spray application. This product is intended for both ground and air applications. When used for air application use 3 to 10 gallons per acre of mixture.

COTTON: For control of Cotton Leafworms, Fleahoppers, Lygus Bug, Red Spider Mites, Thrips and Ground Webworms, use 2/3 to 1 1/3 pints per acre at 4 to 5 day intervals until control is obtained. Do not apply within 5 days of hand picking. Workers entering fields within 24 hours after application should wear protective clothing.

CORN: For control of Rootworm Adults, Aphids, Armyworms, Fall Armyworms, Budworms, Spider Mites, Cutworms and European Corn Borers, use 1/2 to 3/4 pint per acre with sprays at 5 to 7 day intervals.

SOYBEANS: For control of Two-Spotted Mites, Stink Bugs, Bollworms, Fall Armyworms, Green Caterpillars, Velvet Bean Caterpillars and Webworms, use 1/2 to 1 pint per acre. 20 days limitation prior to harvest.


TOMATOES: For control of Aphids, Armyworms, Cabbage Loopers, Grasshoppers, Leaf Miners, Spider Mites, Tomato Russet Mites, Whiteflies, Plant Bugs and Psyllids, use 1/2 pint per acre. 10 days limitation prior to harvest.

TURNIPS: For control of Aphids, Cabbage Loopers, Cabbage Webworms and Vegetable Weevils, use 1/4 to 1/2 pint per acre. 10 days limitation or 21 days if tops are to be used for food.

CABBAGE AND COLE CROPS (Broccoli, Brussels Sprouts, Cauliflower, Kohlrabi): For control of Aphids, Cabbage Loopers and Vegetable Weevils, use 1/4 to 1/2 pint per acre. For Loopers, Armyworms and Flea Beetles, use 1/2 to 1 pint per acre. 10 days limitation at 1/2 pint dosage, 21 days limitation at 1 pint dosage.

PEACHES: For control of Aphids, Peach Tree Borers, Plum Cuckoo and Oriental Fruit Moths, use 1/2 pint per 100 gallons of water. For Peach Tree Borers apply 2 or 3 sprays to trunk from ground to scaffold limbs timed with moth flight. For use in areas other than California. Do not apply more than 5 pounds actual per acre per year. 14 days limitation prior to harvest.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply when weather conditions favor drift from area treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes.



PARRAMORE & GRIFFIN
Valdosta, Georgia