



# CHLORDANE 10% GRANULAR

ACTIVE INGREDIENT:

°Technical Chlordane	-----	10%
INERT INGREDIENTS	-----	90%
		100%

\*equivalent to 6% Octachloro-4, 7-Methanotetrahydroindane and 4% Related Compounds.

## CAUTION: KEEP OUT OF REACH OF CHILDREN

**CAUTION:** Hazardous if swallowed, inhaled, or absorbed through skin! In case of contact, immediately remove contaminated clothing and flush skin or eyes with plenty of water; for eyes, flush with plenty of water for at least 15 minutes, get medical attention. Wash thoroughly with soap and water after handling and before eating or smoking; wear clean clothing. If material is to be handled, wear protective clothing, goggles, and clean synthetic rubber gloves. During commercial or prolonged exposure in loading operations wear a mask or respirator of a type passed by the U. S. Department of Agriculture for Chlordane protection. Keep away from children and domestic animals. Do not store near feed and foodstuffs.

Do not apply to areas occupied by unprotected humans or beneficial animals. This product is poisonous to fish and wildlife. Keep out of lakes, ponds, and streams. Do not apply in any manner not specified on the label. Birds feeding on treated areas may be killed.


### RECOMMENDATIONS

**INSECT CONTROL ON TURF, NURSERY AND GREENHOUSE SOILS:** For the control of Ants, Armyworms, Chiggers, Chinch Bugs, Crickets, European Earwigs, Grasshoppers, Japanese Beetle Grubs, Mosquitos, Rose Chafer, Sod Webworm, Spiders, Ticks, White-fringed Beetle, White Grubs, and Wireworms in Turf (such as Golf Courses, Athletic Fields, Large Estate Nurseries, Cemeteries and Park Sod), and in Nursery and Greenhouse soils where ornamental plants are grown, use at the rate of 50 lbs. of Chlordane 10% Granular per acre (1¼ lbs. to each 1000 sq. ft.) to cover surface uniformly, using a lawn fertilizer spreader. Treatment should be applied in the spring or when insects are observed to be troublesome. Do not treat soil when it is wet. Apply before rain or sprinkling. Do not apply to newly seeded lawns.

**INSECT CONTROL ON HOME LAWNS:** For the control of the above listed insects on home lawns apply 50 lbs. of Chlordane 10% Granular per acre (1¼ lbs. to each 1000 sq. ft.) to cover surface uniformly using a lawn fertilizer spreader. Treatment should be applied in the spring, or when insects are observed to be troublesome. Immediately after treatment, drench lawn with water or sprinkle heavily to make treatment more effective and to reduce the hazard to children and pets. Do not treat when soil is wet. Apply before rain or sprinkling.

**CAUTION:** Do not permit children or pets to go on treated lawn until the insecticide has been washed off the grass into the soil by drenching or sprinkling with water and grass has completely dried.

Manufactured By  
**PARRAMORE & GRIFFIN**  
Valdosta, Georgia



E.P.A. Reg. No. 1812-164