

TOBACCO 5% SEVIN*

ACTIVE INGREDIENTS:

*Carbaryl (1-naphthyl N-methylcarbamate) 5.0%
 **Erdosulfan (hexachloronexahydrometnano-
 2,4,3-benzodioxathiepin oxide) 1.5%
 INERT INGREDIENTS: 93.5%

*Sevin, U.S. Pat. Nos. 2,903,478; 3,009,855

**Thiodan, U.S. Pat. No. 2,799,685

100.0%

Insecticide Dust

DUST 1.5% THIODAN®

CAUTION Keep out of reach of children

DIRECTION FOR USE

Apply dust when insects first appear and repeat at 7 to 10 day intervals as required to maintain control. Apply thoroughly to upper and lower leaf surfaces. Observe days interval between last application and harvest indicated by number in parenthesis () following crop listed.

TOBACCO: (Field) Hornworms, Budworms, Flea Beetles, June Beetles, Japanese Beetles - Use 20 to 40 lbs. of dust per acre. Apply when insects first appear and repeat at 7 to 10 day intervals as required to maintain control. Use the lower dosage on small plants and the higher dosage as plants increase in size. Avoid heavy applications just before priming leaves. Avoid excessive applications in the bud of the plant as injury may result. Avoid applications during rain or during periods of high humidity as injury to tender foliage may result. Do not apply within 5 days of priming or cutting.

(Plant Beds): Tobacco Flea Beetle - Use 1 to 2 lbs. per 100 square yards when plants emerge and repeat at 10 to 14 day intervals up to 5 days before pulling transplants.

COTTON: Boll Weevil, Bollworm, Cotton Fleahopper, Lygus Bugs, Thrips, Pink Bollworm - Use 30 to 40 lbs. of dust per acre. Use the lower dosage when plants are small and the higher dosage when plants increase in size. Apply when insects first appear and repeat at 3 to 7 day intervals as required to maintain control. Do not graze meat or dairy animals in treated fields. Do not apply after 25% of the bolls open.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Birds feeding on treated areas may be killed. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

Highly toxic to fish and wildlife. Do not contaminate any body of water nor apply when conditions favor drift from the crop area treated.

Hazardous if swallowed, inhaled or absorbed through skin. Avoid breathing of dust. Avoid contact with skin, eyes or clothing. In case of contact with skin or eyes, flush with plenty of water. For eyes, get medical attention. Wash with soap and water after handling and before eating or drinking; wear clean clothing. Avoid contamination of feed and foodstuffs. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals.

Do not use empty container. Destroy it by burning with waste or burying, away from crops and water.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. It may cause ataxia, tremor. Thiodan produces central nervous system depression, respiratory depression, tachycardia and dilatation of the pupils.

CAUTION: Do not use in or near water for 24 hours after application. Do not use in or near water.

E.P.A. No. 1812-166

*Trademark of Union Carbide Corporation for the active ingredient carbaryl.

The dusts are registered trademarks of Farbwerke Hoechst A.G.



PARRAMORE & GRIFFIN CO.
 VALDOSTA, GEORGIA

NET WEIGHT 50 LBS.

ACCEPTED

6/26/72

1812-166

TOBACCO 5% SEVIN*

ACTIVE INGREDIENTS:	
Chlorpyrifos methyl (90% active)	5.0%
Inert ingredients	95.0%
TOTAL	100.0%

DUST THIODAN®

CAUTION Keep out of reach of children.

DIRECTION FOR USE

Apply dust when insects first appear and repeat at 7 to 10 day intervals as required to maintain control. Apply thoroughly to upper and lower leaf surfaces. Observe days interval between last application and harvest indicated by number in parenthesis () following crop listed.

TOBACCO (leaf) Fleaworms, Budworms, Flea Beetles, June Beetles, Japanese Beetles. Use 20 to 40 lbs. of dust per acre. Apply when insects first appear and repeat at 7 to 10 day intervals as required to maintain control. Use the lower dosage on small plants and the higher dosage as plants increase in size. Avoid heavy applications just before picking leaves. Avoid excessive applications in the bud of the plant as rot may result. Avoid applications during rain or during periods of high humidity as injury to tender foliage may result. Do not apply within 7 days of picking or cutting.

(Plant Bed) Tobacco Flea Beetle Use 1 to 2 lbs. per 100 square yards when plants emerge and repeat at 7 to 10 day intervals up to 10 days before setting transplants.

COTTON (leaf) Bollworm, Cotton Fleaopper, Lygus Bugs, Thrips, Pink Bollworm. Use 20 to 40 lbs. of dust per acre. Use the lower dosage on small plants and the higher dosage when plants are in full bloom. Apply when insects first appear and repeat at 7 to 10 day intervals as required for control. Do not graze or feed to livestock within 30 days of application. Use 20 lbs. of

6/20/72
1812166



PARVAVOIRE & GRIFFIN CO.

NEW YORK, N.Y.