



SUL-CO-TOX

ACCEPTED

AUG 7 1972

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. 1812-168 SUBJECT
TO ATTACHED COMMENTS.

ACTIVE INGREDIENTS:

Toxaphene (Technical Chlorinated Camphene)	
(Chlorine Content 67% to 69%)	20.0%
Copper, (as Basic Copper Oxide) Expressed as Metallic	3.4%
Sulphur	36.0%

INERT INGREDIENTS: **40.6%**

TOTAL **100.0%**

CAUTION: KEEP OUT OF REACH OF CHILDREN

DIRECTIONS: PEANUTS: To control fall armyworms, cutworms and thrips on peanuts, apply 15 lbs. per acre. This dust is also a fungicide dust for the control of Peanut Leaf Spot. Consult State Extension Service Specialist for complete details on the best peanut dusting program for your local conditions.

These are general suggestions: Apply dust about sixty days after planting, or as soon as the spots appear on the leaf. First Dust—apply 15 to 20 pounds per acre. Later Dusts—apply 20 to 30 pounds per acre. A minimum of three to four applications should be made at two week intervals. Peanuts which have received late application of this insecticide should not be grazed by dairy animals or by animals being finished for slaughter.

Harmful if Swallowed. Harmful dust may be absorbed through the skin, may cause irritation of skin and eyes. Avoid inhaling or getting into eyes. Avoid skin contact. Avoid contamination of food and feed products. Keep clothing free from residues. For eyes, flush with water and get prompt medical attention. Wash after handling. Do not store near or contaminate feed or foodstuffs.

ANTIDOTE: INTERNAL: Give one tablespoon of salt in a glass of water and repeat until vomit is clear.
EXTERNAL: Wash with soap and water.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

This product is toxic to fish, birds and other wildlife. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.



SUL-CO-TOX

ACCEPTED
AUG 7 1972
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. 1812-168 SUBJECT
TO ATTACHED COMMENTS.

ACTIVE INGREDIENTS:

Toxaphene (Technical Chlorinated Camphene) (Chlorine Content 67% to 69%)	20.0%
Copper, (as Basic Copper Oxide) Expressed as Metallic	3.4%
Sulphur	36.0%
INERT INGREDIENTS:	40.6%

TOTAL **100.0%**

CAUTION: KEEP OUT OF REACH OF CHILDREN

DIRECTIONS: **PEANUTS:** To control fall armyworms, cutworms and thrips on peanuts, apply 15 lbs. per acre. This dust is also a fungicide dust for the control of Peanut Leaf Spot. Consult State Extension Service Specialist for complete details on the best peanut dusting program for your local conditions.

These are general suggestions: Apply dust about sixty days after planting, or as soon as the spots appear on the leaf. First Dust—apply 15 to 20 pounds per acre. Later Dusts—apply 20 to 30 pounds per acre. A minimum of three to four applications should be made at two week intervals. Peanuts which have received late application of this insecticide should not be grazed by dairy animals or by animals being finished for slaughter.

Harmful if Swallowed. Harmful dust may be absorbed through the skin, may cause irritation of skin and eyes. Avoid inhaling or getting into eyes. Avoid skin contact. Avoid contamination of food and feed products. Keep clothing free from residues. For eyes, flush with water and get prompt medical attention. Wash after handling. Do not store near or contaminate feed or foodstuffs.

ANTIDOTE: **INTERNAL:** Give one tablespoon of salt in a glass of water and repeat until vomit is clear.
EXTERNAL: Wash with soap and water.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

This product is toxic to fish, birds and other wildlife. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

PARRAMORE & GRIFFIN

E.P.A. REG. NO. 1812-168

Valdosta, Georgia