



Quality

ZINEB 6.5

Shade Tobacco Dust



Pesticides

ACTIVE INGREDIENT(S)	
Zineb (zinc ethylene bisdithiocarbamate)*	6.5%
INERT INGREDIENT(S)	<u>93.5%</u>
TOTAL	100.0%
<small>*(metallic zinc equivalent 1.54%)</small>	

DIRECTIONS

IN THE FIELD, to control Blue Mold and Anthracnose diseases, apply on newly-set plants at the rate of 7 to 8 pounds per acre, increasing the rate to 15 to 20 pounds per acre as the plants mature. Repeat applications as necessary. For more detailed information consult your State Experiment Station or Extension Service.

IN THE PLANT BED, apply to plants before the disease strikes, but in any event prior to the time when the plants reach 4-leaf stage at the rate of 1½ to 2 pounds per 100 square yards, increasing to 3½ pounds per 100 square yards when the plants are large or at the first sign of Tobacco Blue Mold. Repeat applications twice a week until the crop is transplanted to the field. Repeat applications following rain and be sure to give uniform and thorough coverage. For more detailed information consult your State Experiment Station or Extension Service.

CAUTION

May cause irritation of nose, throat and skin. Do not breathe dust. In case of prolonged or frequent use, wear clean, protective clothing and bathe after each exposure. Keep containers closed. Keep away from fire and sparks.

DO NOT REUSE CONTAINER. DESTROY WHEN EMPTY.

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

PARRAMORE & GRIFFIN
VALDOSTA, GEORGIA
E.P.A. Reg. No. 1812-174

CAUTION

Keep out of reach of children

USE ONLY AS SPECIFIED ON THIS LABEL

Tested

50 Lbs. Net

Brand

Dec 8, 1972
1812-174



Quality

ZINEB 6.5

Shade Tobacco Dust



Pesticides

ACTIVE INGREDIENT(S)	
Zineb (zinc ethylene bisdithiocarbamate)*	6.5%
INERT INGREDIENT(S)	<u>93.5%</u>
TOTAL	100.0%
*(metallic zinc equivalent 1.54%)	

DIRECTIONS

IN THE FIELD, to control Blue Mold and Anthracnose diseases, apply on newly-set plants at the rate of 7 to 8 pounds per acre, increasing the rate to 15 to 20 pounds per acre as the plants mature. Repeat applications as necessary. For more detailed information consult your State Experiment Station or Extension Service.

IN THE PLANT BED, apply to plants before the disease strikes, but in any event prior to the time when the plants reach 4-leaf stage at the rate of 1½ to 2 pounds per 100 square yards, increasing to 3½ pounds per 100 square yards when the plants are large or at the first sign of Tobacco Blue Mold. Repeat applications twice a week until the crop is transplanted to the field. Repeat applications following rain and be sure to give uniform and thorough coverage. For more detailed information consult your State Experiment Station or Extension Service.

CAUTION

May cause irritation of nose, throat and skin. Do not breathe dust. In case of prolonged or frequent use, wear clean, protective clothing and bathe after each exposure. Keep containers closed. Keep away from fire and sparks.

DO NOT REUSE CONTAINER. DESTROY WHEN EMPTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

PARRAMORE & GRIFFIN
VALDOSTA, GEORGIA
E.P.A. Reg. No. 1812-171

Dec 8, 1972

CAUTION

Keep out of reach of children

USE ONLY AS SPECIFIED ON THIS LABEL

Tested

1812-174

50 Lbs. Net

Brand