



ACCEPTED  
 May 15, 1975  
 FARMER'S UNION  
 1512-175

**DIRECTIONS**

This material may be used with conventional ground or air equipment, ensuring the wind velocity not to be greater than 5 miles per hour.  
 Begin applications when peanuts are 40 to 45 days beyond planting. Use 1 to 1 1/2 gallons in 15 to 25 gallons of water per acre. Recommended application periods each 14 days after first treatment.  
 There is no tolerance time on this material and it may be applied through the entire growing season.  
 These directions are broad in their recommendations, and we suggest your local extension service be contacted for more detail information concerning frequency and number of applications per season.

**COPPER SULFUR**

ACTIVE INGREDIENTS: Cupric Hydroxide 20.50%  
 Sulfur 17.30%  
 INERT INGREDIENTS: 62.20%  
 TOTAL 100.00%

**CAUTION!**

**Keep Out of Reach of Children**

Do not swallow. Avoid breathing any mist or fumes from the throat or skin.  
 Avoid contact with eyes.  
 This product is toxic to fish. Keep out of any body of water. Decontaminate by cleaning of equipment or disposal of wastes. Apply the product only as directed on this label.  
 Keep away from heat and open flame.  
 NOTICE: Do not use on ornamentals or plants in flower.

**CERCOSPORA  
 LEAF SPOT  
 CONTROL  
 FOR PEANUTS**



**Mark of Quality**

FARRAMORE & GRIFFIN  
 DALHOUSA, GEORGIA

Net Contents 5 Gallons

F.P.C. 151-112-115