

**NEW**

# SUPER SIX Liquid Sulfur

## RECOMMENDATIONS

**PEANUTS:** For rust use 2 to 4 pints per acre of Super Six Sulfur in enough water for good coverage. Begin 40 to 60 days after planting. Use 7 to 14 day intervals, as long as needed to control rust.

**CITRUS:** Oranges, Lime, Grapefruit, Lemons, Tangelos, and Tangerines. To control Rust Mite use 5 pints per acre in enough water for good coverage.

**PEACHES:** To control Powdery Mildew, Brown Rot, Scab. For Petal Fall and Shuck Spray, begin at Pink Bloom apply 1½ pints per acre Super Six in enough water for good coverage. Cover Spray, use 1 to 1¾ pints per acre in enough water for good coverage. May be used as long as needed.

**DO NOT USE WITHIN 2 WEEKS OR 14 DAYS OF AN OIL SPRAYING.**

June 11, 1975  
 UNDER THE FEDERAL INSECTICIDE  
 FUNGICIDE AND RODENTICIDE ACT  
 FOR ECONOMIC POISON REGISTERED  
 UNDER NO. 1812-196

## CAUTION:

**Keep out of reach of Children**

Harmful if swallowed. Avoid breathing spray mist. Contact with skin may cause irritation. Avoid contact with eyes and skin.

**Read the entire label**

Do not reuse container. Dispose by burying in areas away from water supply.

**NOT FOR HOME USE**

**ACTIVE INGREDIENT:** Sulfur as S. . . 52.0%

**INERT INGREDIENT:** . . . . . 48.0%

**Contains 6.0 pounds Sulfur per Gallon**

**LIQUID**

**PARRAMORE & GRIFFIN — Bainbridge, Ga.**

## CAUTION

Sulfur may cause injury if used improperly or under unfavorable weather conditions. During periods of high temperature Sulfur may burn foliage.

**DO NOT STOP AGITATION WITH SULFUR IN SPRAY EQUIPMENT. SULFUR IS HIGHLY CORROSIVE, AND EQUIPMENT SHOULD BE CLEANED THOROUGHLY AFTER EACH DAYS SPRAYING.**



E.P.A. Reg. No. 1812-196

E.P.A. Est. No. 1812-Ga.-2