



PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
CAUTION

Harmful if swallowed, absorbed through the skin, or inhaled. Avoid breathing vapor or spray mist. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Do not apply this product in a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by or protected persons. Do not enter area without protective clothing until sprays have dried. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

RE-ENTRY STATEMENT

Do not enter treated areas without protective clothing for 24 hours after application. Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information. Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. (Indicate specific oral warnings which inform workers of areas or fields that may not be entered without specific protective clothing, period of time field must be vacated and appropriate actions to take in case of accidental exposure). When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: **CAUTION. Area treated with SUPER SIX[®] on (date of application).** Do not enter without appropriate protective clothing for 24 hours. For further information see Precautionary Statements on the label.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

PRODUCT INFORMATION

GRIFFIN SUPER SIX[®]: Liquid Sulfur is a flowable sulfur formulation that may be applied as a ground or aerial application. Consult your State Agricultural Experiment Station or Extension Specialist for advice in selecting treatments from this label to best fit local conditions. Do not combine with emulsifiable liquids except on dormant sprays or delayed dormant spray applications.

USE ONLY ON CROPS LISTED AND ONLY AS SPECIFIED ON THIS LABEL. EXCEPT ON COTTON, do not apply when temperatures exceed or are likely to exceed 90°F. Do not use sulfur with oil or within 14 days of an oil spray (for citrus 21 days). Do not combine with emulsifiable liquids.

NOTE: Sulfur will cause severe fruit and leaf injury to sulfur-sensitive crops. Do not apply or allow to drift to apricots, d'Anjou and Comice pears, cranberries, cucurbits (cucumber, cantelopes, melons, squash), filberts, spinach, tung trees, walnuts or other sensitive plants. Sulfur may burn foliage and fruit during periods of high temperatures and under certain climate conditions.

SUPER SIX[®]: Liquid Sulfur can be mixed and applied with liquid fertilizer or water. **DO NOT STOP AGITATION WITH SULFUR IN SPRAY EQUIPMENT. SULFUR IS HIGHLY CORROSIVE, AND EQUIPMENT SHOULD BE CLEANED THOROUGHLY AFTER EACH DAY'S SPRAYING.**

RECOMMENDATIONS

TREE FRUITS

Unless otherwise specified for specific crops, dosage rates are given as pints of **SUPER SIX[®]:** Liquid Sulfur per 100 gallons water for use in a thorough coverage spray. The total amount of **SUPER SIX[®]:** Liquid Sulfur per acre should be 2 to 4

Super Six®

Liquid Sulfur

ACTIVE INGREDIENT:

Sulfur as S 52.0%

INERT INGREDIENTS: 48.0%

TOTAL 100.0%

This product contains 6 lbs. of Sulfur per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 to 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

IF IN EYES: Flush with plenty of water. Get medical attention if irritation persists.

Griffin

GRIFFIN CORPORATION

VALDOSTA, GEORGIA 31601

NET CONTENTS 30 Gallons

EPA Reg. No. 1812-196

EPA Est. No. 1812-GA-3

gallons depending on the size of the trees. Because of variations in the types of spray equipment used, a range of low and high rates is listed. For high volume sprayers (output 800 to 1000 gallons spray per acre) use the low rate; for low volume sprayers (output 20 to 300 gallons spray per acre) use the high rate. This product should be applied by ground equipment, aircraft or sprinkler irrigation systems in sufficient water to provide thorough coverage.

APPLE, PEAR:

Scab:
Pre-bloom through calyx sprays 1 1/2 to 3 1/2 pints
Cover Sprays 1/2 to 2 pints

Powdery Mildew:

Pre-bloom through calyx sprays 3 pints
Cover Sprays 2 pints

CHERRY, PLUM, PRUNE:

Brown Rot, leafspot:
Pink and Bloom Sprays 1 1/2 to 3 1/2 pints
Petal fall, shuck & cover sprays 1/2 to 1 1/2 pints

GRAPES

CITRUS:

Rust Mite, clover mite:

Apply sprays November thru May

1/2 to 4 1/2 pints

THRIPS: Early spring spray for the reduction of Thrips population. Apply 1 to 2 gallons per acre in sufficient water for thorough coverage. Make application after spring flush has made 3 to 4 inch growth. Repeat if necessary. Do not apply within 21 days of an oil application.

PEACH, NECTARINE:

Powdery Mildew, brown rot, scab:

Pink and Bloom Sprays:

1 1/2 to 3 1/2 pints

Petal Fall, shuck and cover sprays

1/2 to 1 1/2 pints

Application to mature nectarines may cause discoloration.

FIELD AND VINE CROPS:

Unless otherwise specified for specific crops, dosage rates are given as pints of SUPER SIX® Liquid Sulfur per acre. This product should be applied by ground equipment, aircraft or sprinkler irrigation systems in sufficient water to provide thorough coverage.

BEANS (dry, colored, green, snap, lima): Powdery mildew, Rust: Use 4 to 8 pints of SUPER SIX® Liquid Sulfur per acre in sufficient water for thorough coverage. Begin application at the first sign of disease. Repeat at 7 to 14 day intervals to keep new growth protected.

COTTON: Atlantic Mite. Use 2 pints for early season control and up to 1 gallon for mid to late season control.

GRAPES: Powdery Mildew. Use 1 to 2 pints per 100 gallons water, maximum of 8 pints SUPER SIX® Liquid Sulfur per acre in spring and summer. Make first application when shoots are 6 to 8 inches long, second when shoots are 12 to 15 inches long, and third at about bloom stage. The timing, rate and number of applications will vary with local conditions. Consult State Agricultural Extension Service or State Agricultural Experiment Stations in your region for specific information. Follow the USE PRECAUTIONS given on this label.

PEANUTS: For rust use 2 to 4 pints per acre of SUPER SIX® Liquid Sulfur in enough water for good coverage. Begin 40 to 60 days after planting. Use 7 to 14 day intervals as long as needed to control rust.

POTATOES: Powdery Mildew. Use 3 to 4 pints in 8 gallons water per acre by air. Apply at the first sign of infection and repeat at 3 week intervals.

SUGAR BEETS: For control of powdery mildew, apply the recommended rate in 10 gallons of water by air, 10-20 gallons of water by ground at first sign of infection. West, Northwest and Midwest, use 8 pints of SUPER SIX® with recommended rates of water. Repeat applications should be made at 10-30 day intervals or as necessary throughout the season. Under most disease conditions only one or two applications will be needed.

Sprinkler Irrigation Water — Use 1-2 gallons per acre by injecting into sprinkle system at or near the irrigation pump. For center pivot systems, apply during one full revolution of the system. For portable or solid set systems, apply during the first 15 minutes of the irrigation period. Do not operate irrigation system after application until deposit has thoroughly dried. Apply the product through irrigation system containing: 1) anti-siphon and check valves which will prevent water source contamination and overflow of the slurry tank, 2) interlocking controls between the metering device and the water pump to insure simultaneous shut-off.

STRAWBERRIES: Powdery Mildew. Use 2 pints. Apply at first sign of infection and repeat at 3 week intervals. For Strawberries that will be canned, discontinue application of sulfur well before harvest in accordance with local canners' recommendations.

TOMATOES: Tomato russet mite. Use 1/2 to 1 gallon per acre depending upon size of plants. Use full coverage sprays.

NOTICE OF WARRANTY

GRIFFIN warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of GRIFFIN. In no case shall GRIFFIN be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. GRIFFIN MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

NEWARK, N.J.
GRIFFIN