

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
CAUTION

If swallowed, Avoid inhalation of dust or spray mist. Use with care. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling. Do not feed treated food or feed products. Keep out of reach of children.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and wildlife. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply where weather conditions favor drift from treated area. Do not apply to water with fish or wildlife. Do not dispose of this product in water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with the directions on the label.

STORAGE AND DISPOSAL

Store in a cool, dry place. Protect from excessive heat.

Do not contaminate water, food or feed by storage or disposal. Pesticide spray mixture or residue that cannot be used in accordance with label instructions must be disposed of according to Federal, State or local procedures under the Resource Conservation and Recovery Act. Dispose of empty containers according to approved Federal, State or local procedures under the Resource Conservation and Recovery Act.

GENERAL INFORMATION

Read all precautions and directions for use before using. Use only for claims listed and only as specified on this label.

Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers.

Fire hazard may at times occur to Red Delicious, Winesap and other sensitive varieties of apples. Do not apply under conditions involving possible drift to food for other plantings that might be damaged or the crops thereof rendered unfit for sale or consumption.

Compatibility: Griffin Captan 50W may be combined with most commonly used insecticides, adjuvants and fungicides with the exception of strongly alkaline materials such as hydrated lime, which reduces the fungicidal activity. Do not use this product in combination with oil sprays or with solvent formulations of Parathion products. Do not use this product on foliage closely following oil sprays.

NOTE: For concentrate sprayer applications, the same quantity (per acre) of Griffin Captan 50W as would be normally applied when spraying dilute mixture should be applied to each acre in sufficient water for coverage.

ALMONDS: Brown Rot, Shot Hole, Scab, Leaf Blight - 2 lbs. per 100 gals. spray. Apply at popcorn, bloom, petal fall stages and up to 5 weeks after petal fall. May be used up to 12 days of harvest if hulls are not fed to dairy animals or animals being finished for slaughter.

APPLES: Primary Scab Infection - 1 1/2 to 2 lbs. per 100 gals. spray. Secondary Apple Scab, Brooks Fruit Spot, Flyspecking - 1 lb. per 100 gals. spray. Severe Infection - 2 lbs. per 100 gals. spray. Bitter Rot, Black Rot, Botryosphaeria, Black Pox - 2 lbs. per 100 gals. spray. Bull's Eye Rot, Botrytis Rot - 2 lbs. per 100 gals. spray. Make 1 or 2 applications with late cover sprays and 1 or 2 preharvest applications. Where lead arsenate has been used in the apple maggot cover sprays, use Griffin Captan 50W 2 lbs. per 100 gals. spray. Do not use lead or arsenate within 30 days of harvest. Griffin Captan 50W should not be used in combination and/or closely following or in alternation with wettable sulfur products or sulfur sensitive varieties of apples such as Red Delicious, Stayman, Baldwin, King, etc. as severe foliage injury and defoliation may occur.

APRICOTS: Brown Rot (Twig Blight), Jacket Rot - 2 lbs. per 100 gals. spray. Apply at red bud stage, bloom and repeat at 75 per cent petal fall.

ASPARAGUS: Preplanting Root Dip Stimulation of shoot growth - 1 1/2 qts. to 100 gals. water. Dip root sections for 1 minute, drain and plant.

BEANS, PEAS: Damp Off on Beans, Root Rot on Beans and Peas - Broadcast 10 to 12 lbs. per acre. Work into upper 3 or 4 inches of soil or apply 5 to 6 lbs. per acre in row at planting time.

CANTALOUPE, CUCUMBERS, SUMMER SQUASH, WINTER SQUASH, WATERMELONS, PUMPKINS: Anthracnose, Downy Mildew - 2 lbs. per 100 gals. spray or apply 4 lbs. per acre in sufficient water for coverage. Apply at first sign of disease. Repeat at 5 to 7 day intervals or as necessary to keep diseases under control.

CITRUS: Septoria Leaf Spot, Cercospora Blight - 2 lbs. per 100 gals. spray. Apply 4 lbs. per acre in water to cover. Apply at first sign of disease. Repeat at 6 to 10 day intervals.

CELERY: Late Blight (Septoria), Pink Rot (Sclerotinia) - 2 lbs. per 100 gals. spray. Apply 200 diluted gals. per acre or equivalent of 4 to 4 1/2 lbs. per acre in water for coverage. Apply at first signs of diseases. Repeat at 7 to 10 day intervals as necessary to maintain control.

CELERY SEEDBED TREATMENT: Damp Off, Pink Rot. **PEPPERS:** Damp Off - 2 lbs. per 100 gals. spray. Spray or drench at 10 to 15 gals. per 1000 sq. ft. of seed bed. Apply when all seeds have germinated. Repeat as necessary.

CHERRIES: Cherry Leaf Spot (West only), Brown rot, Botrytis rot - 2 lbs. per 100 gals. spray. Spray at prebloom, bloom, petal fall.

DAMP OFF on Cuttings of Azaleas, Carnations and Chrysanthemums: Dip cuttings in a suspension of 4 lbs. per 100 gals. spray before bedding.

GRAPES: Downy Mildew, Blackrot - 2 lbs. in 100 gals. spray. Make first application just before bloom, and repeat immediately after bloom. Depending upon the susceptibility of the grape variety, from 1 to 3 more sprays may be needed at 7 to 10 day intervals. **Dead Arm Disease (California)** - 2 lbs. per 100 gals. spray. 3 to 6 lbs. per acre. First spray at bud break to 1 inch shoot length. Second spray 2 weeks later or shoots 4 to 8 inches in length at 4 to 6 lbs. per acre. **(Northeast)** - 2 lbs. per 100 gals. spray. Make first application when new growth is 4 to 6 inches long, repeat when growth is 8 to 10 inches long.

LEMONS, LIMES, ORANGES, GRAPEFRUIT: Brown Rot (Phytophthora) - 2 to 4 lbs. per 100 gals. spray. Apply as a skirt spray prior to winter rains, thoroughly wetting tree to heights of 3 to 4 feet. Repeat 10 weeks after first application. **ORANGES, TANGELOS:** Scab, Melanose, Tree Response such as increased Fruit Set - Apply 2 lbs. in 100 gallons in post bloom spray from petal fall to full fruit reaches approximately 2 inches. Do not feed treated raw citrus by products to livestock.

LETTUCE, SPINACH: Downy Mildew - 2 lbs. per 100 gals. spray. Apply 100 to 200 gals. of spray per acre. Apply at first signs of disease. Repeat at 7 to 10 day intervals as necessary to maintain control.

Griffin

Captan 50W

50% CAPTAN WETTABLE POWDER

ACTIVE INGREDIENT	BY WT.
*Captan	50%
INERT INGREDIENTS	50%
TOTAL	100%

*N-[[trichloromethylthio]-4-cyclohexene-1,2-dicarboximide]

KEEP OUT OF REACH OF CHILDREN
CAUTION

STATEMENT OF PRACTICAL TREATMENT

If swallowed, drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

If inhaled, remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth to mouth. Get medical attention.

If on skin, wash immediately with soap and water.

If in eyes, flush eyes with plenty of water for at least 15 minutes.

See Side Panel For Additional Precautionary Statements.



GRIFFIN CORPORATION
VALDOSTA, GEORGIA 31601
NET CONTENTS

EPA Reg. No. 1812-261
EPA Est. No. 1812 GA 3

MANGOES

ORNAMENTAL
CHRYSANTHEMUMS

PEACHES, Pears
per 100 gals. spray
cover sprays
on day of harvest

PEARS, Pears
100 gals. spray
TION 1 lb.
100 gals. spray
Pears, Do not

PINEAPPLES

Griffin Captan

Rot Root Rot

plants, immerse
water for 24 hours
height of 15

PRUNES, PLUMS
Prune Russet

POTATOES
plants are 2 to

POTATO SEEDS
water to
with 6 hours
should be stored

POTATO STORAGE
a slurry of 2 lbs.

RASPBERRIES
blossoms are

later. Apply 2 lbs. to 7

starts. Second

STRAWBERRIES
per acre (or 100

starts in spring. Continue the

SOIL AND GRAPE
Rot Diseases

Many Flowering
Apply as a

gals. to each

TARO (Hawaiian)
fin Captan 50W

before taro is
not use if leaf

only

TOMATOES
Spot - 2 lbs.

tion 4 lbs. per
of disease. Repeat

TURF, LAWN
spray. Apply

Apply prior to
areas or feet

PACKING B
Rhizopus B

ing 10 apples
each 8 hours

GRASSES
Copper Spot

per 100 gals.
gal. of water

Repeat at 7
feed clipping

BEANS, CA
to 1/2 bushel

BEETS, CH
per pound of

CONDITION

The Direction

tests. The direction

effectiveness, in

conditions, beyond the

buyer

Griffin Corporation

label and is not

inherent risk

ty of fitness

Corporation

the use or handling

Buyer and User

Griffin Corporation

Griffin Corporation

51

ACTIVE INGREDIENT	BY WT.
*Captan.....	50%
INERT INGREDIENTS.....	50%
TOTAL.....	100%

KEEP OUT OF REACH OF CHILDREN
CAUTION

If swallowed, drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

If on skin, wash immediately with soap and water.

If in eyes: flush eyes with plenty of water for at least 15 minutes.

See Side Panel For Additional Precautionary Statements.



EPA Reg No 1812-267
EPA Est No 1812 GA 3

PACKING BOXES of fruit and vegetable reduction of Storage Rots, Molds due to *Rhizopus*, *Botrytis*, *Cladosporium* - 2 lbs. per 500 gals. Spray, distribute on inside packing in 1" applied as a dip, keep solution agitated continuously. Change dip tank's water after each 8 hour shift.

GRASSES form an entral in nonpastured areas only. Leaf Spot, Damping off, Brown Patch, Copper Spot, Melting Out, Seeding 8, blights and Brown Spot on St. Augustine Grass. 2 lbs. per 100 gals. applying 10 gals. per 1000 sq. ft. (for smaller amounts, mix 8 teabags of herb per gal. of water and apply 1 gal. per 100 sq. ft. of turf). Begin when growth starts in spring. Repeat at 7 to 14 day intervals throughout season. Limit. Do not graze treated areas. Do not feed clippings to livestock.

BEANS, CABBAGE CORN, MELONS PEAS, SQUASH Seed rot, damping off
to 1 least info per bc and dry seed

BEETS, CHARD, SPINACH, GRASS Seed Rot Damping off 1 lb. = 2 bushels
per pound of dry seed. Mix thoroughly in paper bag or glass jar.

CONDITION OF SALE AND WARRANTY

CONDITION OF SALE AND WARRANTY
The Directions for use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherent in the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Griffin Corporation. A buyer shall be assumed by the buyer.

Griffin Corporation warrants that this product conforms to the formal description on the label and is reasonably fit for the purposes referred to in the Directions for Use Subject to the inherent risks referred to above. Griffin Corporation makes no other express or implied warranty. In no case shall Griffin be liable for any other express or implied warranty. In no case shall Griffin be liable for any other damages resulting from the use or handling of this product. Griffin Corporation and the seller accept this product and the Buyer and User accept it subject to the foregoing conditions of Sale and Warranty.

$\mu_0 = 1.4$
 $\mu_1 = 1.4$