



SUL-CO-FLO

6 LB. SULFUR
-plus-

.81 LB. CUPRIC HYDROXIDE

Read the entire label

Do not reuse container. Dispose by burying in areas away from water supply.

CAUTION

This product is toxic to fish. Keep out of lakes, streams, or ponds.

Sulfur may cause injury if used improperly or under unfavorable weather conditions. During periods of high temperature Sulfur may burn foliage.

Do not contaminate water by cleaning of equipment, or disposal of wastes.

Apply this product only as specified on this label.

DO NOT STOP AGITATION WITH SULFUR IN SPRAY EQUIPMENT. SULFUR IS HIGHLY CORROSIVE, AND EQUIPMENT SHOULD BE CLEANED THOROUGHLY AFTER EACH DAYS SPRAYING.



Mark of Quality

RECOMMENDATIONS:

PEANUTS:

To control Rust and Leaf Spot use 2 to 3 quarts Sul-Co-Flo per acre in at least 10 gallons of water to maintain good coverage. Begin applying the above mixture on peanuts 30 to 40 days after planting, when disease first appears. Maintain this schedule each 14 days through the growing season. Do not use within two weeks of an oil spray.

Sulfur is highly corrosive to equipment. After each spraying, clean spray equipment thoroughly.

NOT FOR HOME USE

Parramore & Griffin - Valdosta, Ga.



ACCEPTED

UNDER THE FEDERAL FUNGICIDE ACT FOR THE U.S. DEPT. OF AGRICULTURE

CAUTION:

Keep out of reach of Children

Harmful if swallowed. Avoid breathing spray mist. Contact with skin may cause irritation. Avoid contact with eyes and skin.

ACTIVE INGREDIENT:	
Sulfur as S	50.0%
*Copper (as Metallic)	4.4%
INERT INGREDIENT:	45.6%
TOTAL	100.0%

*Derived from Cupric Hydroxide, 100%

Contains 6.0 pounds Sulfur - .81 pound Cupric Hydroxide per gal.

Net Contents: 5 GALLONS

E.P.A. Reg. No. 1812-215 • E.P.A. EST. No. 100-1000