NOTE: All marked corrections will be made in next printing.

ATRAZINE 80W

ACTIVE INGREDIENT:	
Atrazine (2-chloro-4-ethylamino-6-isopropylamino-s-triazine)	80%
Related compounds	4%
INERT INGREDIENTS	20%

EPA Registration No. 464-481-AA

CAUTION

KEEP OUT OF REACH OF CHILDREN
HARMFUL IF SWALLOWED
AVOID CONTACT WITH EYES, PROLONGED CONTACT WITH SKIN, INHALATION OF DUST, AND CONTAMINATION OF FOOD AND FEED.

Do not contaminate domestic or irrigation water supplies or lakes, streams or ponds.

Do not reuse container. Destroy when empty.

GENERAL INFORMATION

Ground Application

Aerial Application

NOTICE

CORN

Soil Texture	Rate per acre of Atrazine 80W: Broadcast in 20-40 gals. of water per acre	20-inch Band in 10-20 gals. of water per acre	3-inch Band in 7-14 gals. of water per acre

Pre-emergence

Post-emergence

NOTE

Layby Treatment

Ground application

Procedures

NOTE

Preplant

Northern Nutgrass or Yellow Nutsedge (Cyperus esculentus) Control Broadcast 5 lbs. ... preplant, pre-emergence or post-emergence before nutgrass is more than 1½ inches ... applied before nutgrass begins growth. Cultivate with in 2-3 weeks. If applied after ... emerged, cultivate within 10-14 days.

Quackgrass Control on Land Going into Corn Production Split application: Broadcast 2 ... lbs. per acre in the fall or spring and plow 1-3 weeks later. Broadcast a second application at the rate of 2 lbs. per acre in the spring before, during or after planting but before weeds ... inches high. This split application will control both quackgrass and most annual broadleaf and grass weeds.

Single Application Broadcast 3½ lbs. to 5 lbs. per acre in the fall or spring. Plow 1-3 weeks after application.

Use Precautions (Corn) 1) Do not apply more than 5 lbs. Atrazine 80W to corn in any one year.

2) Land treated with Atrazine should not be planted to any crop except corn until the following ... injury may occur.

3) Following harvest of a treated crop, plow (moldboard or disk plow) and thoroughly till the soil in the fall of spring to minimize possible injury to rotational spring seeded crops regardless of the crop used.

4) If Atrazine is applied after June 10 do not rotate with crops other than corn or sorghum the ... year or injury may occur.

5) If Atrazine 80W is used at a broadcast rate higher than 3½ lbs. per acre (or comparable ... band application), a crop of untreated corn or sorghum should precede the next ... crop.

6) ... high P.H. and intermountain areas of the West where rainfall is sparse and erratic ... application is required use Atrazine to control weeds in corn only when corn is no taller ... inches. If untreated corn or sorghum is to precede other rotational crops.

7) In western Minnesota and eastern parts of the Dakotas, Nebraska and Kansas corn treated with Atrazine should not be followed with soybeans if the broadcast rate applied was more ... lbs. per acre (or comparable rate in a band) or injury may occur.

8) Do not plant sugar beets, tobacco, vegetables (including dry beans), spring seeded small ... seeded legumes and grasses the year following Atrazine application or injury may occur.

9) Do not graze treated area or feed treated forage to livestock for 21 days following appli...

TANK-MIX COMBINATION OF ATRAZINE 80W AND SIMAZINE FOR CONTROL OF BOTH BROADLEAF WEEDS AND GRASSES IN CORN

Weeds Controlled Atrazine 80W ... combination with Simazine herbicide controls most ... broadleaf weeds and grasses that compete with corn, including:

annual morningglory, barnyardgrass, Brachiaria spp., bur cucumber, carpetweed, cocklebur, cottonweed, crabgrass, fall panicum, Florida beggarweed, Florida pusley, giant foxtail, green foxtail, jimsonweed, lambsquarters, nightshade, northern nutgrass (yellow nutsedge), pigweed, prickly sida (teaweed), purslane, ragweed, smartweed, velvetleaf (buttonweed), wild cucumber, wild mustard, wild oats, witchgrass, yellow foxtail.

Weeds Not Controlled Atrazine + Simazine does not control perennial weeds such as milkweed, bindweeds and perennial vines.

When to Apply Apply ... before the weeds ... up. However, the following recommendations ... prior to the most effective weed control.

Preplant application Broadcast Atrazine + Simazine ... the spring after plowing during ... other soil surface preparation. Apply ... before planting within 2 weeks of planting.

... Atrazine + Simazine ... that will not ... effectiveness ... will remain on the soil surface with ... moves it down into the soil ...

... the necessity to incorporate ... the Atrazine + Simazine into the ... weeds or grass roots before the dry treated surface ... is ... weather ... prior to the herbicide in the weed germinating zone.

Preemergence application Apply ... after planting, before either ... apply preemergence ... Simazine 5 may ... not to ... germination ...

How to Apply ... your Atrazine + Simazine ... with ... the ... to ... the ... Atrazine 80W + Simazine 80W ... of the ... the best ... to the ... without ... And ... plot ...

... with ... Simazine and ... may ... the ... the ...

each ... as ...

86-1298 PRINTED IN U.S.A. IN JULY, 1973.

THIS IS THE INITIAL PRINTING OF THIS SPECIMEN LABEL.

1812-239

AC

BEST DOCUMENT AVAILABLE