

SUPER-TIN 4L

PRECAUTIONARY STATEMENTS HAZARDOUS TO HUMANS & DOMESTIC ANIMALS

DANGER

Spray may be fatal if swallowed or absorbed through the skin. May cause skin irritation. Do not allow contact of spray with skin or eyes, wear goggles or face shield when handling. In case of skin contact, wash immediately with soap and water. For eye contact, flush with water and get medical attention immediately. Do not breathe spray mist. Spray is hazardous if inhaled. Avoid contamination of food stuffs and feed. In case of ingestion, induce vomiting. Immediately consult a physician. Keep out of reach of children.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Pesticide, spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Dispose of in an incinerator or landfill approved for pesticide containers or bury in a safe place. Consult Federal, State or Local Disposal Authorities for approved alternative procedures such as limited open burning.

GRIFFIN SUPER TIN Fungicide can be applied as a ground or aerial spray to control fungicidal infestations on listed crops. Application rates are for general use. The state agricultural extension or agricultural experiment station specialists should be consulted for specific applications and timing recommendations. With any spray application thorough coverage is essential for good control.

Apply this product only as specified on this label.

PRECAUTIONS:

GRIFFIN SUPER TIN Fungicide can be applied in combination with GRIFFIN SUPER S-X for additional fungicidal control. Do not add surfactants, spreaders or stickers. Do not graze dairy or meat animals in treated areas.

GENERAL INSTRUCTIONS FOR APPLICATION

SUGAR BEETS: Cercospora Leafspot

Ground application, apply 3.8 - 9.5 fl. ounces of GRIFFIN SUPER-TIN per acre in at least 15 gallons of water. Full coverage of the foliage is necessary for best results. **Aerial application,** (helicopter or fixed wing aircraft) apply 3.8 - 9.5 fl. ounces of SUPER-TIN per acre in 5 - 10 gallons of water. Diluted spray should be directed uniformly to all parts of the plant. Use lower gallonage when plants are small and increase volume with plant size. Use the lower rate for protective sprays and the higher rates later in the season or during high infection periods. Application should begin when leafspot conditions appear or when the disease is in the area and repeated at 10 to 14 day intervals. Do not treat within 14 days of harvest. Do not graze or feed beet tops to livestock.



ACTIVE INGREDIENT:

Triphenyltin hydroxide 40.0%

INERT INGREDIENTS: 60.0%

TOTAL 100.0%

Contains 4 pounds Triphenyltin hydroxide per gallon.

KEEP OUT OF REACH OF CHILDREN

DANGER

If on skin, wash immediately with soap and water.

If in eyes, flush with water and get medical attention.

If ingested, induce vomiting — use Syrup of Ipecac if available. Do not use salt. Consult Poison Control Center for assistance and see physician immediately.

See Side Panel For Additional Precautionary Statements.

EPA REG. NO. 1812-244

EPA EST. NO. 1812-GA-3

GRIFFIN CORPORATION

VALDOSTA, GEORGIA 31601

NET CONTENTS 1 GALLON

POTATOES: Early and Late Blight

Ground application, apply 4.7 - 9.5 fl. ounces of SUPER-TIN per acre in at least 15 gallons of water. Full coverage of the foliage is necessary for best results. A spray pressure of less than 200 pounds is recommended. **Aerial application,** apply 4.7 - 9.5 fl. ounces per acre in 3 - 10 gallons of water. For helicopter application, fly high enough so as not to whip the vines. Diluted spray should be directed uniformly to all parts of the plant and the gallonage increased according to the size of the plants. Application should begin with the appearance of blight weather conditions and continue on a 7 day schedule. Do not treat within 7 days of harvest when applied by ground or aerial application equipment. The lower rate of application should be used early in the season and the high rate mid to late season or when blight infection is in the area.

PEANUTS: Cercospora Leafspot

Ground application, apply 4.7 - 7.6 fl. ounces per acre in 10 - 100 gallons of water. **Aerial application,** apply 4.7 - 7.6 fl. ounces per acre in 3 - 10 gallons of water. Use lower rate when leafspot is light, increasing as disease pressure increases.

Apply in sufficient water for uniform, full coverage of foliage. Use lower gallonage in early season when plants are small and increase spray volume as plant size increases. Application should begin approximately six (6) weeks after planting or as soon as first sign of leafspot appear. Spray application should continue on a 10 - 14 day schedule. Do not apply within 14 days of harvest. Do not allow hogs to feed on peanuts in treated fields. Hulls from treated peanuts may be used in feed for livestock. Do not use vines for feed.

CARROTS: Cercospora Leafspot and Alternaria Blight

Apply 3.8 - 7.6 fl. ounces per acre. Use lower rate when diseases are light, increasing rate when disease pressure increases. Apply in sufficient water for uniform, full coverage of foliage 25 - 100 gallons per acre. Use lower volumes in early season when plants are small and increase spray volume as plant size increases. Applications should begin 6 weeks after planting or as soon as first signs of disease appear. Spray applications should continue at 7 day intervals or as needed for continued control. Do not apply within 14 days of harvest. Do not use tops as feed for livestock.

TOBACCO: Brown Spot (Alternaria longipes)

Mix 15.2 fl. ounces of SUPER-TIN per 100 gallons of water and apply 40 - 60 gallons per acre (6.1 - 9 fl. ounces of SUPER-TIN per acre) as a full coverage spray. Diluted spray should be directed to all parts of the plant. Application should begin immediately after first priming (harvest) and repeated immediately after each weekly priming.

PECANS: Scab, Brown Leafspot, Powdery Mildew, Liver Spot, Sooty Mold and Leaf Blotch

Aerial application, apply 11.4 - 22.8 fl. ounces per acre in a minimum 20 gallons of water. **Ground application,** mix 3 - 6 fl. ounces of SUPER-TIN for each 100 gallons of water 15.2 - 45.6 fl. ounces per 100 gallons tank and apply as a full coverage spray. Diluted spray should be directed to all parts of the tree. Applications should begin at pre-pollination stages when the young leaves are unfolding, and a second application made when the small nuts are forming. Repeat spray applications at 2 - 4 week intervals as needed to maintain control. Use the lower rate for the first two applications or until the disease becomes severe, or during dry weather. Use the higher rate during wet weather or during severe scab, powdery mildew, or other disease infections. Do not apply after shucks have started to open.

WARRANTY STATEMENT

Seller makes no warranty, express or implied, concerning the use of this product other than indicated on the label. This material when such use and handling is in accordance with the label instructions.

Seller makes no warranty, express or implied, concerning the use of this product other than indicated on the label. This material when such use and handling is in accordance with the label instructions.