


ACCEPTED
with COMMENTS
in EPA Letter Dated

JUL 11 1984

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
1512-244



Triphenyltin Hydroxide Flowable Fungicide

BEST DOCUMENT AVAILABLE

ACTIVE INGREDIENT:		
Triphenyltin hydroxide	40.0%
INERT INGREDIENTS:	60.0%
TOTAL	100.0%

Contains 4 pounds Triphenyltin hydroxide per gallon

**KEEP OUT OF REACH OF CHILDREN
DANGER**

If on skin, wash immediately with soap and water.
If in eyes, flush with water and get medical attention.
If swallowed, drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth an unconscious person. Get medical attention.
See Side Panel For Additional Precautionary Statements

Griffin

GRIFFIN CORPORATION
VALDOSTA, GEORGIA 31601
NET CONTENTS 1 Gallon

EPA Reg. No. 1512-244
EPA Est. No. 1812 GA 3

GENERAL INSTRUCTIONS FOR APPLICATION

SUGAR BEETS: Cercospora Leafspot
Ground application, apply 3.8 - 9.5 fl. ounces of SUPER-TIN 4L™ per acre in at least 15 gallons of water. Full coverage of the foliage is necessary for best results. Aerial application, (helicopter or fixed wing aircraft) apply 3.8 - 9.5 fl. ounces of SUPER-TIN 4L™ per acre in 5 - 10 gallons of water. Diluted spray should be directed uniformly to all parts of the plant. Use lower gallonage when plants are small and increase volume with plant size. Use the lower rate for protective sprays and the higher rates later in the season or during high infection periods. Application should begin when leafspot conditions appear or when the disease is in the area and repeated at 10 to 14 day intervals. Do not treat within 14 days of harvest. Do not graze or feed beet tops to livestock.

POTATOES: Early and Late Blight, Suppression of Colorado Potato Beetle
Ground application, apply 4.7 - 9.5 fl. ounces of SUPER-TIN 4L™ per acre in at least 15 gallons of water. Full coverage of the foliage is necessary for best results. A spray pressure of less than 200 pounds is recommended. Aerial application, apply 4.7 - 9.5 fl. ounces per acre in 3 - 10 gallons of water. For helicopter application, fly high enough so as not to whip the vines. Diluted spray should be directed uniformly to all parts of the plant and the gallonage increased according to the size of the plants. Application should begin with the appearance of blight weather conditions and continue on a 7 day schedule. Do not treat within 7 days of harvest when applied by ground or aerial application equipment. The lower rate of application should be used early in the season and the high rate mid to late season or when blight infection is in the area. IRRIGATION SYSTEM APPLICATION (solid-set or center-pivot system only) — Apply 5.5 to 9.5 fl. oz. per acre. Do not apply through center-pivot (circle) system which uses water pressure for movement of system. Irrigation system must be equipped with safety valves or other devices to prevent back-siphoning of pesticide into water source. Any irrigation water treated with this product should be held on the treated area until it is absorbed by the soil. It should not be turned into the tail water or fed back into the irrigation system. For complete directions regarding application, refer below. Application should begin with the appearance of blight weather conditions and continue on a 7 day schedule. Do not treat within 7 days of harvest when applied by ground or aerial application equipment and not within 33 days when applied by irrigation system. The lower rate of application should be used early in the season and the high rate mid to late season or when blight infection is in the area.

CARROTS: Cercospora Leafspot and Alternaria Blight
Apply 3.8 - 7.6 fl. ounces per acre. Use lower rate when diseases are light, increasing rate when disease pressure increases. Apply in sufficient water for uniform, full coverage of foliage 25 - 100 gallons per acre. Use lower volumes in early season when plants are small and increase spray volume as plant size increases. Applications should begin 6 weeks after planting or as soon as first signs of disease appear. Spray applications should continue at 7 day intervals or as needed for continued control. Do not apply within 14 days of harvest. Do not use tops as feed for livestock.

PECANS: Scab, Brown Leafspot, Downy Spot, Powdery Mildew, Liver Spot, Sooty Mold and Leaf Blotch
Aerial application, apply 11.4 - 22.8 fl. ounces per acre in a minimum 20 gallons of water. Ground application, mix 3.9 fl. ounces of SUPER-TIN 4L™ for each 100 gallons of water 15.2 - 45.6 fl. ounces per 500 gallons tank and apply as a full coverage spray. Diluted spray should be directed to all parts of the tree. Applications should begin at pre-pollination stages when the young leaves are unfolding, and a second application made when the small nuts are forming. Repeat spray applications at 2 - 4 week intervals as needed to maintain control. Use the lower rate for the first two applications or until the disease becomes severe, or during dry weather. Use the higher rate during wet weather or during severe scab, powdery mildew or other disease infections. Do not apply after shucks have started to open.

WARRANTY STATEMENT

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

GNC 01280
CPK 05A15



$\frac{346}{17}$ $\frac{121884}{4}$
 $\frac{300}{37}$ $\frac{120098}{5}$
 $\frac{301}{17}$ $\frac{121413}{5}$

JUL 11 1984

Griffin Corporation
 Rocky Ford Rd.
 Valdosta, GA 31601

Attention: L. Elliott

Gentlemen:

Subject: Super Tin 4L
 EPA Registration No. 1812-244
 Your Request for Alternate Formulation Dated
 April 24, 1984 and Amended Label Dated
 April 5, 1984.

There is no objection to the use of the alternate formulation requested in your letter of the above date. The amendment has been inserted in the file for the subject product.

The addition of "Suppression of Colorado Potato Beetle" on the label is acceptable. Submit 5 copies of the final printed label and include:

1. In the Environmental Hazard Statement change "Keep...water to" "Do not apply directly to water or wetlands."
2. Comply with PR Notice 83-3.

Find enclosed a stamped copy of the label for your files.

Sincerely yours,

Henry M. Jacoby
 Product Manager (21)
 Fungicide-Herbicide Branch
 Registration Division (TS-767)

Enclosure

RD:Jacobyl:DCR-17092:WANG-0062K:mar:Raven:479-2013:6/22/84

BEST DOCUMENT AVAILABLE

CONCURRENCES							
SYMBOL ▶							
SURNAME ▶							
DATE ▶							