

MAR 06 1995

Mr. Richard H. Collier
Griffin Corporation
P.O. Box 1847
Valdosta, Georgia 31603-1847

Dear Mr. Collier:

Subject: Manex
EPA Reg. No. 1812-251
Your Submission of February 6, 1995

The amendment referred to above, submitted in connection with registration under FIFRA sec. 3(c)(7)(A), is acceptable provided that you:

1. Submit and/or cite all data required for registration and reregistration under FIFRA sec. 3(c)(5) and sec. 4 when the Agency requires all registrants of similar products to submit such data.

2. You may add, under potatoes, the states of Delaware, Michigan, Ohio and Rhode Island to the states with a 3-day PHI.

3. Submit five (5) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

Connie B. Welch
Connie B. Welch
Acting Product Manager (21)
Fungicide-Herbicide Branch
Registration Division (7505C)

Enclosure

7505C:C.Grable:cg:2/27/95

Manex®

Fungicide

Flowable with Zinc

ACCEPTED
with COMMENTS
In EPA Letter Dated

MAR 06 1995

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

1812-251

ACTIVE INGREDIENT:

MANEB (MANGANESE ETHYLENEBISDITHIOCARBAMATE).....37%

(Total manganese as metallic.....7.6%)

INERT INGREDIENTS:.....63%

TOTAL

100%

4 Pounds of Maneb Per Gallon

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

IF INHALED: Remove victim to fresh air; if not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

IF ON SKIN: Remove by washing.

IF IN EYES: Flush with plenty of water. Call a physician.

GRIFFIN CORPORATION
VALDOSTA, GEORGIA 31601

EPA REG. NO. 1812-251
EPA EST. NO. _____

NET CONTENTS

GALLONS

15

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS (AND DOMESTIC ANIMALS)

CAUTION

Precautionary Statements: May cause irritation of eyes, nose, throat and skin. Avoid breathing spray mist. Do not rub eyes or mouth with hands.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Drift or runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Cover or incorporate spilled treated seed. Do not contaminate water when disposing of equipment washwaters.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers (other than mixing and loading) must wear:

- coveralls over long-sleeved shirt and long pants
- waterproof gloves
- shoes plus socks

Mixers and Loaders must wear:

- coverall over long-sleeved shirt and long pants
- waterproof gloves
- shoes plus socks
- protective eyewear
- chemical-resistant apron when mixing and loading

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering controls statements:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240 (d)(4-6), the handler PPE requirements may be reduced or modified as specified in the WPS.

The closed systems or enclosed cabs must be used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240 (d)(4-6). The handler PPE requirements may be reduced or modified as specified in the WPS.

Human flaggers must be in enclosed cabs.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

~~This product is toxic to fish. Drift or runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Cover or incorporate spilled treated seed. Do not contaminate water when disposing of equipment washwaters.~~

PHYSICAL AND CHEMICAL HAZARDS

~~Keep away from fire and sparks. Store in a cool dry place.~~

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- coveralls over long-sleeved shirt and long pants
- waterproof gloves
- shoes plus socks

In case of accidental exposure see Statement of Practical Treatment:

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides 40 CFR part 170.

The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

STORAGE AND DISPOSAL

Store in a cool dry place. Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities by burning. If burned, stay out of smoke.

PHYSICAL AND CHEMICAL HAZARDS

Keep away from fire and sparks. Store in a cool dry place.

APPLICATION DIRECTIONS FOR USE AS A SPRAY

Manex is approved for application in sufficient water to provide good coverage with available equipment in either dilute sprays or in concentrate ground or aerial sprays. Rates listed are based on 100 gallons of dilute spray unless otherwise noted. Rates of product for concentrate and aerial applications should be equal to the rates for dilute sprays on a per acre basis.

FOR GROUND APPLICATIONS use at the rate indicated in sufficient water for thorough coverage, minimum of 10 gallons. Increased volume of water may be necessary as foliage density increases.

FOR AERIAL APPLICATIONS use at the rate indicated in sufficient water for thorough coverage, a minimum of 3 gallons per acre for field crops and 10 gallons per acre for orchard crops. Apply with properly calibrated aerial equipment, arrange nozzles so that spray delivery is uniform over the entire spray swath

When dosage ranges are given, use the higher rate and shorter intervals under severe disease pressure, but do not exceed the maximum rate or apply more frequently than the minimum interval given in the directions for that crop.

FOLIAR APPLICATIONS

Where EBDC Products Used Allow the Same Maximum Poundage of Active Ingredients Per Acre Per Season:

If more than one product containing an EBDC active ingredient (maneb, mancozeb, or metiram) is used on a crop during the same growing season and the EBDC products used allow the same maximum poundage of active ingredient per acre per season, then the total poundage of all such EBDC products used must not exceed any one of the specified individual EBDC product maximum seasonal poundage of active ingredient allowed per acre.

Where EBDC Products Used Allow Different Maximum Poundage of Active Ingredients Per Acre Per Season:

If more than one product containing an EBDC active ingredient is used on a crop during the same growing season and the EBDC products used allow different maximum poundage of active ingredient per acre per season, then the total poundage of all such EBDC products used must not exceed the lowest specified individual EBDC product maximum seasonal poundage of active ingredient allowed per acre.

SEED TREATMENT

In addition to the maximum number of foliar applications permitted by the formula stated above, a single application of seed treatment may be made on crops which have registered seed treatment uses.

CHEMIGATION

Apply Manex only through sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set or hand move. Do not apply Manex through any other type of irrigation system.

Crop injury, lack of effectiveness or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.

If you have questions about calibration you should contact State Extension Service Specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label prescribed safety devised for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Specific Instructions for Public Water Systems

Public water systems means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone backflow preventor (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public waier system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top of the overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent the water fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injections pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

Specific Instructions for Sprinkler Irrigation Systems

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quickcheck valve to prevent the flow of liquid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

7.4.72

The irrigation line or water pump must include a functional pressure switch which will stop the water pump when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

Good agitation is required in the injection tank. In moving systems, apply specified dosage of Manex as a continuous injection. In nonmoving systems inject Manex for 15 to 30 minutes at end of cycle. Use the least amount of water possible with uniform coverage.

Mix the amount of Manex needed for acreage to be treated into the quantity of water determined during prior calibration. For moving systems inject into the system continuously for one complete revolution of the field. For nonmoving systems inject into system for the time established during calibration.

Stop injection equipment after treatment is completed and continue to operate irrigation equipment until all Manex is flushed from system.

APPLICATION INSTRUCTIONS FOR APPLICATION

FRUIT AND NUT CROPS

CROPS	DISEASES	QUARTS PER ACRE	MAXIMUM QUARTS PER ACRE PER SEASON	USE INSTRUCTIONS
Almonds	Brown Rot Blossom and Twig Blight, Fungus (Leaf Blight), Scab, Shothole	4.8 - 6.4 or (1.2 - 1.6 per 100 gal in a dilute spray)	25.6	Apply in popcorn, full bloom and petal fall or every 7 to 10 days if bloom is staggered. Omit petal fall spray if only brown rot is present. Do not apply later than 5 weeks after petal fall or within 145 days of harvest. Do not allow livestock to graze in almond orchards.

Apples

Use either the Pre-bloom or Extended Application schedules. DO NOT COMBINE OR INTEGRATE THE TWO TREATMENT SCHEDULES.

Rusts (including Cedar apple), Fabrea Leaf Spot, Flyspeck, Scab, Sooty Blotch 4.8 or (1.2 per 100 gal in dilute spray) 19.2 per year

Pre-bloom: Begin applications at 1/4 to 1/2 inch green tip and continue on a 7 to 10 day application schedule through bloom. Do not graze livestock in treated areas. It is recommended that this product be used in an Integrated Pest Management Program.

2.4 or (0.6 per 100 gal in dilute spray) 16.8 per year

Extended application or tank mix: For implementation of IPM programs, applications based on tree-row volume or for use as a resistance management tool: begin applications at 1/4 to 1/2 inch green tip and continue applications on a 7 to 10 day schedule through the second cover spray or to within 77 days of harvest. Do not apply within 77 days of harvest. Do not graze livestock in treated areas. It is recommended that this product be used in an Integrated Pest Management Program.

Add spreader/sticker to spray mixture for cover sprays and tank mix with an effective systemic protectant/curative fungicide for more effective control of diseases.

Bananas Sigatoka 1.6 - 2.4 24

Apply when leaves first appear and repeat as needed on a 14 to 21 day interval. May be applied up to the day of harvest.

10-21-02

Cranberries	Fruit Rot	2.43.8 - 4.8	14.4	Spray on a 7 to 10 day interval. Do not apply within 30 days of harvest.
Grapes	Black Rot, Bunch Rot, Deadarm, Downy Mildew	1.2 - 3.2 or (0.3 - 0.8 per 100 gal in a dilute spray)	19.2	East of the Rocky Mountains: Apply in sufficient water to provide thorough coverage starting when new shoots are 1/2 to 1 1/2 inches long and continue at 7- to 10 day intervals until fruit is set. Do not apply within 66 days of harvest.
		1.2 - 2.0 or (0.3 - 0.5 per 100 gal in a dilute spray)	6.0	West of the Rocky Mountains: Apply when shoots are 1/2 to 1 1/2 inches long and continue at 7 to 10 day intervals. Do not apply within 66 days of harvest except in California. California: Do not apply after bloom.
Kadota Figs	Surface Mold (Cladosporium), Surface Rot (Alternaria)	0.6 per 100 gal	2.4	Apply when disease threatens in 100-400 gallons of water per acre. Do not apply with in 10 days of harvest.
Papayas	Anthrachnose, Phytophthora, Fruit Rot, Cercospora Black Rot	1.6 - 2.0 or (0.4 - 0.5 per 100 gal in a dilute spray)	28	Apply when disease first threatens and repeat at 14 to 21 day intervals. May be applied up to the day of harvest.

1-

FIELD AND VEGETABLE CROPS

CROPS	DISEASES	QUARTS PER ACRE	MAXIMUM QUARTS PER ACRE PER SEASON	USE INSTRUCTIONS
Beans (Dry)	Anthracnose, Downy Mildew, Rust	1.2 - 1.6	9.6	Begin when plants are small. Spray on a 5 to 7 day interval. Do not apply within 30 days of harvest.
Broccoli, Brussels Sprout, Cauliflower	Alternaria Leaf Spot, Downy Mildew	1.2 - 1.6	9.6	Begin when diseases threaten. Spray on a 7 to 10 day interval. Do not apply within 7 days of harvest.
Cabbage, Kohlrabi, Chinese Cabbage (tight headed only)	Alternaria Leaf Spot, Downy Mildew	1.2 - 1.6	9.6	Plant beds and direct seeded fields: Spray on a 7 to 10 day interval. Do not apply within 7 days of harvest.
Chinese Cabbage (loose head)	Alternaria Leaf Spot, Downy Mildew	0.8 - 1.2	7.2	Plant beds and direct seeded fields: Spray on a 7 to 10 day interval. California: Do not apply within 7 days of harvest. Hawaii: Do not apply within 10 days of harvest.
Corn (sweet corn, popcorn, sweet corn used for seed production)	Common Rust, Helminthosporium Blight	1.2	18	East of the Mississippi including AR and LA: Use sufficient water for thorough coverage. Start applications when disease first appears and repeat at 3 to 10 day intervals. Add a recommended surfactant or spreader/sticker if needed for better coverage. Do not apply within 7 days of harvest. Do not feed treated forage to livestock.

10/15

		1.2	6	West of the Mississippi except AR and LA: Use sufficient water for thorough coverage. Start applications when disease first appears at 3 to 10 day intervals. Add a recommended surfactant or spreader/sticker if needed for better coverage. Do not apply within 7 days of harvest. Do not feed treated forage to livestock.
Cucumbers	Alternaria (Macrosporium) Leaf Spot, Angular Leaf Spot, Anthracnose, Downy Mildew, Pythium Fruit Rot	1.2 - 1.6	12.8	Begin when diseases threaten or plants begin to run. Spray on a 7 to 10 day interval. Do not apply within 5 days of harvest.
Eggplant	Anthracnose, Septoria Leaf Spot, Cladosporium Leaf Mold, Early Blight, Late Blight, Gray Leaf Spot (Stemphylium)	1.2 - 1.6	11.2 (per crop)	Begin at first fruit cluster and repeat at 7 to 10 day intervals. Do not apply within 5 days of harvest.
Kale	Alternaria Leaf Spot, Downy Mildew	1.2 - 1.6	3.2 (per cutting)	Begin when disease threatens and apply on a 7 to 10 day interval. Do not apply with 10 days of harvest.
Lettuce (Head and Leaf), Endive	Downy Mildew	1.2 - 1.6	9.6	Apply when disease appears. Spray on a 7 to 10 day interval. Remove residues from head lettuce by stripping and trimming. Do not apply within 10 days of harvest.
			6.4	California: Spray on a 7 to 10 day interval. Do not apply within 14 days of harvest.

200

Melons (Cantaloupe, Casabas, Crenshaws, Honeydew, Watermelons)	Alternaria Leaf Spot, Anthracnose, Cercospora Leaf Spot, Downy Mildew, Gummy Stem Blight	1.2 - 1.6	12.8	Apply as soon as plants begin to run or when disease first appears. Spray on a 7 to 10 day interval. Do not apply within 5 days of harvest.
Onions (Dry Bulb)	Botrytis Leaf Blight, Downy Mildew, Neck Rot, Purple Blotch	1.6 - 2.4	24	Begin applications when diseases are first reported in the area at 7 day intervals. Do not apply within 7 days of harvest. Do not apply to exposed bulbs.
Onions (Green)	Botrytis Leaf Blight, Downy Mildew, Neck Rot, Purple Blotch	1.6 - 2.4	16.8	Begin application when diseases are first reported in the area at 7 day intervals throughout the season. Do not apply within 7 days of harvest. Do not apply to exposed bulbs.
	Smut (except in California)	2.4	2.4 (per 29,000 linear feet of row - 18 inch spacing)	Apply as a furrow drench at time of planting onion seeds. Use 75 to 100 gallons of water/acre.
Peppers	Anthracnose, Cercospora Leaf Spot (Frogeye Spot), Phytophthora Blight, Ripe Rot	1.2 - 2.4	14.4	East of the Mississippi: Begin when disease threatens. Spray on a 7 to 10 day interval. Do not apply within 7 days of harvest.
		1.2 - 1.6	9.6	West of the Mississippi: Begin when disease threatens. Spray on a 7 to 10 day interval. Do not apply within 7 days of harvest.

1-12-20

Potatoes	Early and Late Blights	0.8-1.2 - 1.6	11.2	Begin applications when plants are 2 to 6 inches high by applying 0.8 qts. pre acre. As the vines increase in size, apply 1.2 to 1.6 qts. per acre. Apply on a 5 to 10 day interval. Do not apply within 14 days of harvest except for FL, CT, ME, MA, NH, NY, PA, VT and WI where the PHI is 3 days. It is recommended that this product be used with an Integrated Pest Management Program. Vine Kill should occur 14 days before harvest.
Potatoes (Seed Pieces)	Fusarium Seed Piece Decay	0.8 quart/10 gallons		Dip whole or cut tubers. Spread in cool place if held before planting. Seed piece treatment only. Do not use treated seed pieces for food, feed or oil purposes.
Pumpkins	Angular Leaf Spot, Downy Mildew	1.2 - 1.6	12.8	Begin when disease threatens. Spray on a 7 to 10 day interval. Do not apply within 5 days of harvest.
Squash (Winter and Summer)	Anthracoise, Downy Mildew	1.2 - 1.6	12.8	Begin when plants start to run. Spray on a 7 to 10 day interval. Do not apply within 5 days of harvest.
Sugar Beets	Cercospora Leaf Spot	1.2 - 1.6	11.2	Apply when disease appears. Spray on a 7 to 10 day interval. Do not apply within 14 days of harvest.
Tomatoes (Greenhouse and Field)	Anthraenose, Cladosporium Leaf Mold, Early and Late Blights, Gray Leaf Spot (Stemphyllum), Septoria Leaf Spot	1.2 - 2.4	16.8	East of the Mississippi: Begin at first fruit cluster and repeat at 7 to 10 day intervals. Do not apply within 5 days of harvest.

15.75

1.2 - 1.6 6.4

West of the Mississippi: Begin at first fruit cluster and repeat at 7 to 10 day intervals. Do not apply within 5 days of harvest.

ORNAMENTALS

CROP	DISEASES	RATE PER ACRE IN 100 GAL WATER	USE INSTRUCTIONS
Carnations	Alternaria Leaf Spot, Anthracnose, Botrytis Blight	0.8 - 1.2 quarts	Begin when new growth starts. Repeat weekly.
Chrysanthemums	Ascochyta Ray Blight, Botrytis Petal Spot, Septoria Leaf Spot	12.8 fl oz plus 3/4 pound Captan 50W	Apply twice weekly during the blooming season for petal spot and ray blight. For septoria leaf spot apply weekly throughout the season.
Dahlias	Alternaria Leaf Spot, Botrytis Blight	0.8 - 1.2 quarts	Begin when new growth starts. Repeat weekly.
Dogwood	Anthracnose	0.8 - 1.2 quarts	Begin when buds open. Repeat when bracts fall 4 weeks later and in late summer.
Gladiolus	Botrytis Blight, Curvularia and Stemphylium Leaf Spots	1.2 quarts	Begin when flower spikes are developing. Repeat 2 to 3 times at weekly intervals.
Lilies	Botrytis Blight	0.8 - 1.2 quarts	Begin with new growth. Repeat weekly.
Pansies	Anthracnose	0.8 - 1.2 quarts	Begin with new growth. Repeat weekly.

Peonies	Alternaria Leaf Spot, Botrytis Blight, Phytophthora Blight	0.8 - 1.2 quarts	Apply to foliage and soil in early spring and early fall and 7 to 10 day intervals during the growing season.
Roses	Black Spot, Cercospora Leaf Spot (Texas), Rust (California)	0.8 - 1.2 quarts	Begin when first leaves unfold. Repeat at 7 to 10 day intervals.
Snapdragons	Rust	0.8 - 1.2 quarts	Begin with emergence. Repeat weekly.
Zinnias	Alternaria Blight, Leaf Spot, Botrytis Blight	0.8 - 1.2 quarts	Begin with emergence. Repeat weekly.

TURF GRASSES

CROP	DISEASES	USE RATE	USE INSTRUCTIONS
Grasses	Brown Patch	4.8 fl oz in sufficient water/1000 square feet	Begin at first sign of disease. Repeat at 7 to 14 day intervals. Do not use on pasture or range grasses. Do not feed clippings to livestock. Do not use if grass is grown for seed.
	Dollar Spot	9.6 - 12.8 fl oz in sufficient water/1000 square feet	Begin at first sign of disease. Repeat at 7 to 14 day intervals. Do not use on pasture or range grasses. Do not feed clippings to livestock. Do not use if grass is grown for seed.

Melting-Out	4.8 - 6.4 fl oz in sufficient water/1000 square feet	Begin at first sign of disease. Repeat at 7 to 14 day intervals. Do not use on pasture or range grasses. Do not feed clippings to livestock. Do not use if grass is grown for seed.
Leaf, Stem and Stripe Rusts	2.4 quarts per acre in 100 gal water	Begin when rust pustules are first seen. Repeat at 7 to 14 day intervals. Do not graze treated areas or feed clippings to livestock.

Our recommendations for use are based on tests believed reliable. Since the use is beyond our control, we can not guarantee the results if such use is not in accordance with directions. We disclaim any responsibility for damages resulting from careless or improper handling or use.

WARRANTY STATEMENT

GRIFFIN warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of GRIFFIN. In no case shall GRIFFIN be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. GRIFFIN MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESSED OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

Manex[®] is a trademark of Griffin Corporation

MAR 06 1995

Mr. Richard H. Collier
Griffin Corporation
P.O. Box 1847
Valdosta, Georgia 31603-1847

Dear Mr. Collier:

Subject: Manex Fungicide
EPA Reg. No. 1812-251
Your Submission of November 21, 1994

The amendment referred to above, submitted in connection with registration under FIFRA sec. 3(c)(7)(A), is acceptable provided that you:

1. Submit and/or cite all data required for registration and reregistration under FIFRA sec. 3(c)(5) and sec. 4 when the Agency requires all registrants of similar products to submit such data.

2. Make the labeling changes listed below before you release the product for shipment:

a. The maximum number of applications per season for several crops exceeds the maximum allowed by the Special Review PD4 for EBDC containing products. Revise the maximum number of applications to the following:

Almonds; 4
Apples, prebloom; 4
Apples, extended; 7
Grapes, east of Rockies; 6
Grapes, west of Rockies; 3


b. You may add, under potatoes, the states of Delaware, Michigan, Ohio and Rhode Island to the list of states with a 3-day PHI.

3. Submit five (5) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,



Connie B. Welch
Acting Product Manager (21)
Fungicide-Herbicide Branch
Registration Division (7505C)

Enclosure

) 7505C:C.Grable:cg:2/27/95

12/22/93 11/21/94

HOMEOWNER

Manex™ FUNGICIDE

Flowable with Zinc

ACCEPTED
with COMMENTS
to EPA Letter Dated

MAR 06 1995

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

1812-251

ACTIVE INGREDIENT:

MANEB (MANGANESE ETHYLENEBISDITHIOCARBAMATE).....37%

(Total manganese as metallic.....7.6%)

INERT INGREDIENTS:.....63%

) TOTAL 100%

4 Pounds of Maneb Per Gallon

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

IF INHALED: Remove victim to fresh air; if not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

IF ON SKIN: Remove by washing.

IF IN EYES: Flush with plenty of water. Call a physician.

GRIFFIN CORPORATION
VALDOSTA, GEORGIA 31601

EPA REG. NO. 1812-251
EPA EST. NO. 1812-4A33

NET CONTENTS

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS (AND DOMESTIC ANIMALS)

CAUTION

Precautionary Statements: May cause irritation of eyes, nose, throat and skin. Avoid breathing spray mist. ~~After sprays have dried, do not enter or allow entry into treated areas where there is danger of drift until the 24-hour re-entry interval has expired unless wearing the personal protective equipment listed on the label.~~ Do not enter treated areas until sprays have dried. Keep all unprotected persons, children, livestock, and pets away from treated area or where there is danger of drift. Do not rub eyes or mouth with hands.

) Home gardeners applying this product must wear long-sleeved shirt, long pants and chemical resistant gloves. The gloves must be washed thoroughly with soap and water before removing. Clothes must be changed immediately after use of this product and must be laundered separately from other laundry items before reuse.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Drift or runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Cover or incorporate spilled treated seed. Do not contaminate water when disposing of equipment washwaters.

PHYSICAL AND CHEMICAL HAZARDS

) Keep away from fire and sparks. Store in a cool dry place.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers (other than mixing and loading) must wear:

- coveralls over long-sleeved shirt and long pants
- waterproof gloves
- shoes plus socks

Mixers and Loaders must wear:

- coverall over long-sleeved shirt and long pants
- waterproof gloves
- shoes plus socks
- protective eyewear
- chemical resistant apron when mixing and loading

~~Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.~~

~~USER SAFETY RECOMMENDATIONS~~

~~Users should:~~

- ~~—Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.~~
- ~~—Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.~~
- ~~—Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.~~

~~ENVIRONMENTAL HAZARDS~~

~~This product is toxic to fish. Drift or runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Cover or incorporate spilled treated seed. Do not contaminate water when disposing of equipment washwaters.~~

~~PHYSICAL AND CHEMICAL HAZARDS~~

~~Keep away from fire and sparks. Store in a cool dry place.~~

~~DIRECTIONS FOR USE~~

~~It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.~~

~~RE-ENTRY STATEMENT~~

~~Do not enter treated areas until sprays have dried.~~

~~STORAGE AND DISPOSAL~~

~~Store in a cool dry place. Do not contaminate water, food or feed by storage or disposal.~~

~~**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.~~

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities by burning. If burned, stay out of smoke.

APPLICATION DIRECTIONS FOR USE AS A SPRAY

Manex is approved for application in sufficient water to provide good coverage. Rates listed are based on 1 gallon of dilute spray unless otherwise noted. Use at the rate indicated in sufficient water for thorough coverage. Increased volume of water may be necessary as foliage density increases.

When dosage ranges are given, use the higher rate and shorter intervals under severe disease pressure, but do not exceed the maximum rate.

~~If this product is used on a crop, no other product containing a different EBDC active ingredient may be used on the same crop during the same growing season.~~

APPLICATION INSTRUCTIONS FOR APPLICATION

FRUIT AND NUT CROPS

CROPS	DISEASES	TSP PER /GALLON	MAXIMUM TSP/GALLON NUMBER OF APPLICATIONS PER SEASON	USE INSTRUCTIONS
Almonds	Brown Rot Blossom and Twig Blight, Fungus (Leaf Blight), Scab, Shothole	2 1/3 - 3	49 16	Apply at popcorn, full bloom and petal fall or every 7 to 10 days if bloom is staggered. Omit petal fall spray if only brown rot is present. Do not apply later than 5 weeks after petal fall or within 145 days of harvest. Do not allow livestock to graze in almond orchards.
Apples				Use either the Pre-bloom or Extended Application schedules. DO NOT COMBINE OR INTEGRATE THE TWO TREATMENT SCHEDULES.

Rusts (including Cedar apple),
 Fabrea Leaf Spot, Flyspeck,
 Scab, Sooty Blotch

2 1/3

37 15

Pre-bloom: Begin applications at 1/4 to 1/2 inch green tip and continue on a 7 to 10 day application schedule through bloom. Do not graze livestock in treated areas. It is recommended that this product be used in an Integrated Pest Management Program.

1 1/4

32 25

Extended application or tank mix: For implementation of IPM programs, applications based on tree-row volume or for use as a resistance management tool: begin applications at 1/4 to 1/2 inch green tip and continue applications on a 7 to 10 day schedule through the second cover spray or to within 77 days of harvest. Do not apply within 77 days of harvest. Do not graze livestock in treated areas. It is recommended that this product be used in an Integrated Pest Management Program.

Add spreader/sticker to spray mixture for cover sprays and tank mix with an effective systemic protectant/curative fungicide for more effective control of diseases.

Bananas Sigatoka 3-4 1/2 46

Apply when leaves first appear and repeat as needed on a 14 to 21 day interval. May be applied up to the day of harvest.

Cranberries Fruit Rot 7 1/2-9 1/4 28

Spray on a 7 to 10 day interval. Do not apply within 30 days of harvest.

Grapes	Black Rot, Bunch Rot, Deadarm, Downy Mildew	½- 1½	37 24	<u>East of the Rocky Mountains:</u> Apply in sufficient water to provide thorough coverage starting when new shoots are ½ to 1½ inches long and continue at 7 to 10 day intervals until fruit is set. Do not apply within 66 days of harvest.
		½- 1	11½	<u>West of the Rocky Mountains:</u> Apply when shoots are ½ to 1½ inches long and continue at 7 to 10 day intervals. Do not apply within 66 days of harvest except in California. <u>California:</u> Do not apply after bloom.
Kadota Figs	Surface Mold (Cladosporium); Surface Rot (Alternaria)	1¼	4¾	Apply when disease threatens. Do not apply within 10 days of harvest.
Papayas	Anthraenose; Phytophthora; Fruit Rot; Cercospora Black Rot	¾-1	5¾	Apply when disease first threatens and repeat at 14 to 21 day intervals. May be applied up to the day of harvest.

FIELD AND VEGETABLE CROPS

CROPS	DISEASES	TSP PER GALLON	MAXIMUM TSP/GALLON NUMBER OF APPLICATIONS PER SEASON	USE INSTRUCTIONS
Beans (Dry)	Anthraenose; Downy Mildew; Rust	2¼-3	18½	Begin when plants are small. Spray on a 5 to 7 day interval. Do not apply within 30 days of harvest.

Broccoli, Brussels Sprout, Cauliflower	Alternaria Leaf Spot, Downy Mildew	2 1/3 - 3	18 1/2 6	Begin when diseases threaten. Spray on a 7 to 10 day interval. Do not apply within 7 days of harvest.
Cabbage, Kohlrabi, Chinese Cabbage (tight headed only)	Alternaria Leaf Spot, Downy Mildew	2 1/3 - 3	18 1/2 6	Plant beds and direct seeded fields: Spray on a 7 to 10 day interval. Do not apply within 7 days of harvest.
Chinese Cabbage (loose head)	Alternaria Leaf Spot, Downy Mildew	1 1/2 - 2 1/3	13 3/4 5	Plant beds and direct seeded fields: Spray on a 7 to 10 day interval. <u>California:</u> Do not apply within 7 days of harvest. <u>Hawaii:</u> Do not apply within 10 days of harvest.
Corn (dent corn, popcorn, sweet corn used for seed production)	Common Rust, Helmintho- sporium Blight	2 1/3	34 1/2 14	<u>East of the Mississippi</u> <u>including AR and LA:</u> Use sufficient water for thorough coverage. Start applications when disease first appears and repeat at 3 to 10 day intervals. Add a recommended surfactant or spreader/sticker if needed for better coverage. Do not apply within 7 days of harvest. Do not feed treated forage to livestock.
		2 1/3	44 1/2 4	<u>West of the Mississippi except</u> <u>AR and LA:</u> Use sufficient water for thorough coverage. Start applications when disease first appears at 3 to 10 day intervals. Add a recommended surfactant or spreader/sticker if needed for better coverage. Do not apply within 7 days of harvest. Do not feed treated forage to livestock.

Cucumbers	Alternaria Leaf Spot, Angular Leaf Spot, Anthracnose, Downy Mildew, Pythium Fruit Rot	2 $\frac{1}{3}$ - 3	24 $\frac{1}{2}$ 8	Begin when diseases threaten or plants begin to run. Spray on a 7 to 10 day interval. Do not apply within 5 days of harvest.
Eggplant	Anthracnose, Septoria Leaf Spot, Cladosporium Leaf Mold, Early Blight, Late Blight, Gray Leaf Spot (Stemphylium)	2 $\frac{1}{3}$ - 3	24 $\frac{1}{2}$ 7 (per crop)	Begin at first fruit cluster and repeat at 7 to 10 day intervals. Do not apply within 5 days of harvest.
Kale	Alternaria Leaf Spot, Downy Mildew	2 $\frac{1}{3}$ - 3	6 $\frac{1}{4}$ 2 (per cutting)	Begin when disease threatens and apply on a 7 to 10 day interval. Do not apply with 10 days of harvest.
Lettuce (Head and Leaf), Endive	Downy Mildew	2 $\frac{1}{3}$ - 3	48 $\frac{1}{2}$ 6	Apply when disease appears. Spray on a 7 to 10 day interval. Remove residues from head lettuce by stripping and trimming. Do not apply within 10 days of harvest.
			12 $\frac{1}{3}$ 4	California: Spray on a 7 to 10 day interval. Do not apply within 14 days of harvest.
Melons (Cantaloupe, Cassabas, Crenshaws, Honeydew, Watermelons)	Alternaria Leaf Spot, Anthracnose, Cercospora Leaf Spot, Downy Mildew, Gummy Stem Blight	2 $\frac{1}{3}$ - 3	24 $\frac{1}{2}$ 8	Apply as soon as plants begin to run or when disease first appears. Spray on a 7 to 10 day interval. Do not apply within 5 days of harvest.

Onions (Dry Bulb)	Botrytis Leaf Blight, Downy Mildew, Neck Rot, Purple Blotch	3 - 4 $\frac{2}{3}$	46 9	Begin applications when diseases are first reported in the area at 7 day intervals. Do not apply within 7 days of harvest. Do not apply to exposed bulbs.
O n i o n s (Green)	Botrytis Leaf Blight, Downy Mildew, Neck Rot, Purple Blotch	3 - 4 $\frac{2}{3}$	32 $\frac{1}{2}$ 6	Begin application when diseases are first reported in the area at 7 day intervals throughout the season. Do not apply within 7 days of harvest. Do not apply to exposed bulbs.
(F u r r o w drench)	Smut (except in California)	4 $\frac{2}{3}$	4$\frac{2}{3}$ per 435 sq-ft 1	Apply as a furrow drench at time of planting onion seeds. Use $\frac{3}{4}$ to 1 gallon of water per 435 square feet.
Peppers	Anthracnose, Cercospora Leaf Spot (Frogeye Spot), Phytophthora Blight, Ripe Rot	2 $\frac{1}{3}$ - 4 $\frac{2}{3}$	27 $\frac{2}{3}$ 5	<u>East of the Mississippi:</u> Begin when disease threatens. Spray on a 7 to 10 day interval. Do not apply within 7 days of harvest.
		2 $\frac{1}{3}$ - 3	18 $\frac{1}{2}$ 6	<u>West of the Mississippi:</u> Begin when disease threatens. Spray on a 7 to 10 day interval. Do not apply within 7 days of harvest.
Potatoes	Early and Late Blights	2 $\frac{1}{3}$ - 3	21 $\frac{1}{2}$ 7	Begin when plants are 2 to 6 inches high. Apply on a 5 to 10 day interval. Do not apply within 14 days of harvest except for FL, CT, ME, MA, NH, NY, PA, VT and WI where the PHI is 3 days. It is recommended that this product be used with an Integrated Pest Management Program. Vine Kill should occur 14 days before harvest.

Potatoes (Seed Pieces)	Fusarium Seed Piece Decay	15 $\frac{1}{3}$	1	Dip whole or cut tubers. Spread in cool place if held before planting. Seed piece treatment only. Do not use treated seed pieces for food, feed or oil purposes.
Pumpkins	Angular Leaf Spot, Downy Mildew	2 $\frac{1}{3}$ - 3	24 $\frac{1}{2}$ 8	Begin when disease threatens. Spray on a 7 to 10 day interval. Do not apply within 5 days of harvest.
Squash (Winter and Summer)	Anthrachnose, Downy Mildew	2 $\frac{1}{3}$ - 3	24 $\frac{1}{2}$ 8	Begin when plants start to run. Spray on a 7 to 10 day interval. Do not apply within 5 days of harvest.
Sugar Beets	Cereospora Leaf Spot	2$\frac{1}{3}$ - 3	21$\frac{1}{2}$	Apply when disease appears. Spray on a 7 to 10 day interval. Do not apply within 14 days of harvest.
Tomatoes (Greenhouse and Field Garden)	Anthrachnose, Cladosporium Leaf Mold, Early and Late Blights, Gray Leaf Spot (Stemphylium), Septoria Leaf Spot	2 $\frac{1}{3}$ - 4 $\frac{2}{3}$	32 $\frac{1}{4}$ 6	<u>East of the Mississippi:</u> Begin at first fruit cluster and repeat at 7 to 10 day intervals. Do not apply within 5 days of harvest.
		2 $\frac{1}{3}$ - 3	12 $\frac{1}{4}$ 4	<u>West of the Mississippi:</u> Begin at first fruit cluster and repeat at 7 to 10 day intervals. Do not apply within 5 days of harvest.

ORNAMENTALS

CROP	DISEASES	TSP PER #GALLON	USE INSTRUCTIONS
Carnations	Alternaria Leaf Spot, Anthracnose, Botrytis Blight	1½ - 2⅓	Begin when new growth starts. Repeat weekly.
Chrysanthemums	Ascochyta Ray Blight, Botrytis Petal Spot, Septoria Leaf Spot	¾ plus 2¼ oz Captan 50W	Apply twice weekly during the blooming season for petal spot and ray blight. For Septoria leaf spot apply weekly throughout the season.
Dahlias	Alternaria Leaf Spot, Botrytis Blight	1½ - 2⅓	Begin when new growth starts. Repeat weekly.
Dogwood	Anthracnose	1½ - 2⅓	Begin when buds open. Repeat when bracts fall 4 weeks later and in late summer.
Gladiolus	Botrytis Blight, Curvularia and Stemphylium Leaf Spots	2⅓	Begin when flower spikes are developing. Repeat 2 to 3 times at weekly intervals.
Lilies	Botrytis Blight	1½ - 2⅓	Begin with new growth. Repeat weekly.
Pansies	Anthracnose	1½ - 2⅓	Begin with new growth. Repeat weekly.
Peonies	Alternaria Leaf Spot, Botrytis Blight, Phytophthora Blight	1½ - 2⅓	Apply to foliage and soil in early spring and early fall and 7 to 10 day intervals during the growing season.

Roses	Black Spot, Cercospora Leaf Spot (Texas), Rust (California)	1½ - 2⅓	Begin when first leaves unfold. Repeat at 7 to 10 day intervals.
Snapdragons	Rust	1½ - 2⅓	Begin with emergence. Repeat weekly.
Zinnias	Alternaria Blight, Leaf Spot, Botrytis Blight	1½ - 2⅓	Begin with emergence. Repeat weekly.

TURF GRASSES

CROP	DISEASES	USE RATE	USE INSTRUCTIONS
Grasses	Brown Patch	3 tsp in sufficient water/100 square feet	Begin at first sign of disease. Repeat at 7 to 14 day intervals. Do not use on pasture or range grasses. Do not feed clippings to livestock. Do not use if grass is grown for seed.
	Dollar Spot	5 ¼ - 7⅔ tsp in sufficient water/100 square feet	Begin at first sign of disease. Repeat at 7 to 14 day intervals. Do not use on pasture or range grasses. Do not feed clippings to livestock. Do not use if grass is grown for seed.
	Melting Out	2 ¼ - 3 ¼ tsp in sufficient water/100 square feet	Begin at first sign of disease. Repeat at 7 to 14 day intervals. Do not use on pasture or range grasses. Do not feed clippings to livestock. Do not use if grass is grown for seed.
	Leaf, Stem and Stripe Rusts	4½ tsp in sufficient water/100 square feet	Begin when rust pustules are first seen. Repeat at 7 to 14 day intervals. Do not graze treated areas or feed clippings to livestock.

Our recommendations for use are based on tests believed reliable. Since the use is beyond our control, we can not guarantee the results if such use is not in accordance with directions. We disclaim any responsibility for damages resulting from careless or improper handling or use.

NOTICE OF WARRANTY

GRIFFIN warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of GRIFFIN. In no case shall GRIFFIN be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. GRIFFIN MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESSED OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

Manex™ is a trademark of Griffin Corporation
Captan is a registered trademark of ICI Americas Inc.