

SUPER 16

SULFUR-TIN FLOWABLE FUNGICIDE

ACTIVE INGREDIENTS:

| | |
|------------------------|--------|
| Sulfur | 40.0% |
| Triphenyltin Hydroxide | 5.1% |
| NET WT INGREDIENTS | 54.0% |
| TOTAL | 100.0% |

Formulated to contain no free sulfur.

KEEP OUT OF REACH OF CHILDREN

DANGER

Use only as directed on label. Do not eat or drink while using product. Do not apply to food stuffs and immediately con-

tact of spray with skin or eyes. If contact occurs, wash thoroughly with soap and water. If eye contact occurs, flush eyes with water for at least 15 minutes. If skin contact occurs, wash with soap and water. If swallowed, do not induce vomiting. Call a physician or poison control center for assistance and see doctor if necessary.

For additional Product Safety Information see label.

DO NOT DRINK

DO NOT EAT

DO NOT DRINK

GRiffin CORPORATION
VALDOSTA, GEORGIA 31601

NET CONTENTS 1 GALLON



GRiffin SUPER 16 Fungicide can be applied as a ground or aerial spray to control fungal infestations on listed crops. Application rates are for general use. The state agricultural extension or agricultural experiment station specialists should be consulted for specific applications and timing recommendations. With any spray, application thorough coverage is essential for good control.

Apply this product only as specified on this label.

PRECAUTIONS

Do not add sulfur to other substances or stickers. Do not graze dairy or meat animals in treated areas.

GENERAL INSTRUCTIONS FOR APPLICATION

PEANUTS *Cucumis sativus L.* *Rust*

Ground application: Apply 2-3 pints per acre in 10-120 gallons of water. Aerial application: Apply 2-3 pints per acre in 3-10 gallons of water. Use lower rate when leafspot is light, increasing as disease pressure increases.

Apply in sufficient water for uniform coverage. Use lower pressure in early season when plants are smaller and increase spray volume as plant size increases. Application should begin at planting, six to 6½ weeks after planting, or as soon as first signs of leafspot appear. Spray application should continue until 10-14 day schedule. Do not apply within 14 days of harvest. Do not allow hogs to feed on peanuts or treated hulls. Hulls from treated peanuts may be used for livestock. Do not use water to feed.

WARRANTY STATEMENT

Griffin Corporation warrants to the original purchaser that this product will contain the active ingredients indicated in the label. If any active ingredient is found to be less than the labeled amount, the difference will be paid by the manufacturer.

2073

3083

BEST DOCUMENT AVAILABLE