

# Yellow Jacket Wettable Sulfur II

For Spraying Fruits,  
Vegetables & Ornamentals.

ACTIVE INGREDIENT:	
Sulfur .....	90.0%
INERT INGREDIENTS: .....	10.0%
TOTAL .....	100.0%

FINENESS: Not less than 93% passing 325 mesh.

## KEEP OUT OF REACH OF CHILDREN CAUTION

If on skin, wash immediately with soap and water.

If in eyes, immediately flush eyes with plenty of clear water for at least 15 minutes.

If swallowed, drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

See Side Panel For Additional Precautionary Statements.

APR 8 1981

GRIFFIN CORPORATION

EPA REG. NO. 1812-255

VALDOSTA, GEORGIA 31601

EPA EST. NO. 6325-GA 1

NET CONTENTS 50 lbs.

## RECOMMENDATIONS

Consult State Agricultural Experiment Stations or the State Agricultural Extension Service for additional information as the timing and number of applications needed will vary with local conditions.

All dosages given on this label are for dilute sprays (IX concentrations). For use in concentrate sprays, increase the dosage of Yellow Jacket Wettable Sulfur II proportionately. Spray often enough to keep new growth covered.

**IMPORTANT:** Use the higher dosage during periods of frequent and heavy rainfall. Spray often enough to keep susceptible growth covered. This may require weekly or bi-weekly applications during periods of rapid growth and frequent rainfall. When using Lead Arsenate with Yellow Jacket Wettable Sulfur II always add the same amount of Spray Lime, or other safener, as would be used when applying Lead Arsenate without sulfur. When using Zinc Sulfate and Spray Lime as a safener on peaches, always add Zinc Sulfate to the spray tank first, followed by Spray Lime, Lead Arsenate and sulfur in order given.

**Apple, Pear: Powdery Mildew, Scab** — Use 6 to 8 lbs. per 100 gals. of water in the pre-bloom, bloom and petal fall sprays. Use 4 to 6 lbs. in the cover sprays.

**Cherry: Brown Rot, Leaf Spot** — Use 6 to 8 lbs. per 100 gals. of water in pink, bloom, petal fall, shuck, cover and post-harvest sprays.

**Citrus: Rust Mite** — Use 5 lbs. per 100 gals. of water. Begin applications at first signs of damage and repeat as necessary. Rust mites should be controlled at all times of the year in order to prevent russetting of fruit and greasy spot on leaves. **Scale Crawlers** — Add 5 to 8 lbs. to each 100 gals. of tank strength lime sulfur solution.

**Peach: Brown Rot, Scab** — Use 6 to 8 lbs. per 100 gals. of water in pink, bloom, shuck, cover and pre-harvest sprays.

**Plum, Prune: Brown Rot, Leaf Spot** — Use 6 to 8 lbs. per 100 gals. of water in pink, bloom, shuck, cover and pre-harvest sprays.

**Beans, Peas: Powdery Mildew, Rust** — Use 16 lbs. per acre in sufficient water for thorough coverage. Repeat applications in 7 to 14 day intervals as necessary.

**Roses: Black Spot, Powdery Mildew** — Use 6 to 8 lbs. per 100 gals. of water (3 to 4 tablespoons per 1 gal. of water). Make first application prior to first signs of disease and maintain preventive schedule throughout the season, repeating applications every 7 to 14 days. Keep new growth protected.

**Perennials, Shrubs: Powdery Mildew, Rust** — Use 6 to 8 lbs. per 100 gallons of water (3 to 4 teaspoons per 1 gallon of water). Make first application prior to first signs of disease and maintain preventive schedule throughout the season, repeating applications every 7 to 14 days. Keep new growth protected.

## WARRANTY STATEMENT

GRIFFIN CORPORATION warrants that its product in its unopened package conforms to the chemical description on the label and is reasonably fit for the purposes set forth on the label when used according to directions under normal use conditions on the crops specified. There are no other warranties, expressed or implied, concerning the use of this product other than indicated on the label. This warranty does not extend to the handling or use of this product contrary to label instructions or under abnormal conditions or under conditions not reasonably foreseeable to seller, and buyer assumes all risk of any such use.

Custom Printing No. 6072

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### WARNING

Harmful if swallowed. Avoid contact with skin or clothing. Causes eye irritation. Do not get in eyes. Wash after handling. Avoid breathing vapor. Do not spray on humans or pets. Do not spray where food or drink is exposed or being consumed.

#### ENVIRONMENTAL HAZARDS

Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. When used on ships do not wash or flush freshly sprayed surfaces into adjacent waters.

#### FIRST AID

If swallowed, induce vomiting by sticking finger in back of throat or by administration of syrup of ipecac; get medical attention. If in eyes, flush with water for 15 minutes and see a physician. For skin, wash with plenty of soap and water. If irritation develops, get medical attention.

#### PHYSICAL OR CHEMICAL HAZARDS

#### DIRECTIONS FOR USE GENERAL CLASSIFICATION

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

#### STORAGE AND DISPOSAL

Store at temperatures above 32°F. Do not reuse empty container. Wrap container and put in trash collection.

Do not use in edible product areas of food processing plants, restaurants, or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed. In the home cover or remove exposed foods. Avoid contamination of cooking utensils and food preparation surface.

## SYNTHETIC INSECTICIDE W.B. 2%

### A SYNTHETIC PYRETHROID DUAL ACTION INSECT KILLER KILLS AS CONTACT AND RESIDUAL SPRAY KILLS ROACHES UP TO 4 WEEKS

#### ACTIVE INGREDIENTS

\*+ resmethrin (5-(phenylmethyl)-  
3-(2-methyl-1-propenyl) cyclopro-  
panecarboxylate.....2.000%w/w  
Related compounds.....0.274%

INERT INGREDIENTS.....97.726%  
100.000%w/w

\*Cis/trans isomers ratio: max. 30% (+) cis and min.  
70% (+) trans.  
+U.S. Patent No. 3,465,007.  
+U.S. Patent No. 3,683,078.  
#SBP-1382© of Penick Corporation, 1050 Wall Street  
West, Lyndhurst, N.J. 07071

Contains 0.17 pounds per gallon

## KEEP OUT OF REACH OF CHILDREN WARNING

HAVE CLEAN WATER SUPPLY AT HAND  
IN CASE OF AN EYE EMERGENCY  
WHEN DILUTING THIS CONCENTRATE  
SEE SIDE PANEL FOR ADDITIONAL  
PRECAUTIONARY STATEMENTS

CAN BE USED IN NON-FOOD AREAS  
OF FOOD HANDLING ESTABLISHMENTS

MANUFACTURED BY  
**BIXON CHEMICAL CO.**  
50-1997 PLACE  
CORONA, NEW YORK 11368

EPA Reg. No. 6125-42 EPA Est. 6125NY01

NET CONTENTS  
NET WEIGHT

GALLONS  
POUNDS

## DILUTION INSTRUCTIONS

For an emulsion containing 0.35% dilute

Concentrate Fluid Ounces	To Volume Below With Water
1 1/4	8 fluid ounces
3	15 fluid ounces
6	32 fluid ounces
11.5	1/2 gallon
23	1 gallon

MIX WELL BEFORE USING.

#### APPLICATIONS:

##### INSTRUCTIONS FOR CONTACT AND RESIDUAL SPRAYS

**FOR INDOOR USE:** In non-food areas of food handling establishments, commercial, industrial, office, residential, public, institutional and storage buildings, homes, hotels, ship cabins and hospitals. In hospitals patients should be removed prior to treatment. Also for use in buses, automobiles, and non-food areas of trucks, boxcars and ship cargo holds. Do not use on electrical equipment or on surfaces that can be harmed or stained by water. Also for use indoors in dog kennels and horse stables. In horse stables, not for application if horses will be used for food.

**CONTACT SPRAY - KILLS AND REDUCES INFESTATIONS:**  
**COCKROACHES, CRICKETS, SILVERFISH, ANTS:** Apply as a contact spray. Fill spray tank with 0.35% Transparent Emulsion Spray formulation. Keep spray nozzle about 8 to 12 inches from surface to be treated. Adjust pressure and nozzle to deliver a coarse wet spray. When possible, spray directly on pests to reduce infestations. Spray thoroughly into hiding places such as cracks, crevices, moist areas, openings around pipes and sinks, under refrigerators and other appliances, baseboards, and storage areas. Apply spray until area is moist, but not to the point of run-off. Spray directly on insects where possible. Repeat as necessary.

**ANTS:** To kill ants: Spray trails, nests and points of entry within buildings and homes. Spray on ants where possible. Repeat as necessary.

**FLEAS:** Aids in the control of fleas on pet bedding and in horse stalls. Hold spray nozzle about 18 inches from areas to be treated. Direct spray on resting areas and floor around these areas. Apply spray until area is moist. Treat pet bedding the same way. Also spray entire inside surface of dog houses. Repeat as necessary. Do not allow pets to contact treated areas until spray has dried. Remove or cover animal feed before spraying. Also, treat dogs, cats and horses with a registered flea control product before they re-enter the treated area.

**RESIDUAL SPRAY:** Aids in the control of: **COCKROACHES, CRICKETS, SILVERFISH, ANTS.** Kills for up to 4 weeks: Spray baseboards, cracks and crevices, around door and window frames, and other hiding places such as closets, in back of and under appliances, sinks and cupboards in kitchens and bathrooms. In basements, spray hiding places in cracks and crevices, lower walls and dark places. Apply as a coarse wet spray in a 4 to 6 inch band until area is thoroughly wet but not to the point of run-off. Repeat as necessary.

## WARRANTY STATEMENT

Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such, or in combination with other materials.