### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

## CAUTION

Harmful if swallowed, inhaled or absorbed through the skin. Do not gl eyes or on skin. Do not breathe spray mist. Wash thoroughly after handling. If swallowed, induce vomiting. Get medical attention of inhaled, remove victim to fresh air. If on skin, remove by washing. Get medical attention if irritation persists. If in eyes, flush with plenty of water. Get medical attention.

## **ENVIRONMENTAL HAZARDS**

This product is toxic to fish and wildlife. Keep out of any body of water. Do not apply where weather conditions favor drift from areas treated. Do not contaminate, water by cleaning of equipment or disposal of wastes.

#### DIRECTIONS FOR USE

It is eviolation of Federal law to use this label in a manner inconsistent with its label-

## STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not rause empty container. Pesticide, spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of in landfill approved for pesticides or buried in a safe place away from water supplies.

Container Disposal: Triple rinse (or equivalent) and dispose in an incinerator or landfill approved for pesticide containers, or bury in a safe place. Consult Federal, State or Local Disposal Authorities for approved alternate procedures such as limited open burning.

## **GENERAL INFORMATION**

Apply this product only as specified on this label.

Local conditions may effect the use of this chemical. Consult State Agricultural Extension Service or Experiment Station weed specialists for specific recommendations for local weed problems...


DI-Trol 4L is to be mixed with water and applied as a spray for control of weeds. Effects are slow to appear and will not become apparent until the chemical has been carried into the root system by mcisture.

IMPORTANT: Do not apply (except as recommended for crop use) or drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend or in locations where the chemical may be washed or moved into contact with their roots. Do not use on lawns, walks, driveways, tennis courts, or similar areas. Prevent drift of spray to desirable plants. Do not contaminate domestic waters. Keep from contact with fertilizers, insecticides, fungicides and seeds.

Thoroughly clean all traces of Di-Trol 4L from application equipment immediately after use: otherwise crop injury may result when equipment is used again. Flush tank, pumps, hose and boom with several changes of water after removing notice. tips and screens (clean parts separately).

#### Shake or stir well before using

Before spraying, calibrate equipment to determine quentity of water necessary to uniformly cover measured area to be restrict. Carefully measure proper amount and mix into necessry volume of water water serves only as a carrier. Material must be kept in suspension at all times by continuous agitation. Except for small areas, use



Diuron Flowable de bicide

For Control of Many A super case Perennial Grasses and Herbeeneous Warrant

**ACTIVE INGREDIENT:** Diuron (3-13, 4-dichlorophen // 1-aimethylurea)\*.....40.8% 

\*Contains 4.0 lbs. of Diuron per gallon.

# CAUTION KEEP OUT OF REACH OF CHILDREN

if swallowed, drink 1 or 2 glasses of water. Induce vomiting, by placing finger in back of throat. Call a physician. Never give anything by mouth to an unconscious person.

If inhaled, remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth to mouth. Get medical attention.

If on skin, remove by washing.

If in eyes, flush with plenty of water. Get medical attention if imitation persists.


 $I_{j}I_{j}$ 

See Side Panel For Additional Precautionary Statements.



**GRIFFIN CORPORATION** MET CONTENTS 5 Gallons

> EPA Reg. No. 1812-257 EPA Est. No. 1812-GA-3



## Diuron Flowable Herbiade

For Control of Many Annual and Perennial Grasses and Herbecaeous Weeds

**ACTIVE INGREDIENT:** Diuron (3-[3, 4-dichlorophenyl] -1-dimethylurea)\* . . . . . 40.9% INERT INGREDIENTS:.....60.0%

\*Contains 4.0 lbs. of Diuron per gallon.

# **CAUTION** KEEP OUT OF REACH OF CHILDREN

If swallowed, drink 1 or 2 glasses of water. Induce vomiting, by placing finger in back of throat. Call a physician. Never give anything by mouth to an unconscious person.

If inhaled, remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth to mouth. Get medical attention.

If on skin, remove by washing.

If in eyes, flush with plenty of water. Get medical attention if irritation persists.

See Side Panel For Additional Precautionary Statements.



#### GRIFFIN CORPORATION VALDOSTA, GEORGIA 31601 NET CONTENTS 5 Gallons

EPA Reg. No. 1812-257

EPA Est. No. 1812-GA-3

fixed boom power sprayers properly calibrated to insure constant rate of application Openings in screens should be equal to or larger than 50 mesh. Agitate by mechanical or hydraulic means in the spray tank. If by pass or return line is used, it should terminate at bottom of tank to minimize foaming. Do not use air agitation. For general weed control in small areas, tank type hand sprayer or sprinkling can may be used; shake or stir frequently.

#### NON-CROP WEED CONTROL

Di-Trol 4L is an effective general herbicide for the control of many annual and perennial grasses and herbaceous weeds on non-cropland creas where bare ground is desired. The degree of control and duration of effect will vary with amount of chemical applied, soil type, rainfall and other conditions. May be used at any time for non cropland weed control, except when ground is frozen, provided adequate moisture is supplied by rainfall or artificial means. Best results are obtained if applied shortly before weed growth begins. If dense growth is present, remove tops and spray the ground.

To control most weeds for an extended period of time on non-cropland such as: utility, highway, pipeline and railroad rights-of-way, petroleum tank farms, lumberyards, storage areas, industrial plant sites and around farm buildings -- Apply 1 to 4 gallons per acre to control most annual weeds. Use 4 to 8 gallons per acre to control both annual and perennial weeds. Repeat treatment may be required where a longer peiod of control is desired or when hard to kill, deep-rooted perennial weeds such as Johnsongrass alle present.

For irrigation and drainage ditches, apply 1% plats per 1,000 sq. ft. (8 gallons per acrel. For irrigation ditches, apply during the non-crop season, and when ditch is not in use. To minimize movement of Di-Trol 4L with irrigation water and avoid crop injury, it is essential that it be fixed in the treated soil by moisture. Apply before expects I seasonal rainfall, if possible when soil in the ditch is still moist. If rainfall has not totaled at least 4 inches following treatment and before intended use of irrigation ditch, fill with water and allow to stand for 72 hours; drain off and waste remaining water before using ditch. Do not treat any ditch area into which roots of trees or other desirable plants may extend.

For small areas, apply 1/2 pint per 270 sq. ft. (approximately 10 gallons per acre).

## **COTTON (Lay-By)**

Use 1 ½ to 2 pints (1 ½ to 3 pints in California and Arizons) in 25 to 40 gallons of water per acre; use the higher rates or heavier soil types (those high in clay and/or organic matter). Do not use on light (sandy or low organic) soils as injury to the cotton may result. Apply when cotton is at least 12 inches high (at least 20 inches for Pinia S-21. Keep contact of cotton plants with spray or drift to a minimum.

Do not spray over top of cotton.

For control of germinating weed seedlings, apply immediately after the last cultivation, directing spray to cover the soil beneath cotton plants and between rows. Alternatively, for control of emerged annual weeds (4 inches or less in height) at layby time, apply as a directed spray to cover weed foliage beneath cotton plants and

In impated cotton, best weed control is obtained if the field is irrigated within 3 to 4 days after application; thoroughly wet the surface of the ground over the row to carry the chemical into the root zone of germinating weeds.

Do not replant treated areas to any crop within one year after last application as injury to subsequent crops may result, except that cotton, corn, or sorghum (grain sorghums only; not sorgos or forage sorghums, or grass sorghums) may be planted in the spring of the year following treatment.

WARRANTY STATEMENT: :

GRIFFIN CORPORATION warrants that this product in its unopened package conforms to the chimical description on the label and is reasonably fit for the purposes set forth on the label when used according to directions under normal use conditions on the crops sepecifed. There are no other warranties, expressed or implied, concerning the use of this product other than indicated on the label. This warranty does not extend to the handling of use of this product contrary to label instructions or under abnormal conditions or index conditions not reasonably foreseeable to seller and buyer assumes all risk of eny soch use.

GCN 00281 CPN 06517

ramount and terial must be all areas, use

necessary to

ALS

fter handl-

inhaled,

dical at-

ter. Get

ater. Do

Do not

with its tabel-

ıl. Open

pray mix-

hould be

ace away

n an in-

ı⊓ a safe pproved

oricultural Ex-eccimmenda-

of weeds. Efmical has been

drain or flush

re their roots

ved into con-

is courts, or

nate domestic

t immediately again. Flush

noving nozzle

i seeds.