

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS**

CAUTION

In case of contact with skin, wash thoroughly with soap and water. In case of contact with the eyes, flush with clean water and seek medical attention. If swallowed, do not induce vomiting. If inhaled, remove victim to fresh air. If not breathing, give artificial respiration. If necessary, use mouth-to-mouth respiration. Get medical attention. For more information, see the label. **Note to Physicians:** This product is toxic to the central nervous system.

ENVIRONMENTAL HAZARDS

Keep out of reach of children. Do not use near water. Do not apply to water bodies.

DIRECTIONS FOR USE

Use as a preplant or preemergence herbicide in sorghum, milo and sweet sorghum.

STORAGE & DISPOSAL

Do not contaminate water. Do not feed or use as fertilizer. Do not use in areas where it may be applied to water. Do not use in areas where it may be applied to water. Do not use in areas where it may be applied to water.

GENERAL INFORMATION

Failure to follow all precautions on this label may result in crop loss or injury to humans and animals. This product is a selective herbicide that can be applied to sorghum, milo and sweet sorghum. It is effective against many broadleaf weeds and grasses. It is also effective against some narrowleaf weeds. It is also effective against some grasses. It is also effective against some broadleaf weeds. It is also effective against some grasses. It is also effective against some broadleaf weeds. It is also effective against some grasses.

MIXING PROCEDURES

This product is a water dispersible granule. It is mixed with water. It is mixed with water. It is mixed with water.

Unless otherwise specified, use a minimum of 10 gallons of spray mixture. For preemergence applications, use the amount specified on the label. For postemergence applications, use the amount specified on the label.

Band width in inches. Rate: Apply to the band. Apply to the band. Apply to the band.

Be sure sprayer is clean and not contaminated with any material. Do not use in areas where it may be applied to water. Do not use in areas where it may be applied to water. Do not use in areas where it may be applied to water.

Liquid Fertilizer: If a liquid fertilizer is used, it should be applied to the soil. It should be applied to the soil. It should be applied to the soil.

APPLICATION EQUIPMENT

Ground Equipment: Use a pump sprayer or backpack sprayer. Use a pump sprayer or backpack sprayer. Use a pump sprayer or backpack sprayer.

Use a pump with capacity of 10 gallons. Use a pump with capacity of 10 gallons. Use a pump with capacity of 10 gallons.

Use a nozzle that is suitable for the application. Use a nozzle that is suitable for the application. Use a nozzle that is suitable for the application.

AERIAL APPLICATION PROCEDURE

Use only where broadcast application is indicated. Apply a minimum of 10 gallons of water per acre. Apply a minimum of 10 gallons of water per acre. Apply a minimum of 10 gallons of water per acre.

INSTRUCTIONS FOR SAFE HANDLING

Do not apply directly to water. Do not apply directly to water. Do not apply directly to water.

PREPLANT APPLICATION (PRE-PLANT)

For preplant application, apply to the soil. For preplant application, apply to the soil. For preplant application, apply to the soil.

ACCEPTED

FEB 12 1981

U.S. Environmental Protection Agency, Washington, D.C. 20460
Approved for the pesticide
propazine under
EPA Reg. No. 1812-CA-1

Milo-PRO 90

**DRY FLOWABLE HERBICIDE
For Weed Control in Sorghum
(milo and sweet sorghum)**

ACTIVE INGREDIENT	
Propazine: 2-chloro-4,5-bis (isopropylamino) s-triazine	90.0%
INERT INGREDIENTS	10.0%
TOTAL	100.0%

CAUTION

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

- If on skin, wash immediately with soap and water.
- If in eyes, flush with water for at least 15 minutes. Get medical attention if irritation persists.
- If swallowed, drink promptly a large quantity of milk, egg whites or gelatin solution. If these are not available, drink large quantities of water. Induce vomiting. Get medical attention.
- If inhaled, remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

See Side Panel For Additional Precautionary Statements.



GRIFFIN CORPORATION
VALDOSTA, GEORGIA 31601
NET CONTENTS

EPA Reg. No. 1812-CA-1
EPA Reg. No. 1812-CA-1

ACCEPTED
FEB 12 1981
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, extended for the pesticide registered under EPA Reg. No. 1812-200

Milo-Pro 90

**DRY FLOWABLE HERBICIDE
 For Weed Control in Sorghum
 (milo and sweet sorghum)**

ACTIVE INGREDIENT:
 Propazine: 2-chloro-4,6-bis(isopropylamino)-s-triazine 90.0%
INERT INGREDIENTS 10.0%
TOTAL 100.0%

CAUTION

**KEEP OUT OF REACH OF CHILDREN
 CAUTION**

If on skin, wash immediately with soap and water.

If in eyes, flush with water for at least 15 minutes. Get medical attention if irritation persists.

If swallowed, drink promptly a large quantity of milk, egg whites or gelatin solution. If these are not available, drink large quantities of water. Induce vomiting. Get medical attention.

If inhaled, remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

See Side Panel For Additional Precautionary Statements.



GRIFFIN CORPORATION
 VALDOSTA, GEORGIA 31601
 NET CONTENTS

EPA Reg. No. 1812-200
 EPA Est. No. 1812-01-001

Soil texture	Broadcast rate per acre
Sand	DO NOT USE
Sandy loam	2.2 lbs
Loam	2.7 lbs
Heavy clay	DO NOT USE

PREPLANT APPLICATIONS (OK, NM, TX, The Gulf Coast States, AR, GA)
 Apply this product to the soil surface before or after planting sorghum. Do not apply this product to the soil surface after sorghum has emerged. Apply this product to the soil surface before or after planting sorghum. Do not apply this product to the soil surface after sorghum has emerged. Apply this product to the soil surface before or after planting sorghum. Do not apply this product to the soil surface after sorghum has emerged.

Soil texture	Broadcast rate of this product per acre
Sand	DO NOT USE
Sandy loam	1.33 lbs

Soil texture	Broadcast rate of this product per acre
Sand	DO NOT USE
Sandy loam	2.2 lbs
Loam	2.7 lbs
Heavy clay and high organic matter soils of Arkansas, Georgia, and the Gulf Coast States including Texas Gulf Coast	3.6 lbs

Except in the states of Arkansas, Georgia, and the Gulf Coast States including Texas Gulf Coast, broadcast rate of 1.33 lbs of this product per acre is recommended for Broadleaf Weed Control.

PREEMERGENCE APPLICATIONS
 Apply this product to the soil surface before sorghum emerges for weed control before sorghum emerges.

Soil texture	Broadcast rate of this product per acre
Sand	DO NOT USE
Sandy loam	1.33 lbs

Soil texture	Broadcast rate of this product per acre
Sand	DO NOT USE
Sandy loam	2.2 lbs
Loam	2.7 lbs
Heavy clay and high organic matter soils of Arkansas, Georgia, and the Gulf Coast States including Texas Gulf Coast	3.6 lbs

Except in the states of Arkansas, Georgia, and the Gulf Coast States including Texas Gulf Coast, broadcast rate of 1.33 lbs of this product per acre is recommended for Broadleaf Weed Control.

See the General Information section to calculate amount per acre for band treatment.

CROP ROTATION SUGGESTIONS
 Do not rotate with corn, cotton, soybeans, or other crops in the same family as sorghum except as noted below. Do not rotate with corn, cotton, soybeans, or other crops in the same family as sorghum except as noted below. Do not rotate with corn, cotton, soybeans, or other crops in the same family as sorghum except as noted below.

WARRANTY STATEMENT
 Griffin Corporation warrants that the product in its original container and package conforms to the chemical analysis and specifications set forth on the label when used according to the directions on the label. There are no other warranties, expressed or implied, made by Griffin Corporation. This warranty does not extend to abnormal conditions or under abnormal conditions of use.