

Needs controlled

Downy brome (cheatgrass),
tansymustard, prickly lettuce,
pepperweed, seedling shepherd's
purse, yellow flower pepperweed

Soil texture	Rate per acre
Sand and loamy sand	DO NOT USE
Sandy loam	0.9 lb
Loam soil	0.9-1.3 lbs
Clay soil	1.3-1.75 lb

Henbit, wild mustard, common chickweed, knawel (German moss),
Alyssum, downy brome (cheatgrass),
wild oats, pepperweed, tansymustard,
ard, pigweed, speedwell, and seedlings of yellow rocket, shepherd's
purse, white cockle, fanweed (pennycress),
dandelion, Russian thistle,
kochia

Soil texture	Rate per acre
Sand and loamy sand	DO NOT USE
Sandy loam	0.9 lb
Loam soil	1.3 lbs
Clay soil	1.75 lbs

Pigweed, common chickweed,
mouseear chickweed, speedwell,
henbit, knawel (German moss),
pepperweed, downy brome, alyssum,
common lambsquarters, wild oats,
and seedlings of shepherd's
purse, fanweed (pennycress), yellow rocket,
wintercress, Russian thistle,
dandelion, kochia, white cockle

Soil texture	Rate per acre
Sand and loamy sand	DO NOT USE
Sandy loam	0.9 lb
Loam soil	1.1 lbs
Clay soil	1.4 lbs

Year Old (Northeastern U.S. Only)

Apply in the spring before June 1 and has become well established and is
germinating but before permanently frozen ground conditions. Apply in the fall
when dormant when this product is applied. Apply 0.9 lb per acre in the
fall. Apply 0.9 lb per acre in the fall. Apply 0.9 lb per acre in the fall.
lambsquarters, common chickweed, German moss, downy brome, wild
mustard, pigweed, speedwell, and pepperweed

Grass Mixtures (Northeastern U.S. Only)

Apply to grass mixtures which have been established at least one year. Mixtures
in orchards or pastures which have gone dormant. Apply 0.9 lb per acre in the
fall. Apply 0.9 lb per acre in the fall. Apply 0.9 lb per acre in the fall.
Alyssum, downy brome (cheatgrass), wild oats, shepherd's purse

- 1) Do not use on gravelly, sandy or heavy clay soils where crop injury may occur.
- 2) Applications made to active growing crops may result in crop injury.
- 3) Use only on crops which are in dormant or dormant state.
- 4) Do not apply on crops.
- 5) This product will control shed perennial or biennial weeds such as Canada goose, wild carrot, and dock.
- 6) Temporary crop chlorosis may appear after application. The sprayer should be shut off when turning around at the end of a row. Do not apply to any crops except corn for the growing season after application.

Application and grazing of dairy and beef cattle and sheep for hay

Use as Coastal and Midland

Apply to most annual broad leaf and grass weeds including hairy vetch, chickweed, tail, carelessweed, carpetweed, annual meadow grass, pigweed, chickweed, henbit, and burdock, apply to persistent weeds including
rudgrass from fall to spring depending on the weed and soil conditions. Apply at the appropriate rate in the following table in a minimum of 150 gal. of water per acre. A minimum of 1 gal. of water for each 0.9-1.3 lbs. of product applied.

	Rate per acre
Organic matter	0.9 lb
Clay matter	1.3-1.75 lbs
Organic matter	2.2-2.6 lbs
Clay matter	2.6 lbs

Apply to cattle or sheep for 30 days after 0.9-2.6 lbs. per acre application or for 60 days after 2.2-2.6 lbs. per acre application. Do not cut for hay for 60 days after 0.9-2.6 lbs. per acre application or for 90 days after 2.2-2.6 lbs. per acre application.

Liquid fertilizers may replace all or part of the water as a carrier. Do not apply if there is danger of liquid fertilizers causing crop injury.

Apply to weeds and corn emergence. Use the appropriate rate in the table below. Apply before plowing at the appropriate rate on the table below. Apply before plowing. If soil is tilled or work is done after application, apply this product is applied within 2 weeks after application.

Soil texture	Rate per acre
Silt and clay	0.9 lb
Clay matter	1.3-1.75 lbs
Organic matter	2.2-2.6 lbs
Clay matter	2.6 lbs

Apply to high nitrogen soils. Apply to high nitrogen soils. Apply to high nitrogen soils.

Apply to high nitrogen soils. Apply to high nitrogen soils. Apply to high nitrogen soils.

Apply to high nitrogen soils. Apply to high nitrogen soils. Apply to high nitrogen soils.

Apply to high nitrogen soils. Apply to high nitrogen soils. Apply to high nitrogen soils.

Apply to high nitrogen soils. Apply to high nitrogen soils. Apply to high nitrogen soils.

Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall.

Grasses Grown for Seed (Northeastern U.S. Only)

Apply to ryegrass, timothy, and other grasses. Apply to ryegrass, timothy, and other grasses. Apply to ryegrass, timothy, and other grasses.

Apply to ryegrass, timothy, and other grasses. Apply to ryegrass, timothy, and other grasses. Apply to ryegrass, timothy, and other grasses.

NURSERIES, CHRISTMAS TREE PLANTINGS, SHELTERBELTS

Nurseries (see list below). Nurseries (see list below). Nurseries (see list below).

Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall.

Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall.

Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall.

Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall.

Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall.

Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall.

Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall.

Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall.

Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall.

Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall.

Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall.

Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall.

Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall.

Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall.

Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall.

Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall.

Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall.

Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall.

Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall.

Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall.

Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall.

Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall.

Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall.

Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall.

Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall.

Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall.

Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall.

Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall.

Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall.

NONSELECTIVE WEED CONTROL ON NONCROP LAND

Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall.

Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall.

Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall.

Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall. Apply 2.2-4.4 lbs. per acre in the fall.

