

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
CAUTION**

Use only as directed. Do not breathe dust or spray. Do not take internal. Avoid contact with eyes, nose, mouth, and skin. Wash thoroughly after handling. Do not eat, drink, or use tobacco during or after use. Wear regular long-sleeved shirt and pants. Avoid contact with bare skin. Do not use in confined areas. Do not use in areas where children or animals are present. Do not use on crops or animals for which it is not labeled. Do not use on crops or animals for which it is not labeled. Do not use on crops or animals for which it is not labeled.

ENVIRONMENTAL HAZARDS

This insecticide is highly toxic to fish, birds, and bees. Do not apply to water bodies. Do not apply to areas where bees are present. Do not apply to areas where birds are present. Do not apply to areas where fish are present. Do not apply to areas where crops or animals are present for which it is not labeled. Do not use on crops or animals for which it is not labeled. Do not use on crops or animals for which it is not labeled.

DIRECTIONS FOR USE

Read and follow all directions on the product in a manner consistent with its labeling.

STORAGE AND DISPOSAL

Store in original container in a cool, dry place. Do not store in areas where children or animals are present. Do not store in areas where crops or animals are present for which it is not labeled. Do not use on crops or animals for which it is not labeled. Do not use on crops or animals for which it is not labeled.

GENERAL INFORMATION

SAVIT BAW is a registered trademark of Griffin Corporation. All rights reserved. Do not use on crops or animals for which it is not labeled. Do not use on crops or animals for which it is not labeled. Do not use on crops or animals for which it is not labeled.

SPECIAL DIRECTIONS FOR PREPARING LOW-GALLONAGE SPRAYS

In low-gallonage ground equipment, use 1/2 gallon of water per gallon of concentrate. In low-gallonage backpack sprayers, use 1/2 gallon of water per gallon of concentrate. Do not use on crops or animals for which it is not labeled. Do not use on crops or animals for which it is not labeled. Do not use on crops or animals for which it is not labeled.

For application to crops or animals, use the amount of concentrate specified on the label. Do not use on crops or animals for which it is not labeled. Do not use on crops or animals for which it is not labeled. Do not use on crops or animals for which it is not labeled.

Do not use on crops or animals for which it is not labeled. Do not use on crops or animals for which it is not labeled. Do not use on crops or animals for which it is not labeled. Do not use on crops or animals for which it is not labeled.

Do not use on crops or animals for which it is not labeled. Do not use on crops or animals for which it is not labeled. Do not use on crops or animals for which it is not labeled. Do not use on crops or animals for which it is not labeled.

GENERAL CAUTION

Do not use on crops or animals for which it is not labeled. Do not use on crops or animals for which it is not labeled. Do not use on crops or animals for which it is not labeled. Do not use on crops or animals for which it is not labeled. Do not use on crops or animals for which it is not labeled. Do not use on crops or animals for which it is not labeled.

**IMPORTANT PRE-HARVEST
AND GRADING USE INFORMATION AND LIMITATIONS**

Do not use on crops or animals for which it is not labeled. Do not use on crops or animals for which it is not labeled. Do not use on crops or animals for which it is not labeled. Do not use on crops or animals for which it is not labeled. Do not use on crops or animals for which it is not labeled. Do not use on crops or animals for which it is not labeled.

DIRECTIONS FOR USE

Read and follow all directions on the product in a manner consistent with its labeling.

Use only as directed. Do not breathe dust or spray. Do not take internal. Avoid contact with eyes, nose, mouth, and skin. Wash thoroughly after handling. Do not eat, drink, or use tobacco during or after use. Wear regular long-sleeved shirt and pants. Avoid contact with bare skin. Do not use in confined areas. Do not use in areas where children or animals are present. Do not use on crops or animals for which it is not labeled. Do not use on crops or animals for which it is not labeled. Do not use on crops or animals for which it is not labeled.

Use 1 1/2 to 2 1/2 pounds for cucumber control. On clovers in Texas use 2 pounds for clover leaf weevil. To avoid possible injury or tender forage do not apply when forage is wet or when rain or excessive humidity is expected during the next 2 days.

ASPARAGUS
Use 1 1/2 to 2 1/2 pounds for asparagus beetle on seedlings or spears.
Use 2 1/2 to 3 1/2 pounds for asparagus beetle on spears and Asparagus ferns. Do not use on Asparagus ferns during the growing period. Do not apply to ferns after the first 3 to 4 weeks.
Do not apply within 3 days of harvest.

BEANS
Use 1 1/2 to 2 1/2 pounds for bean beetle on seedlings or pods.
Use 2 1/2 to 3 1/2 pounds for bean beetle on pods and Asparagus ferns. Do not use on Asparagus ferns during the growing period. Do not apply to ferns after the first 3 to 4 weeks.
Do not apply within 3 days of harvest.

CABBAGE, BRUSSELS SPROUTS, SPRING ONION, CAULIFLOWER AND KONJAK
Use 1 1/2 to 2 1/2 pounds for cabbage root maggot, cabbage root fly, and cabbage root weevil.
Use 2 1/2 to 3 1/2 pounds for cabbage root maggot, cabbage root fly, and cabbage root weevil.
Do not apply within 14 days of harvest.

CHINESE CABBAGE, SQUASHES, HANGOVER SALAD, NOPSEADISH, RALE, MUSTARD GREENS, RADISHES, RUTABAGA AND TURNIPS
Use 1 1/2 to 2 1/2 pounds for flea beetle and leafhoppers.
Use 2 1/2 to 3 1/2 pounds for flea beetle and leafhoppers.
Do not apply within 14 days of harvest of Chinese cabbage, rutabaga, and turnips.
Do not apply within 14 days of harvest of Chinese cabbage, rutabaga, and turnips.

CARROTS, FANSHIPS, PARSLEY
Use 1 1/2 to 2 1/2 pounds for flea beetle and leafhoppers.
Use 2 1/2 to 3 1/2 pounds for flea beetle and leafhoppers.
Do not apply within 14 days of harvest of carrot.
Do not apply within 14 days of harvest of carrot.

CORN (field, sweet and pop)
Use 1 1/2 to 2 1/2 pounds for European cornborer, corn earworm, and corn rootworm.
Use 2 1/2 to 3 1/2 pounds for European cornborer, corn earworm, and corn rootworm.
Do not apply within 14 days of harvest of corn.
Do not apply within 14 days of harvest of corn.

COTTON
Use 1 1/2 to 2 1/2 pounds for early and late instars of bollworm, tobacco budworm, and cotton rootworm.
Use 2 1/2 to 3 1/2 pounds for early and late instars of bollworm, tobacco budworm, and cotton rootworm.
Do not apply within 14 days of harvest of cotton.
Do not apply within 14 days of harvest of cotton.

CUCUMBER, MELONS, PUMPKIN AND SQUASH
Use 1 1/2 to 2 1/2 pounds for cucumber beetle, squash vine borer, and squash bug.
Use 2 1/2 to 3 1/2 pounds for cucumber beetle, squash vine borer, and squash bug.
Do not apply within 14 days of harvest of cucumber.
Do not apply within 14 days of harvest of cucumber.

DANDELION, ENDIVE (ESCAROLE), LETTUCE AND SALADY
Use 1 1/2 to 2 1/2 pounds for flea beetle and leafhoppers.
Use 2 1/2 to 3 1/2 pounds for flea beetle and leafhoppers.
Do not apply within 14 days of harvest of dandelion, endive, lettuce, and salad.
Do not apply within 14 days of harvest of dandelion, endive, lettuce, and salad.

FORAGE GRASSES AND PASTURE
Use 1 1/2 to 2 1/2 pounds for armyworm grasshopper, grasshopper, and grasshopper.
Use 2 1/2 to 3 1/2 pounds for armyworm grasshopper, grasshopper, and grasshopper.
Do not apply within 14 days of harvest of forage grasses and pasture.
Do not apply within 14 days of harvest of forage grasses and pasture.

GARDEN BEET, SPINACH AND SWISS CHARD
Use 1 1/2 to 2 1/2 pounds for flea beetle and leafhoppers.
Use 2 1/2 to 3 1/2 pounds for flea beetle and leafhoppers.
Do not apply within 14 days of harvest of garden beet, spinach, and Swiss chard.
Do not apply within 14 days of harvest of garden beet, spinach, and Swiss chard.

ORRA
Use 1 1/2 to 2 1/2 pounds for flea beetle and leafhoppers.
Use 2 1/2 to 3 1/2 pounds for flea beetle and leafhoppers.
Do not apply within 14 days of harvest of orra.
Do not apply within 14 days of harvest of orra.

PEANUTS
Use 1 1/2 to 2 1/2 pounds for peanut rootworm.
Use 2 1/2 to 3 1/2 pounds for peanut rootworm.
Do not apply within 14 days of harvest of peanuts.
Do not apply within 14 days of harvest of peanuts.


PEAS
Use 1 1/2 to 2 1/2 pounds for pea beetle and leafhoppers.
Use 2 1/2 to 3 1/2 pounds for pea beetle and leafhoppers.
Do not apply within 14 days of harvest of peas.
Do not apply within 14 days of harvest of peas.

POPPERS
Use 1 1/2 to 2 1/2 pounds for popper beetle and leafhoppers.
Use 2 1/2 to 3 1/2 pounds for popper beetle and leafhoppers.
Do not apply within 14 days of harvest of poppers.
Do not apply within 14 days of harvest of poppers.

RUTABAGA
Use 1 1/2 to 2 1/2 pounds for flea beetle and leafhoppers.
Use 2 1/2 to 3 1/2 pounds for flea beetle and leafhoppers.
Do not apply within 14 days of harvest of rutabaga.
Do not apply within 14 days of harvest of rutabaga.

SQUASH
Use 1 1/2 to 2 1/2 pounds for cucumber beetle, squash vine borer, and squash bug.
Use 2 1/2 to 3 1/2 pounds for cucumber beetle, squash vine borer, and squash bug.
Do not apply within 14 days of harvest of squash.
Do not apply within 14 days of harvest of squash.

181-247 1013 1453



CARBARYL INSECTICIDE


ACTIVE INGREDIENTS:
Carbaryl (1-naphthyl methylcarbamate) ... 60%
INERT INGREDIENTS ... 40%
TOTAL ... 100%

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

If small and scattered populations of insects are present, use 1 1/2 to 2 1/2 pounds per acre. If large and dense populations are present, use 2 1/2 to 3 1/2 pounds per acre. Do not apply when forage is wet or when rain or excessive humidity is expected during the next 2 days. Do not apply within 14 days of harvest of crops or animals for which it is not labeled.



BORNGRASS (such as milo grass, sorghum, sweet sorghum and hybrids)
Use 1 1/2 to 2 1/2 pounds for armyworm, corn earworm, grasshopper and weevils.
Use 2 1/2 to 3 1/2 pounds for armyworm, corn earworm, grasshopper and weevils.
Do not apply within 14 days of harvest of borngrass.
Do not apply within 14 days of harvest of borngrass.

SOYBEANS
For flight to moderate populations in Southeastern states only.
Use 1 1/2 to 2 1/2 pounds for velvetbean caterpillar, grasshopper, and leafhopper.
Use 2 1/2 to 3 1/2 pounds for velvetbean caterpillar, grasshopper, and leafhopper.
Do not apply within 14 days of harvest of soybeans.
Do not apply within 14 days of harvest of soybeans.

SUGAR BEETS
Use 1 1/2 to 2 1/2 pounds for armyworm, flea beetle, leafhopper and weevil.
Use 2 1/2 to 3 1/2 pounds for armyworm, flea beetle, leafhopper and weevil.
Do not apply within 14 days of harvest of sugar beets.
Do not apply within 14 days of harvest of sugar beets.

TOBACCO IN PLANT BEDS
Use 1 1/2 to 2 1/2 pounds for armyworm, flea beetle, leafhopper and weevil.
Use 2 1/2 to 3 1/2 pounds for armyworm, flea beetle, leafhopper and weevil.
Do not apply within 14 days of harvest of tobacco in plant beds.
Do not apply within 14 days of harvest of tobacco in plant beds.

SAVIT granules should be used in a ratio of 10 parts of granules to 100 parts of water. Granules should be used in a ratio of 10 parts of granules to 100 parts of water. Granules should be used in a ratio of 10 parts of granules to 100 parts of water.

For aircraft application: For pre-harvest spraying of storage tanks, tanks are to be sprayed with a mixture of 10 parts of granules to 100 parts of water. For pre-harvest spraying of storage tanks, tanks are to be sprayed with a mixture of 10 parts of granules to 100 parts of water.

For application: For application of granules to storage tanks and aircraft, tanks are to be sprayed with a mixture of 10 parts of granules to 100 parts of water. For application of granules to storage tanks and aircraft, tanks are to be sprayed with a mixture of 10 parts of granules to 100 parts of water.

COMPATIBILITY

1. Do not mix SAVIT with other pesticides unless your experience indicates that the mixture is effective and safe. If uncertain, prepare a small amount of the mixture and test it on a small area of the crop before applying it to the whole crop.

GENERAL CAUTION

- 1. Do not apply to crops which are to be grazed by livestock.
- 2. Do not apply to crops which are to be used for animal feed.

IMPORTANT PRE-HARVEST AND GRAZING USE INFORMATION AND LIMITATIONS

Do not apply to crops which are to be grazed by livestock. Do not apply to crops which are to be used for animal feed. Do not apply to crops which are to be used for animal feed.

DIRECTIONS FOR USE

Apply to crops which are to be grazed by livestock. Apply to crops which are to be used for animal feed. Apply to crops which are to be used for animal feed.

TOBACCO FIELD AND PLANT BED PEST CONTROL

Apply to tobacco fields and plant beds. Apply to tobacco fields and plant beds. Apply to tobacco fields and plant beds.

PEANUTS

Apply to peanut fields. Apply to peanut fields. Apply to peanut fields.

COTTON

Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre.

Use 1 1/2 to 2 1/2 pounds per acre. Use 1 1/2 to 2 1/2 pounds per acre. Use 1 1/2 to 2 1/2 pounds per acre.

Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre.

Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre.

CUCUMBER

Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre.

Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre.

Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre.

Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre.

Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre.

Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre.

Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre.

Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre.

Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre.

Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre.

Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre.

Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre.

Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre.

Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre.

Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre.

Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre.

Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre.

CAUTION

STATEMENT OF PRACTICAL TREATMENT

If swallowed, drink 1 or 2 glasses of water to induce vomiting. Do not induce vomiting or give anything by mouth to an unconscious person. Seek medical attention.

If on skin, remove contaminated clothing. Wash skin with soap and water.

If in eyes, flush with plenty of water. Do not rub eyes. Seek medical attention.

See label page for additional precautions and first aid instructions.



GRIFFIN CORPORATION
VALDOSTA, GEORGIA 31601
NET CONTENTS

10.5 LBS. (4.75 KG.) 264
10.5 LBS. (4.75 KG.) 264

BURGHUMS (such as milo, grain sorghum, sweet sorghum and hybrids) Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre.

SOYBEANS Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre.

TOBACCO Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre.

PEANUTS Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre.

SUGAR BEETS Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre.

GRASSHOPPERS Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre.

TOBACCO IN PLANT BEDS Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre.

PEANUTS Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre.

SUGAR BEETS Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre.

GRASSHOPPERS Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre.

TOBACCO IN PLANT BEDS Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre.

PEANUTS Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre.

SUGAR BEETS Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre.

GRASSHOPPERS Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre. Use 1/2 to 1 1/2 pounds per acre.

BEST AVAILABLE COPY

may not be used as a foliar insecticide. Do not use for control of insects on plants in the Recovery Act Disposal Category II (Disposal of plants may be controlled by approved Federal, State or local agencies under the Recovery Act provisions of the Recovery Act).

GENERAL INFORMATION

SAVIT BUV is a wettable powder for dispersion in water and application as an insecticidal spray or hydraulic blowers, mist blowers, low-gallonage ground equipment and aircraft. USE IN STRICT ACCORDANCE WITH LABEL DIRECTIONS AND LIMITATIONS.

SPECIAL DIRECTIONS AND PREPARING LOW-GALLONAGE SPRAYS

In low-gallonage ground equipment use the best of the following directions. Mix the concentrate in a bucket or line-type number 30 tank. Use clean equipment with at least 1/2 of the required water volume in the spray tank. Close off (or the start pump and maintain) the desired flow rate of water while slowly adding the proper amount of SAVIT BUV. Add remaining water and continue to mix the material in suspension. Then either apply using at least 3 gallons per acre of the prepared spray. After spraying, flush tank and lines with clean water.

For aircraft application of the prepared spray, the application rate should be above 100 gpm to maintain uniform suspension. Recirculate at intervals and before pouring into aircraft tank. For best results, use low-gallonage equipment and apply at least 3 gallons per acre of the prepared spray. After spraying, flush tanks and lines with clean water.

Combination spray mixtures in low-gallonage equipment and aircraft. Physical compatibility of SAVIT BUV with other pesticides in low-gallonage mixtures is not fully known. Before preparing large amounts, combinations should be tested by fully dissolving SAVIT BUV, then adding the other pesticide and the remaining portion of water. DO NOT USE MIXTURES THAT FORM A SOLID OR PRECIPITATE. Incompatibility will reduce effectiveness, may cause excessive phytotoxicities or plant injury. Use caution when applying on alternate spray mixtures. Under adverse growing conditions, combinations of carbaryl with some phosphate pesticides may cause foliar injury.

COMPATIBILITY

1. SAVIT BUV has been effective when used with the following: insecticides, insecticides, insecticides, fungicides, miticides, herbicides, growth regulators, and other pesticides used on citrus.
2. Compatibility can be affected by water hardness, various surfactants or other materials and ingredients. Possible plant injury may occur under adverse conditions such as high humidity, drought or improper fertilization, etc.
3. Control of SAVIT BUV with different substances and other materials is less effective. It states that the mixture is effective. To know the result of plant injury, if uncertain, prepare a small amount of the mixture and test it on a separate location of the crop.
4. Incompatible with: 1. alkalis. Not effective if used with alkaline materials such as bicarbonates, lime, lime sulfur, and calcium sprayers.

GENERAL CAUTION

1. To avoid possible phytotoxicity, do not apply when foliage is wet or when rain or excess humidity is expected during the next 2 days.
2. Does not control spider mites and is incompatible with most acaricidal pesticides.

IMPORTANT PRE-HARVEST AND GRAZING USE INFORMATION AND LIMITATIONS

Two uses are listed under the Federal Food, Drug, and Cosmetic Act: pre-harvest and grazing. Pre-harvest uses are for CARBAR and for CARBAR. CARBAR is used in all states with different pre-harvest and grazing limitations. CARBAR is used in all states with different pre-harvest and grazing limitations. Where no pre-harvest or grazing time limitations are stated, application can be made to the day of harvest and crops can be grazed without time delay if direction for use on the crop are followed.

For CARBAR, insecticide is used in all states with label directions for pre-harvest and grazing. For CARBAR, insecticide is used in all states with label directions for pre-harvest and grazing. For CARBAR, insecticide is used in all states with label directions for pre-harvest and grazing. For CARBAR, insecticide is used in all states with label directions for pre-harvest and grazing.

DIRECTIONS FOR USE

IMPORTANT: The directions on this label are based on tests and field experience and are intended for general use and to provide a guide to plants and animals and to provide a guide to plants and animals. Follow directions for most effective use.

FORAGE, FIELD AND VEGETABLE CROP INSECT CONTROL

For optimum disease control, apply 1 to 2 pounds of SAVIT BUV per acre. Where a dosage range is indicated, use lower rate on young plants, higher rate on mature plants. Use sufficient spray volume for 100% full coverage. This will usually vary from 3 to 40 gallons per acre depending on equipment, degree of insect infestation, and stage of crop growth. Keep spray nozzles well regulated. Apply when insects are first seen appear and repeat at 7 to 14 day intervals or as necessary unless other intervals are specified below. See specific directions below for grasshopper control.

ALFALFA AND CLOVERS

Use 4 to 10 pounds for beetle beetles and Mexican bean beetle.
Use 1 to 2 pounds for alfalfa caterpillar, leaf beetles, leaf beetles, green leaf beetles, Japanese beetle, leafhoppers, thrips, and velvetbean caterpillar.
Use 1 to 1 1/2 pounds for armyworm, corn earworm, stink bug, and webworm.
Use 1 to 2 pounds (Western United States) and 2 pounds (Eastern United States) for control of alfalfa weevil.

CORN (field, sweet and pop)

Use 2 to 2 1/2 pounds for fall armyworm, corn rootworm adults, European corn borer, fall armyworm, Japanese beetle, leaf beetles and leafhoppers. For larvae of corn rootworm, same levels apply to the field and sweet corn. Repeat as necessary. For insects attacking silks and ears, application is starting when first silks appear and continuing until silks begin to dry. Three or more may be required depending on severity of infestation. Timing and good coverage are absolutely critical.

Use 2 to 4 pounds (at least 1 1/2 gallons of water for 100 square feet) for fall armyworm control. Spray in 12 inch band or to insure adequate stalk coverage. Use of sufficient amounts for narrower bands or broad application of SAVIT BUV to the base region of corn during the pollen shed period will control populations.

COTTON

Use 1 to 1 1/2 pounds for early season thrips, flea beetles, cotton leaf-opper, striped blister beetle, and webworm.

Use 1 to 2 1/2 pounds (depending on the size of the cotton and insect population level) for pest in bollworm, bollworm, fall armyworm, cotton leafperforator, leafhoppers, leaf-opper, plant bug, and for light to moderate infestations of lygus bugs in Western irrigated cotton. Should be made every 5 to 7 days as long as control is necessary. For improved bollworm control of bollworm, molasses feed grade, later use in a spray volume not to exceed 20 gallons avoid staining and do not use molasses after bolls open.

Use 1 to 1 1/2 pounds for pine rootworm on a 5 to 7 day schedule depending on the size of degree of infestation.

Use 2 1/2 pounds for stink bugs and saltmarsh caterpillar.

APHID populations will be suppressed by repeated applications of this insecticide. May be applied.

COWPEAS

Use 1 to 1 1/2 pounds for beetle beetles and Mexican bean beetle.

Use 1 to 1 1/2 pounds for alfalfa caterpillar, bean leaf beetle, cucumber beetle, flea beetles, green Japanese beetle, leafhoppers, thrips, and alfalfa hopper, thrips and velvetbean caterpillar. Use 1 to 1 1/2 pounds for armyworm, corn earworm, cutworm, stink bugs and webworm. Use 2 1/2 pounds for cowpea curculionid. Apply 4 applications beginning at first bloom and thereafter.

In California use 2 1/2 pounds for corn earworm, lima bean pod borer, lygus and stink bugs. To avoid possible injury on tender foliage, do not apply when foliage is wet or when rain or excess humidity is expected during the next 2 days.

CUCUMBER, MELONS, PUMPKIN AND SQUASH

Use 1 to 1 1/2 pounds for cucumber beetles, flea beetles, leafhoppers and squash bug. Avoid phytotoxicity. To avoid possible injury on tender foliage, do not apply when foliage is wet or when excessive humidity is expected during the next 2 days.

DANDELION, ENDIVE (ESCAROLE), LETTUCE AND SALSFY

Use 1 to 1 1/2 pounds for flea beetles, flea beetle, henequen bug and leafhoppers. Use 1 to 1 1/2 pounds in a 5 to 7 day schedule for aster leafhopper. Use 1 to 2 1/2 pounds for armyworm, imported cabbageworm, corn earworm, tarnished plant bug. For optimum control of corn earworm after lettuce heads begin to form use 2 1/2 pound schedule. To avoid possible injury on lettuce, do not apply when foliage is wet or when humidity is expected during the next 2 days.

Do not apply within 3 days of harvest of head endive and salsify roots. Do not apply within 14 days of harvest of dandelion, endive (escarole), leaf lettuce and salsify.

FORAGE GRASSES AND PASTURE

Use 1 to 1 1/2 pounds for armyworm and thrips. For thrips control in grasses grown for seed pressure may help penetration into fruit. Use 2 to 2 1/2 pounds for wire grubs (green June bug).

GARDEN BEET, SPINACH AND SWISS CHARD

Use 1 to 1 1/2 pounds for flea beetles, henequen bug and leafhoppers. Use 1 to 1 1/2 pounds in a 5 to 7 day schedule for aster leafhopper. Use 1 to 2 1/2 pounds for armyworm, corn earworm, tarnished plant bug and stink bugs. Do not apply within 3 days of harvest of garden beetle tops, spinach and Swiss chard. Do not apply within 14 days of harvest of garden beetle tops, spinach and Swiss chard.

OKRA

Use 1 to 2 1/2 pounds for corn earworm and stink bug. Repeat 5 to 7 day intervals or as necessary.

PEANUTS

Use 1 to 1 1/2 pounds for beetle beetles and Mexican bean beetle. Use 1 to 1 1/2 pounds for alfalfa caterpillar, leaf beetles, leaf beetles, cucumber beetle, green Japanese beetle, leafhoppers, thrips, and alfalfa hopper, thrips and velvetbean caterpillar. Use 1 to 1 1/2 pounds for armyworm, corn earworm, stink bug and webworm. To avoid possible injury on tender foliage, do not apply when foliage is wet or when rain or excess humidity is expected during the next 2 days.

PEAS

Use 1 to 2 pounds for leafhoppers and Colorado potato beetle. Use 1 to 1 1/2 pounds for armyworm. Use 3 pounds for alfalfa caterpillar in Washington State only. Repeat as necessary.

POTATO, TOMATO, EGGPLANT AND PEPPER

Use 1 to 1 1/2 pounds for European corn borer, fall armyworm, flea bugs, tomato fruitworm, corn earworm, tarnished plant bug and stink bug. Use 2 1/2 pounds for cutworm control in potato, tomato and eggplant.

RICE

In the Mississippi Delta and Texas use 1 to 1 1/2 pounds for armyworm and stink bug. In California use 2 1/2 pounds for armyworm, grasshoppers and tadpole shrimp. For optimum control, apply to the water when the pest first appears. **WARNING:** Avoid injury to rice. Do not apply preplant (such as Stem F 3 or Rogue before or after SAVIT BUV application). Do not apply within 14 days of harvest.



ACTIVE INGREDIENTS:
 Carbaryl (1-naphthyl methycarbamate) ... 80%

INERT INGREDIENTS: ... 20%

TOTAL ... 100%

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

If swallowed, drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

If in eyes, flush with plenty of water for at least 15 minutes. For further treatment get medical attention.

See side panel for additional precautionary statements

GRIFFIN CORPORATION
 VALDOSTA, GEORGIA 31601
 NET CONTENTS ...

EPA REG. NO. 1812 264
 EPA EST. NO. 1812 GA 3

SORGHUMS (such as milo, grain sorghum, sweet sorghum and hybrids):
 Use 1 1/4 to 1 1/2 pounds for armyworm, corn earworm, stink bugs and webworms.
 Use 1 1/2 pounds for sorghum midge.
 Use 2 1/2 pounds for cutworms. For optimum insect control, in grain sorghum, direct spray into the tassel.
 Needs: Treat for sorghum midge 3 to 4 days after heads have emerged from boot.
 Do not apply within 21 days of harvest of grain.
 No time limitation on sorghums used for forage.

SOYBEANS
 For light to moderate populations in Southeastern states only
 Use 1/2 pound for velvetbean caterpillar, green cloverworm, bean leaf beetle, cucumber beetle and Mexican bean beetle.
 Use 1/2 to 1 pound for cutworm.
 For cleanup of existing populations:
 Use 1/4 to 1/2 pounds for beetle beetles and Mexican bean beetle.
 Use 1 1/2 pounds for alfalfa caterpillar, leaf beetles, cucumber beetles, green cloverworm, Japanese beetle, leafhoppers, three-cornered alfalfa leafhopper, thrips and velvetbean caterpillar.
 Use 1 1/2 to 2 1/2 pounds for armyworm, corn earworm, webworm, stink bug and cutworm.
WARNING! Do not apply to alfalfa or alfalfa hay. Do not apply to alfalfa or alfalfa hay used for livestock feed.
 Butoxone: S-1 to soybeans.

SUGAR BEETS
 Use 1/2 to 2 1/2 pounds for armyworm, flea beetle, leaf beetles and webworms.
 Do not apply within 14 days of harvest.

TOBACCO IN PLANT BEDS
 For tobacco flea beetle: Use 2 1/2 level tablespoons per gallon or 1 1/2 pounds per 50 gallons, and apply 6 gallons per 100 square yards.
 For green June beetle grub: Use 1/2 pound per 100 gallons of water, when insects in their larval stage appear.
 Apply only to areas that larvae have penetrated by sprinkling as a trench with 50 to 100 gallons per 100 square yards. May be applied before or after seeding. Avoid excessive application as plant injury may result.

IN FIELDS
 For budworm, flea beetles, Japanese beetle, June beetles and hornworms: Use 1 1/4 to 2 1/2 pounds per acre in sufficient water for full coverage of plants. Use lower rate on young plants up to knee high; higher rate on mature plants. Use at least 10 gallons per acre of the prescribed spray. Begin treatments when worms are small. Avoid excessive application in bud of plants as injury may result.
 Treated fields may be entered immediately after foliage has dried.
 If late applications are necessary, allow 3 days before pruning or cutting.

GRASSHOPPERS
 Apply 1/2 to 1 1/2 pounds as often as necessary TO CONTROL GRASSHOPPERS IN THE ABOVE FORAGE FIELD AND VEGETABLE CROPS. FOLLOW PRE-HARVEST AND TREATMENT LIMITATIONS LISTED FOR EACH OF THE ABOVE CROPS. A 1/2 to 1 1/2 pounds rate is suggested for nymphs on small plants or

Use Savit 80W ...
 Do not apply when ...

TREE FRUIT AND NUT INSECT CONTROL

Refer to the following for information on SAVIT 80W: 100 per cent carbaryl dust. Apply dust at rate of 10 pounds per acre. Do not apply when foliage is wet or when rain or excessive humidity is present.

ALMOND
 Use 1 1/2 to 2 pounds for almond weevil, peach twig borer, fruit fly, and other insects. Do not apply within 28 days of harvest.

APPLES AND PEARS
 Use 1 1/2 to 2 pounds for apple maggot, European spruce sawfly, codling moth, apple rust, and other insects. Do not apply within 1 day of harvest of apples.

APPLE THINNING
 Many factors influence the degree of apple thinning obtained with SAVIT 80W. Factors such as weather, soil, and fertilizer are important. Do not apply when foliage is wet or when rain or excessive humidity is present.

CITRUS FRUITS (such as grapefruit, lemons, limes, oranges, tangerines, and citrus hybrids):
 Use 1 1/2 to 2 pounds for California orange and citrus thrips, fruit fly, and other insects. Do not apply within 5 days of harvest of these crops.

PEACHES, APRICOTS AND NECTARINES
 Use 1 1/2 to 2 pounds for peach weevil, European sawfly, and other insects. Do not apply more than 10 pounds of SAVIT 80W per acre.

PEACANS
 Use 1 1/2 to 2 pounds for pecan weevil and pecan nut casebearer. Do not apply more than 9 pounds of SAVIT 80W per acre.

PLUMS, PRUNES AND CHERRIES
 Use 1 1/2 to 2 pounds for plum aphid, cherry maggot, and other insects. Do not apply within 1 day of harvest of these crops.

WALNUT
 Use 1 1/2 to 2 pounds for walnut leafhopper, European fruit lecanium, and other insects. Do not apply within 1 day of harvest of these crops.

SMALL FRUIT INSECT CONTROL

BLUEBERRIES
 Use 1 1/2 to 2 pounds for blueberry maggot, cherry and cranberry fruitworm, and other insects. Do not apply within 1 day of harvest.

CRANBERRIES
 Use 1 1/2 to 3 1/2 pounds for cutworm, cranberry fireworm, and other insects. Do not apply within 1 day of harvest.

GRAPES
 Use 1 1/2 to 2 1/2 pounds for European fruit lecanium, grape leafhopper, grapefruit scale, and other insects. Do not apply within 1 day of harvest.

STRAWBERRIES
 Use 1 1/2 to 2 1/2 pounds for meadow springtail, strawberry leafhopper, and other insects. Do not apply within 1 day of harvest.

