

1812-269 PM 21 1 of 1

PRECAUTIONARY STATEMENTS HAZARDOUS TO HUMANS & DOMESTIC ANIMALS DANGER — PELIGRO

DANGER: Fatal if inhaled. Corrosive; causes irreversible eye damage. May be harmful or fatal if swallowed or absorbed through the skin. Do not get in eyes, on or on skin. Do not breathe dust, vapor, or spray mist. When handling either drums containing TPTH or spray treated materials, wear protective clothing: long pants, long sleeve shirt, impermeable gloves, hat, boots and a pesticide respirator jointly approved by the Mining Enforcement and Safety Administration and the National Institute for Occupational Safety and Health. When handling the concentrated product, wear a face shield. Wash thoroughly with soap and water after handling and before eating or smoking. Remove contaminated clothing and wash before reuse. Do not enter treated areas for at least 24 hours after treatment.

The United States Environmental Protection Agency has determined that triphenyltin hydroxide causes birth defects in laboratory animals. Exposure to triphenyltin hydroxide during pregnancy should be avoided.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and wildlife. Do not apply directly to water or wetlands. Drift or runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Cover or incorporate spills. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with the labeling.

RE-ENTRY STATEMENT

Do not apply this pesticide in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing until sprays have dried. Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product.

DANGER: Do not enter treated areas without appropriate protective clothing for 24 hours after application. In case of accidental exposure, contact your nearest Poison Control Center.

Pro-Tex™ is a non-flammable, flowable fungicide.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the hazardous waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple Rinse (or equivalent). Then recycle for recycling or incineration, or put in place and dispose of in a sanitary landfill, or in incineration, or if allowed by state and local authorities, by burning. Do not reuse.

Pro-Tex™ Fungicide can be applied as a ground or aerial spray to control fungal infestations on broad crops. Application rate is for general use. For agricultural extension or agricultural experiment station specialists should be consulted for specific applications and timing recommendations. With any spray application through nozzles, essential for good control.

Apply this product only as specified on this label.

RESTRICTED USE PESTICIDE

Because of the high acute toxicity of triphenyltin hydroxide and its potential for creating teratogenic effects, this product may be applied only by certified applicators or persons directly under their supervision.

PRO-TEX™

ACCEPTED

JUN 26 1985

Federal Insecticide,
Fungicide, and Rodenticide Act

Registered under
EPA Reg. No. 1812-269

ACTIVE INGREDIENTS:

Maneb (Manganese Ethylenebisdithiocarbamate) 32.63%

Triphenyltin hydroxide 4.72%

INERT INGREDIENTS 62.65%

TOTAL 100.00%

This product contains 3.5 lbs. of Maneb.

This product contains 0.5 lbs. of TPTH.

KEEP OUT OF REACH OF CHILDREN

DANGER — PELIGRO



POISON

IF ON SKIN: Wash with plenty of soap and water.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

IF IN EYES: Flush with plenty of water. Call a physician.

IF SWALLOWED: Do not induce vomiting. Drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Call a physician or Poison Control Center.

PRECAUCION AL USUARIO: Sustenido no lee ingles, no use este producto hasta que le etiqueta haya sido explicado ampliamente.

See Side Panel For Additional Precautionary Statements



GRIFFIN CORPORATION
VALDOSTA, GEORGIA 31601

EPA Reg. No. 1812 269

EPA Est. No. 1812 GA 3

Net Contents 2 1/2 gallons

GENERAL INSTRUCTIONS FOR APPLICATION

POTATOES: Early and Late Blight

Ground Application: Apply 0.75 - 1.6 qts of Pro-Tex™ per acre in at least 15 gallons of water. Full coverage of the foliage is necessary for the best results. Diluted spray should be directed uniformly to all parts of the plant and the gallonage increased according to the size of the plant.

Aerial Application: Apply 0.75 - 1.0 qts. of Pro-Tex™ per acre in 3 - 10 gallons of water. For helicopter application fly high enough so as not to whip the vines.

Applications should begin with the appearance of blight weather conditions and continue on a seven (7) day schedule. The lower rate of application should be used early in the season and the high rate mid to late season or when blight infection is in the area. Do not apply within 7 days of harvest.

SUGAR BEETS: Cercospora Leafspot

Ground Application: Apply 1.0 - 2.0 qts of Pro-Tex™ per acre in at least 15 gallons of water. Full coverage of the plant is necessary for best results.

Aerial Application (helicopter or fixed wing aircraft): Apply 1.0 - 2.0 qts. of Pro-Tex™ per acre in 5 - 10 gallons of water.

Use lower gallonage when plants are small and increase volume with plant size. Use the lower rate for protective sprays and the higher rates later in the season or during high infection periods. Application should begin when leafspot conditions appear or when the disease is in the area. Repeat at 10-14 day intervals. Do not treat within 14 days of harvest. Do not graze or feed beet tops to livestock.

BEST AVAILABLE COPY
WARRANTY STATEMENT

GRIFFIN warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of GRIFFIN. In no case shall GRIFFIN be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. GRIFFIN MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

EPA 17263
USE 2/186