



**KOCIDE® SD**  
**SEED DRESSING**  
**AGRICULTURAL FUNGICIDE**

<b>ACTIVE INGREDIENT</b>	
Cupric Hydroxide . . . . .	30%
<b>INERT INGREDIENTS . . . . .</b>	<u>70%</u>
<b>TOTAL . . . . .</b>	100%

Contains 3.2 Lbs. Cupric Hydroxide Per Gallon  
(Metallic Copper Equivalent 19.4%)

**KEEP OUT OF REACH OF CHILDREN**  
**DANGER - PELIGRO**  
**STATEMENT OF PRACTICAL TREATMENT**

**IF IN EYES:** Hold eyelids open and flush with water for 15 minutes. Get medical attention.

**IF SWALLOWED:** Drink promptly a large quantity of milk, egg white, gelatin solution, or if these are not available, large quantities of water. Avoid alcohol. Call a physician or Poison Control Center.

**IF ON SKIN:** Wash with plenty of soap and water. Get medical attention if irritation persists.

**IF INHALED:** Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate use of gastric lavage.

**PRECAUCION AL USUARIO:** Si usted no lee ingles, no use este producto hasta que la etiqueta haya sido explicado ampliamente.

Griffin Corporation  
Valdosta, Georgia 31601

2 1/2 Gallons

U.S. Patent No. 3,428,731

EPA Reg. No. 1812-301  
EPA Est. No. 8901-TX-1

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
DANGER - PELIGRO**

Corrosive. Causes irreversible eye damage. Harmful or fatal if swallowed. Harmful if absorbed through skin or inhaled. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Do not get in eyes, skin or clothing. Avoid breathing vapors. Wear goggles or face shields. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

**ENVIRONMENTAL HAZARDS**

This pesticide is toxic to fish and aquatic organisms. Do not apply directly to water. Runoff from treated areas may be hazardous to fish and aquatic organisms in adjacent aquatic sites. Do not contaminate water when disposing of equipment washwaters.

**GENERAL INSTRUCTIONS**

KOCIDE SD should be used as described below. KOCIDE SD is specially formulated for use as a seed protectant. It is compatible with many fungicides and other materials commonly used as seed disinfectants. Because of the wide number of combinations possible with other pesticides, it is recommended that specific mixtures be evaluated before use with this product.

The following specific instructions are based on general applications. The recommendations of State Agricultural Experiment Stations should be closely followed.

**BEST AVAILABLE COPY**

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling, or reconditioning, or puncture and dispose of in a sanitary landfill, or incinerate, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**WORKER PROTECTION STATEMENT:** Do not apply this product in a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

<b>CROP</b>	<b>DISEASE</b>	<b>RATE/100 LBS OF SEED</b>	<b>USE INSTRUCTIONS</b>
Rice	Water Mold & Seed Rot	4-8 ounces	For ease of handling and when using a seed treating machine, dilute KOCIDE SD with an equal amount of water. Maintain continuous agitation of the mixture throughout the operation. (Phytotoxicity may occur in drill-seeded rice. Consult State Agricultural Experiment Station regarding specific recommendations for your area.)
Wheat & Barley	Bacterial Leaf Blight, Bacterial Leaf Streak, & Common Bunt	4 ounces	It should be diluted with equal parts of water before applying.

Treated seed should not be used for food, feed or oil purposes.

**BEST AVAILABLE COPY**

**WARRANTY STATEMENT**

GRIFFIN warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of GRIFFIN. In no case shall GRIFFIN be liable for consequential, special or indirect damages resulting from the use or handling of this product. **GRIFFIN MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**