


PM 35

1812-320

185



BEST AVAILABLE COPY

LINEX[®] 50 DF

DISPERSIBLE GRANULE

ACTIVE INGREDIENT:

Linuron 3-(3, 4-dichlorophenyl)-1-methoxy-1-methylurea 50%

INERT INGREDIENTS: 50%

TOTAL 100%

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Induce vomiting.

IF ON SKIN: Remove by washing with soap and water.

IF IN EYES: Wash with large amounts of water and get medical attention.

See Additional Precautionary Statements

Griffin

Griffin. The Hardworking Difference™

GRIFFIN CORPORATION

VALDOSTA, GA 31601

EPA REG. NO. 1812-320

2 of 5

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

Harmful if swallowed. May irritate eyes, nose, throat, and skin. Avoid breathing dust or spray mist. Avoid contact with skin, eyes, and clothing. In case of skin contact, wash with soap and water. In case of contact with eyes, immediately flush with copious amounts of water. Get medical attention if irritation persists. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or wetlands. Do not apply where runoff is likely to occur. Do not apply where weather conditions favor drift from areas treated. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

RE-ENTRY STATEMENT

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not re-enter treated areas for 24 hours following application unless protective clothing is worn.

Because certain states may require more restrictive entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. (Indicate specific oral warnings which inform workers of areas or fields that may not be entered without specific protective clothing, period of time field must be vacated and appropriate actions to take in case of accidental exposure). When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: CAUTION: Area treated with LINEX 50 DF on (day of application). Do not re-enter treated areas for 24 hours following application unless protective clothing is worn. In case of accidental exposure see Statement of Practical Treatment.

REQUIRED CLOTHING AND EQUIPMENT FOR MIXING/LOADING AND APPLYING LINURON

One piece overalls which have long sleeves and long pants constructed of finely woven fabric as specified in the USDA/EPA Guide to Commercial Applicators.

Wide brimmed hats.

Chemical resistant gloves which are mid-forearm to elbow length.

Any article worn while applying linuron must be cleaned before re-using. Clothing which has been drenched or has otherwise absorbed concentrated pesticide must be disposed of in a sanitary landfill, or incinerated, or if allowed by state and local authorities, by burning.

Instead of clothing and equipment specified above, the applicator can use an enclosed tractor cab which provides a filtered air supply (as described by Taschenberg and Bourke, 1975).

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product.

GENERAL INFORMATION

LINEX 50 DF is a dispersible granule to be mixed in water and applied as a spray for selective control of weeds on certain crops and for non-selective weed control on non-cropland areas. LINEX 50 DF is non-corrosive to equipment, non-flammable, and non-volatile.

To control susceptible weed seedlings for an extended period of time, apply LINEX 50 DF to soil before weed emergence. The degree of control and duration of effect will vary with the amount of chemical applied, soil texture, rainfall and other conditions. Higher dosages are needed for soils high in clay or organic matter. Soil high in clay or organic matter will require lower dosages to obtain equivalent herbicide performance. Since moisture is needed to activate LINEX 50 DF, rainfall or irrigation is needed within 24 hours of application.

When using LINEX 50 DF to control emerged weeds best results are obtained on succulent weeds growing in temperatures of 70° or higher with high humidity. When recommended, addition of a surfactant to the spray increases contact effects of LINEX 50 DF.

It is suggested that growers limit their first use to small areas as the effect of LINEX 50 DF varies with soils, uniformity of application and environmental conditions. Follow all label directions on this and any product used in mixtures.

DIRECTIONS

LINEX 50 DF should be used only in accordance with recommendations on this label. Do not apply this product through an irrigation system.

Injury to or loss of desirable trees or other plants may result from failure to observe the following:

Equipment should not be flushed or drained or LINEX 50 DF applied near desirable trees or other plants or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. LINEX 50 DF should not be used on ornamental shrubs or herbaceous plants, lawns, walks, driveway, tennis courts or similar areas. Keep drift of dry powder or spray from desirable plants.

Equipment should be cleaned of all traces of LINEX 50 DF immediately after use. Nozzle tips and screens should be removed and cleaned separately. Flush tank, pump, hoses and boom with several changes of water.

WEED CONTROL AND CROP USE

PREEMERGENCE USE — Germinating Weeds: LINEX 50 DF controls weeds as they germinate. LINEX 50 DF will control the following broadleaf weeds and grasses:

Broadleaf Weeds

Carpetweed
Chickweed
Dayflower, Common
Florida Pusley
Galinsoga

Goosefoot, Nettleleaf
Lambsquarters
Mustard
Pigweed

Purslane
Radish, Wild
Ragweed, Common
Smartweed

Grasses

Barnyardgrass (Watergrass)
Canarygrass
Crabgrass

Foxtail (Including Giant)
Goosegrass
Fall Panicum

LINEX 50 DF will provide partial control of the following:

Cocklebur Morningglory, Annual Prickly Sida (Teaweed) Sicklepod Velvetleaf (Buttonweed)

Established perennials such as bermudagrass, Canada thistle, field bindweed, johnsongrass and purple nutsedge will not be controlled. On lighter soils lower dosage rates will be effective and higher rates on heavier soils and the more resistant seedling weeds. Moisture in the form of rainfall or sprinkler irrigation is necessary after treatment to carry the chemical into the root zone of germinating weeds. Best results are obtained if moisture of 1/2" to 1" on moist soils; 1" to 2" on dry soils is applied within two weeks after application.

Crop injury may result if application of LINEX 50 DF is made to ground which is cloddy or compacted resulting in improperly planted seed. A good seed bed must be prepared before application. Seed should be planted to specified depth. Do not disturb or cultivate surface soil after application and before crop emergence as weed control will be reduced and crop injury may result. If there is not enough moisture to activate the herbicide, a shallow cultivation, preferably by rotary hoe, should be made after crop emergence while weeds are small enough to be controlled by mechanical means.

POSTEMERGENCE USE — Emergent Seedling Weeds: LINEX 50 DF will control the following weeds at recommended rates.

Broadleaf Weeds

Carpetweed
Cocklebur
Dayflower, Common
Dog Fennel
Fiddleneck (Amsinckia)
Florida Beggarweed
Florida Purslane (Florida Pusley)

Groundsel
Knapweed
Lambsquarters
Morningglory, Annual
Mustard
Nettleleaf
Goosefoot
Pigweed

Prickly Sida (Teaweed)
Purslane
Sesbania
Sicklepod
Smartweed
Velvetleaf (Buttonweed)
Wild Buckwheat

Grasses

Barnyardgrass (Watergrass)
Broadleaf (Signalgrass)
Canarygrass
Crabgrass
Fall Panicum
Foxtail (Including Giant)
Goosegrass

Rattail Fescue
Ryegrass, Annual
Texas Panicum

BEST AVAILABLE COPY

3 of 5

EQUIPMENT — SPRAY VOLUMES AND PRESSURES: A tractor-mounted fixed-boom power sprayer calibrated to a constant speed and rate of delivery should be used. Openings in screens should be equal to or larger than 50 mesh. To keep the material in suspension continuous agitation in the spray tank is necessary. Agitation can be by hydraulic or mechanical means. If a by-pass or return line is used, it should terminate at the bottom of the tank to minimize foaming. Overlapping should be avoided. Shut spray booms off while starting, turning, slowing or stopping or crop injury may result.

Use 20 to 60 gallons per acre and spray pressure of 30 to 40 psi for preemergence application.

For postemergence application for thorough coverage of weed foliage, use sufficient volume (minimum of 20 gallons per acre) and a spray pressure of 20 to 25 psi to keep spray drift to a minimum.

LINEX 50 DF is labeled for application using ground equipment only. Aerial applications are prohibited.

SPRAY PREPARATION: The tank should be 1/4 filled with water. Start agitation system, add **LINEX 50 DF** and continue adding water. Each component of any tank mix should be added separately while adding water. Surfactant, if used, should be added last when the tank is nearly full. Agitation should continue throughout. If poor mixing should occur with any component, premix the component with two parts water before adding to the spray tank.

Fertilizer Spray Mixtures: For preemergence application, nonpressure nitrogen or fertilizer solution may be used in the spray mixture unless otherwise directed. Small quantities should be tested for compatibility by the following procedure before full scale mixing:

1. Measure one pint of intended spray, water or fertilizer solution into a jar.
2. Check the pH of the liquid and adjust if necessary.
3. Add, in the given order, the intended ingredients, shaking well after each addition:
 - (a) Surfactants (spreaders), acidifiers, compatibility agents, and activators; add one teaspoon for each pint/100 gallons.
 - (b) Dry ingredients (wetttable powders, dry flowables); add one tablespoon for each pound/100 gallons.
 - (c) Soluble ingredients: add one tablespoon for each pound/100 gallons.
 - (d) Flowables: add one teaspoon for each pint/100 gallons.
 - (e) Soluble ingredients: add one tablespoon for each pound/100 gallons.
 - (f) Spreaders — stickers: one teaspoon for each pint/100 gallons.
4. The final mixture should be uniform and smooth with no evidence of coagulation occurring. If compatibility is evident, begin test again with a compatibility agent added first. Six drops is equivalent to four ounces per 100 gallons. If this does not smooth the mixture, try higher concentrations and other compatibility agents.
5. Allow the mixture to stand undisturbed thirty minutes. If separation occurs, shake and observe the resulting mixture.
 - (a) If it is smooth, then proceed with spraying.
 - (b) If not, do not attempt to spray the mixture.
 - (c) more compatibility agents
 - (d) different formulations
 - (e) different ingredients (switch from WP or EC to Flowable or from WP to EC)
 - (f) change active ingredients

Prepare spray by adding the fertilizer solution to the spray tank first if the mixture is compatible. Dilute **LINEX 50 DF** with water, then add the herbicide and add slowly to spray tank with agitator running. Follow directions on the tank mix.

USE RATES: All rates are based on broadcast rates. Use proportionately less for band treatment. Use the lower rate on heavy clay or organic matter (coarser soils) and the higher rate on soils high in clay or organic matter (fine soils). For postemergence application the lower rate should be used on smaller weeds and the higher rate on larger weeds.

SOIL LIMITATIONS: Do not use on sand, loamy sand, gravelly soils or exposed subsoils or soils containing less than 1% organic matter unless otherwise directed as injury to the treated or subsequent crop may result.

REPLANTING: The same crop may be replanted in soil treated preemergence with **LINEX 50 DF** if initial seedling fields to produce a stand. Soil should be reworked before replanting. Injury to the crop may result if field is retreated with a second application. Do not replant treated areas to any other crop within 4 months after last application unless otherwise stated as injury to subsequent crops may result. West of the Rocky Mountains carrots may be planted 4 months after the last application.

Do not rotate crops used for food or feed which are not registered for use with linuron into areas previously treated with this chemical.

CROP ROTATION RECOMMENDATIONS — Unless otherwise directed, any crop may be planted after 4 months except for cereals where only barley, oats, rye and wheat may be planted.

West of the Rocky Mountains: Carrots or celery may be planted 4 months after last application; do not plant any other crop until 1 year after the last application as injury may result.

ASPARAGUS — California, Michigan, Minnesota, North Carolina, Oregon, Washington only.

Direct Seeded or Newly Planted Crops: Do not exceed 4 lbs. total per acre per season; do not use surfactant or fertilizer solution in spray mixture.

Preemergence Application — Make a single application of 2 to 4 lbs. per acre after planting seed 1 1/2" deep in coarse soil and 1" deep in fine soils. During planting operation, spray activated charcoal as a 1" band on soil surface directly over seed rows at the rate of 300 lbs. per acre (equivalent to 15 lbs. per acre of soil surface with 20" row spacing). Preemergence weed control will be reduced in soil with high organic matter (greater than 5% and peat or muck).

Postemergence Application: Make 1 or 2 applications of 1 to 2 lbs. per acre when ferns are in 6" to 18" stage and weeds are not over 4" tall.

Established Beds: Do not apply within 1 day of harvest. Do not exceed 8 lbs. total per acre per season. Do not use surfactant or fertilizer solution in spray mixture. Do not re-enter treated areas for 24 hours following application unless protective clothing is worn.

Preemergence Application: Make a single application of 2 to 4 lbs. per acre. Preemergence weed control will be reduced in soils with high organic matter (greater than 5% and peat or muck).

Postemergence Application: Make 1 to 4 applications of 1 to 2 lbs. per acre before weeds exceed 4" in height. Apply before cutting season or immediately after cutting. Do not re-enter treated areas for 24 hours following application unless protective clothing is worn.

Directed Postemergence Application (Fern Stage): Make a single application of 4 to 8 lbs. per acre as directed; spray to base of plants for control of dudain melon.

Note: If more than 4 lbs. **LINEX 50 DF** per acre is applied per season, do not plant any other crop until 1 year after last application. Do not enter areas treated postemergence for 24 hours following application unless protective clothing is worn.

In California, dudain melon and annual nightshade are controlled.

BULBS — Tulip, Calla Lily, Daffodil and Dutch Iris Only — California: After planting of bulbs, settle the soil with sprinkler irrigation (rainfall will serve the same purpose); then before emergence of plants (bulbs), apply 2 lbs. **LINEX 50 DF** per acre in a minimum of 20 gallons of water per acre. Treat only once during the growing season.

NOTE: Unless otherwise directed, do not use on sand, loamy sand, gravelly soils, or exposed subsoils nor on soils containing less than 1% organic matter as injury to the crop may result. Do not re-enter treated areas for 24 hours following application unless protective clothing is worn.

CARROTS — Florida, Michigan, Ohio and Wisconsin — **Preemergence Application** — A single application of 1 to 3 lbs. per acre should be made after planting but prior to carrots emerging. Seed should be planted at least 1 1/2" deep. The lower rate should be used on lighter soils and the higher rate on heavier soils. Additional postemergence applications may be made as long as the total does not exceed 4 lbs. of **LINEX 50 DF** per acre per season.

NOTE: To prevent possible crop injury in various varieties of carrot, determine tolerance to **LINEX 50 DF** before adoption as a field practice.

Entire U.S. — Postemergence: Apply 1 1/2 to 3 lbs. per acre as a non-directed spray after carrots are at least 3" tall. Apply before annual grasses exceed 2" in height and before broadleaf weeds exceed 8" in height. Repeat application may be made but do not exceed 4 lbs. **LINEX 50 DF** per acre (West of Rocky Mountains, do not exceed 3 lbs. **LINEX 50 DF** per acre). Do not exceed 40 psi spray nozzle pressure as crop injury may result.

LINEX 50 DF may be applied following an application of Stoddard solvent provided treatments are at least one day apart. Stoddard solvent may be applied following application of **LINEX 50 DF** provided treatments are at least 2 weeks apart. Shorter time intervals between applications may result in injury to the crop. Do not apply **LINEX 50 DF** as a tank mixture with Stoddard solvent, surfactants, nitrogen or fertilizer solution, other pesticides, nor when temperature exceeds 85°F as crop injury may result.

NOTE: Because carrot varieties vary in their resistance, determine tolerance to **LINEX 50 DF** prior to adoption as a field practice to prevent possible crop injury. Do not treat susceptible varieties which show an initial burning of foliage following postemergence treatment with **LINEX 50 DF**. Do not re-enter treated areas treated postemergence for 24 hours following application unless protective clothing is worn.

CELERY — East of Rocky Mountains — **Post-transplant Application:** Make a single application of 1 1/2 to 3 lbs. per acre. Apply as a non-directed spray after celery is transplanted and established, but before celery is 8" tall. Apply before annual grasses exceed 2" in height and before broadleaf weeds exceed 8" in height. In the Northeast, use only on celery grown on muck soils.

Do not exceed 40 psi spray nozzle pressure, and do not apply when temperature exceeds 85°F nor as a tank mixture with surfactants, nitrogen or fertilizer solution, or other pesticides as injury to the crop may result. Do not replant to crops other than celery or carrots within 4 months after application as injury to subsequent crops may result. Do not re-enter treated areas for 24 hours following application unless protective clothing is worn.

CORN (Field) — East of the Rocky Mountains — Preemergence Application — Select one of the following herbicide treatments for application as a tank mixture. A single application should be made after planting but prior to crop emergence. Seed should be planted at least 1 3/4" deep on either flat or raised seedbeds only or injury to the crop may result. Do not spray over top of emerged corn.

Tank Mixture LINEX 50 DF + Lasso

Product Per Acre	Sandy Loam	Soil Texture	
		1 to 3 % Organic Matter	Clay Loam
		Silt Loam	
LINEX 50 DF (lbs.)	2/3 to 1 1/4	1 to 1 2/3	1 1/4 to 2
PLUS			
Lasso (quarts)	3/4 to 1	1 to 1 1/2	1 1/2 to 2
3 to 6% Organic Matter			
LINEX 50 DF (lbs.)	1 1/4 to 2	1 2/3 to 2 1/2	2 to 3
PLUS			
Lasso (quarts)	1 to 1 1/2	1 1/2 to 2	2 to 2 1/2

Replanting: Corn or soybeans may be replanted within 4 months. After 4 months, any crop may be planted.

Tank Mixture LINEX 50 DF + Ramrod

Product Per Acre	Sandy Loam	Soil Texture	
		1 to 3 % Organic Matter	Clay Loam
		Silt Loam	
LINEX 50 DF (lbs.)	2/3 to 1 1/3	1 to 1 2/3	1 1/4 to 2
PLUS			
Ramrod (lbs.)	1 to 2	1 1/2 to 2 1/2	2 to 3
3 to 6 % Organic Matter			
LINEX 50 DF (lbs.)	1 1/3 to 2	1 2/3 to 2 1/2	2 to 3
PLUS			
Ramrod (lbs.)	2 to 3	2 1/2 to 4	3 to 4 3/4

Replanting: Corn may be replanted within 4 months. After 4 months, any crop may be planted.

Tank Mixture LINEX 50 DF + Atrazine

Product Per Acre	Sandy Loam	Soil Texture	
		1 to 2 % Organic Matter	Clay Loam
		Silt Loam	
LINEX 50 DF (lbs.)	2/3 to 1	1 to 1 1/2	1 1/3 to 1 2/3
PLUS			
Atrazine 80% (lbs.)	1/2 to 2/3	2/3 to 1	3/4 to 1
2 to 5 % Organic Matter			
LINEX 50 DF (lbs.)	1 to 2	1 1/2 to 2 1/2	1 2/3 to 3
PLUS			
Atrazine 80% (lbs.)	2/3 to 1 1/4	1 to 1 1/2	1 to 2

CORN (FIELD AND SWEET) Directed Postemergence Application: A single application as a directed spray should be made after corn is at least 15" high. Do not spray over top of corn. There should be sufficient differential between height of corn and weeds so that the directed spray thoroughly covers all weed foliage without contact of upper leaves or whorl of corn by spray or drift. Such contact may cause crop injury. Early cultivation will aid in achieving proper differential between height of corn and weeds.

Apply 1 1/4 to 3 lbs. **LINEX 50 DF** per acre. A pint of surfactant for each 25 gallons of spray mixture may be added. For field corn, non-pressurized nitrogen solution may be substituted for all or part of the water. On lighter soils (low in clay or organic matter) the lower rate should be used and when weeds do not exceed 2" in height. The higher rate should be used on heavier soils (soils high in clay or organic matter) and for weeds up to 5" in height.

COTTON: East of Rocky Mountains — Directed Postemergence Application: **LINEX 50 DF** should be applied as a directed spray with nozzles adjusted to minimize contact to cotton leaves with spray or drift so that crop injury does not result. Do not spray over top of cotton.

When cotton is at least 8" tall and weeds are not over 2" tall, apply 1 to 1 1/2 lbs. **LINEX 50 DF**. One pint of surfactant for each 25 gallons of spray mixture may be added. If a second application is needed use the same rate and apply one week or later after first treatment. Alternatively, a single application of 2 to 3 lbs. per acre following last cultivation can be made after cotton is 20" tall. If emerged weeds are present add surfactant as directed above.

NOTE: Should not be used on Pima varieties of cotton. Do not feed forage or gin trash from treated areas to livestock. Do not graze treated fields.

HYBRID POPLAR — Midwest: Apply 2 to 4 lbs. **LINEX 50 DF** per acre before bud break in the spring. For application after bud break, apply 2 to 4 lbs. **LINEX 50 DF** per acre as a directed spray. Spray should be directed to weed growth and to avoid contact with the poplar plant. Do not spray over the top of the poplar as injury to the plant will result.

Use the lower rate on light soils and higher rate on heavier soils. For best results on emerged weeds, treat at the seedling stage.

More than one treatment may be made but not more than 8 lbs. **LINEX 50 DF** per acre should be applied per year.

PARSNIPS — Preemergence Application: A single application of 1 1/2 to 3 lbs. per acre should be made after planting but prior to crop emergence. Seed should be planted at least 1 1/2" deep.

POTATOES — Preemergence Application: A single application of **LINEX 50 DF** should be made as a broadcast spray after planting but prior to crop emergence. Seed should be planted at least 2" deep. Do not spray over top of emerged potato plants. Weeds are to be "hogged" and/or "hilled", apply after the final "hogging" or "hilling" operation. Application should be made before grasses are 2" tall and before broadleaf weeds are 6" tall, preferably just before or when weed seedlings emerge. A pint of surfactant for each 25 gallons of spray mixture can be added if emerged weeds are present. In irrigated areas, best results are obtained when application is made in a moist soil, followed within 2 weeks by 1" to 2" of sprinkler irrigation (or rainfall). If soil is dry and sandy, irrigate before application and follow with sprinkler irrigation to activate herbicide.

East of Rocky Mountains: On sandy loams, silt loams, soils with 1 to 2% organic matter (coarser soils) apply 1 1/2 to 2 1/2 lbs. of **LINEX 50 DF** per acre. On silty, clay loams and soils with 2 to 5% organic matter (fine soils) apply 2 1/2 to 4 lbs. per acre to emerged weeds (before potato emergence).

Wisconsin — Central Sands Area: **LINEX 50 DF** should be applied at the rate of 1 lb. per acre on sands and 2 lbs. per acre on loamy sands.

Northeast: Use the following tank mix for improved control of annual grasses. Follow directions and precautions on the Dual label also.

BEST AVAILABLE COPY

Product Per Acre	Soil Texture	
	Coarse Sandy Loam	Medium Silt, Loam, Silt Loam
LINEX 50 DF (lbs.)	1 to 1 1/2	1 1/2 to 2
PLUS		
Dual (pints)	1 1/2	2

Product Per Acre	Soil Texture	
	Coarse Sandy Loam	Medium Silt, Loam, Silt Loam
LINEX 50 DF (lbs.)	1 1/2 to 2	2 to 2 1/2
PLUS		
Dual (pints)	2	2 1/2 to 3

SORGHUM — Preemergence Application: Use one of the following herbicide treatments for application as a tank mixture. A single application should be made after planting but prior to crop emergence. Seed should be planted at least 1" deep on flat or raised seedbeds only or injury to the crop may result. Do not spray over top of emerged sorghum.

Tank Mixture LINEX 50 DF with Propazine — Southwest

Product Per Acre	Soil Texture	
	Coarse Sandy Loam	Medium Silt, Loam, Silt Loam
LINEX 50 DF (lbs.)	5/8 to 1	1 to 1 1/2
PLUS		
Propazine 80% (lbs.)	3/8 to 5/8	5/8 to 1

Product Per Acre	Soil Texture	
	Coarse Sandy Loam	Medium Silt, Loam, Silt Loam
LINEX 50 DF (lbs.)	1 to 1 1/2	1 to 2
PLUS		
Propazine 80% (lbs.)	5/8 to 1	5/8 to 1 1/4

Replanting: Do not follow treated sorghum with any fall crop, nor with sugar beets, tobacco, vegetables or potatoes in rotation. Before replanting, thorough seedbed preparation including fall or spring plowing is recommended.

Tank Mixture LINEX 50 DF and Ramrod — Great Plains

Product Per Acre	Soil Texture		
	Coarse Sandy Loam	Medium Silt, Loam, Silt Loam	Fine Silt, Clay, Silty Clay Loam, Clay
LINEX 50 DF (lbs.)	2/3 to 1 1/3	1 to 1 2/3	1 1/4 to 2
PLUS			
Ramrod (lbs.)	1 to 2	1 1/2 to 2 1/2	2 to 3

Product Per Acre	Soil Texture		
	Coarse Sandy Loam	Medium Silt, Loam, Silt Loam	Fine Silt, Clay, Silty Clay Loam, Clay
LINEX 50 DF (lbs.)	1 1/3 to 2	1 2/3 to 2 1/2	2 to 3
PLUS			
Ramrod (lbs.)	2 to 3	2 1/2 to 4	3 to 4 3/4

NOTE: Do not feed sorghum forage or silage from treated fields to dairy animals. Do not graze treated area.

Directed Postemergence Application: A single application of LINEX 50 DF should be made as a directed spray. A pint of surfactant for each 25 gallons of spray mixture may be added. If sprayer is equipped with skids, shoes or shields, apply 1 lb. per acre when sorghum is 12" tall (free standing plants) and weeds are up to 2" high. Use 1 to 2 lbs. per acre when sorghum is 15" tall and weeds are 2 to 4" in height. Application should be made only when there is sufficient differential between height of sorghum and weeds so that the directed spray thoroughly covers all weed foliage without contact of upper leaves or whorl of sorghum by spray or drift as such contact may cause crop injury.

Do not graze livestock in treated areas.

SOYBEANS — Preemergence Application: Select one of the following herbicide treatments and make a single application but prior to crop emergence. Seeds should be planted at least 3/4" deep on flat or raised seedbeds only or crop injury may result. If application is made to fields with standing water or fields too wet to cultivate, injury to soybeans may result. Do not spray over top of emerged soybeans. Do not use on sand or loamy sand or any soil containing less organic matter than listed below.

Preemergence Following Preplant Application of Trifluralin, Prowl® or Tolban®: Where Trifluralin, Prowl or Tolban has been used as a preplant incorporation treatment, apply LINEX 50 DF as a separate operation using rates recommended below for LINEX 50 DF alone. For rotation crops, follow instructions on Trifluralin, Prowl or Tolban labels.

LINEX 50 DF Alone — Add 1 pint surfactant for each 25 gallons of spray mixture if weeds have emerged.

Product Per Acre	Soil Texture		
	Coarse Sandy Loam	Medium Silt, Loam, Silt Loam	Fine Silt, Clay, Silty Clay Loam, Clay
LINEX 50 DF (lbs.)	1 to 1 2/3	1 1/4 to 2 1/3	1 1/3 to 2 2/3
PLUS			
Dual 8E (pts.)	1 1/2	2	2 to 2 1/2

Over 5 % Organic Matter, Use 8 lbs.

Replant: Soybeans or field corn may be replanted within 4 months. After 4 months any crop may be planted.

This tank mix controls the following:

Barnyardgrass	Goosegrass	Prickly Sida
Broadleaf Signalgrass	Hemp Sesbania	Ragweed
Fall Panicum	Lambsquarters	Sicklepod
Foxtail	Pigweed	Signalgrass

This tank mix partially controls cocklebur and jimsonweed.

Tank Mixture LINEX 50 DF with Lexone® DF Herbicide and Lasso

Product Per Acre	Soil Texture		
	Coarse Sandy Loam	Medium Silt, Loam, Silt Loam	Fine Silt, Clay, Silty Clay Loam, Clay
LINEX 50 DF (lbs.)	1/3 to 1/2	1/2 to 3/4	3/4 to 1 1/2
PLUS			
Lexone DF (lbs.)	1/8 to 1/4	1/4 to 1/3	1/3 to 1/2
PLUS			
Lasso (qts.)	3/4 to 1	1 to 1 1/2	1 1/4 to 2

Product Per Acre	Soil Texture		
	Coarse Sandy Loam	Medium Silt, Loam, Silt Loam	Fine Silt, Clay, Silty Clay Loam, Clay
LINEX 50 DF (lbs.)	1/3 to 1/2	1/2 to 3/4	3/4 to 1 1/2
PLUS			
Lexone DF (lbs.)	1/8 to 1/4	1/4 to 1/3	1/3 to 1/2
PLUS			
Lasso (qts.)	3/4 to 1	1 to 1 1/2	1 1/4 to 2

Product Per Acre	Soil Texture		
	Coarse Sandy Loam	Medium Silt, Loam, Silt Loam	Fine Silt, Clay, Silty Clay Loam, Clay
LINEX 50 DF (lbs.)	1/3 to 1/2	1/2 to 3/4	3/4 to 1 1/2
PLUS			
Lexone DF (lbs.)	1/8 to 1/4	1/4 to 1/3	1/3 to 1/2
PLUS			
Lasso (qts.)	3/4 to 1	1 to 1 1/2	1 1/4 to 2

NOTE: Soybean varieties such as Altona, Coker 102 and 156, Goven, NKs, 1884, Semmes, Tracy, Vansoy, Terra Vig 505 and 606, and AP 71 are sensitive to Lexone DF. If Lexone DF is used on these varieties, injury may occur. Tolerance to Lexone DF should be determined before use on any other soybean variety. If Lexone DF is used on soils having a calcareous surface layer or pH of 7.5 or higher, or if used in conjunction with soil applied organo phosphate pesticides such as Desanit, Dlayston, Mocap, Nemacur, Thimet, Parathion, Lorsban 15G or Counter, injury to soybeans may occur.

Do not feed treated plant parts to livestock. If atrazine was applied on the soil the year before, injury may occur. Increased possibility of crop injury may occur when soybean seedlings are weakened because of seedling disease, cold weather, deep planting (more than 2"), excessive moisture, high soil pH of 7.5 or higher, higher salt concentration or drought.

Cultivation After Planting: Treated soil may be shallow-cultivated, rotary hoed or hand hoed without reducing the weed control activity of the tank mixture. Do not cultivate deeper than the treated layer

of soil since this may bring untreated soil to the surface and poor weed control may result.

Replanting: Treated fields may be replanted to soybeans if initial seeding fails to produce a stand. Soil should not be reworked. Crop injury may result if second application is applied. Do not replant treated areas to any crop other than soybeans within four months after treatment as injury to subsequent crops may result.

Tank Mixture LINEX 50 DF with Lasso

Product Per Acre	Soil Texture		
	Coarse Sandy Loam	Medium Silt, Loam, Silt Loam	Fine Silt, Clay, Silty Clay Loam, Clay
LINEX 50 DF (lbs.)	2/3 to 1 1/4	1 to 1 2/3	1 1/4 to 2
PLUS			
Lasso (qts.)	3/4 to 1	1 to 1 1/2	1 1/2 to 2

Product Per Acre	Soil Texture		
	Coarse Sandy Loam	Medium Silt, Loam, Silt Loam	Fine Silt, Clay, Silty Clay Loam, Clay
LINEX 50 DF (lbs.)	1 1/4 to 2	1 2/3 to 2 1/2	2 to 3
PLUS			
Lasso (qts.)	1 to 1 1/2	1 1/2 to 2	2 to 2 1/2

For Control of Black Nightshade Apply:

Product Per Acre	Soil Texture		
	Coarse Sandy Loam	Medium Silt, Loam, Silt Loam	Fine Silt, Clay, Silty Clay Loam, Clay
LINEX 50 DF (lbs.)	3/4 to 1 1/2	1 to 1 2/3	1 1/4 to 2
PLUS			
Lasso (qts.)	2	2 1/2 to 3	3

Replanting: Soybeans or field corn may be replanted within 4 months. After 4 months, any crop may be planted.

Tank Mixture LINEX 50 DF with Lexone DF and Dual 8E: Controls common ragweed, prickly sida, sicklepod, hemp sesbania, lambsquarters, pigweed, and partially controls cocklebur, and jimsonweed.

Product Per Acre	Soil Texture		
	Coarse Sandy Loam	Medium Silt, Loam, Silt Loam	Fine Silt, Clay, Silty Clay Loam, Clay
LINEX 50 DF (lbs.)	1/3 to 1/2	1/2 to 3/4	3/4 to 1 1/2
PLUS			
Lexone DF (lbs.)	1/6 to 1/4	1/4 to 1/3	1/3 to 1/2
PLUS			
Dual 8E (pts.)	1 to 1 1/4	1 1/4 to 1 1/2	1 1/2 to 2

Product Per Acre	Soil Texture		
	Coarse Sandy Loam	Medium Silt, Loam, Silt Loam	Fine Silt, Clay, Silty Clay Loam, Clay
LINEX 50 DF (lbs.)	1/3 to 3/4	3/4 to 1 1/2	1 1/2 to 2
PLUS			
Lexone DF (lbs.)	1/4 to 1/3	1/3 to 1/2	1/2 to 2/3
PLUS			
Dual 8E (pts.)	1 1/2 to 2	1/2 to 2/3	2 to 2 1/2

NOTE: Soybean varieties such as Altona, Coker 102 and 156, Goven, NKs 1884, Semmes, Tracy, Vansoy, Terra Vig 505 and 606, and AP 71 are sensitive to Lexone DF. If Lexone DF is used on these varieties, injury may occur. Tolerance to Lexone DF should be determined before use on any other soybean variety. If Lexone DF is used on soils having a calcareous surface layer or pH of 7.5 or higher, or if used in conjunction with soil applied organo phosphate pesticides such as Desanit, Dlayston, Mocap, Nemacur, Thimet, Parathion, Lorsban 15G or Counter, injury to soybeans may occur.

Do not feed treated plant parts to livestock. If atrazine was applied on the soil the year before, injury may occur. Increased possibility of crop injury may occur when soybean seedlings are weakened because of seedling disease, cold weather, deep planting (more than 2"), excessive moisture, high soil pH of 7.5 or higher, high salt concentration or drought.

Cultivation After Planting: Treated soil may be shallow-cultivated, rotary hoed or hand hoed without reducing the weed control activity of the tank mixture. Do not cultivate deeper than the treated layer of soil since this may bring untreated soil to the surface and poor weed control may result.

Replanting: Treated fields may be replanted to soybeans if initial seeding fails to produce a stand. Soil should not be reworked. Crop injury may result if second application is applied. Do not replant treated areas to any crop other than soybeans within four months after treatment as injury to subsequent crops may result.

Tank Mixture LINEX 50 DF with Dual 8E

Product Per Acre	Soil Texture		
	Coarse Sandy Loam	Medium Silt, Loam, Silt Loam	Fine Silt, Clay, Silty Clay Loam, Clay
LINEX 50 DF (lbs.)	2/3 to 1 1/4	1 to 1 2/3	1 1/4 to 2
PLUS			
Dual 8E (pts.)	1 1/4	1 1/2	2

Product Per Acre	Soil Texture		
	Coarse Sandy Loam	Medium Silt, Loam, Silt Loam	Fine Silt, Clay, Silty Clay Loam, Clay
LINEX 50 DF (lbs.)	1 1/4 to 2	1 2/3 to 2 1/2	2 to 3
PLUS			
Dual 8E (pts.)	1 1/2	2	2 to 2 1/2

Do not feed treated plant parts to livestock. Do not graze treated areas.

Replanting: Soybeans or field corn may be replanted within 4 months. For rotation crops follow instructions on the Dual 8E label.

For control of black nightshade in Kentucky, Illinois, Indiana and Ohio:

Product Per Acre	Soil Texture		
	Coarse Sandy Loam	Medium Silt, Loam, Silt Loam	Fine Silt, Clay, Silty Clay Loam, Clay
LINEX 50 DF (lbs.)	3/4 to 1 1/2	1 to 1 2/3	1 1/4 to 2
PLUS			
Dual 8E (pts.)	1 1/2	2	2 to 2 1/2

Do not use on sand or loamy sands.

Replanting: Soybeans or field corn may be replanted within four months. Any crop may be planted after 4 months.

Tank Mixture LINEX 50 DF and Surflan® 75W or Surflan® AS

Product Per Acre	Soil Texture		
	Coarse Sandy Loam	Medium Silt, Loam, Silt Loam	Fine Silt, Clay, Silty Clay Loam, Clay
LINEX 50 DF (lbs.)	2/3 to 1 1/4	1 to 1 2/3	1 1/4 to 2
PLUS			
Surflan 75W (lbs.)	2/3 to 1	1 to 1 1/3	1 1/3 to 1 2/3
OR			
Surflan AS (qts.)	1/2 to 3/4	3/4 to 1	1 to 1 1/4

Do not feed treated plant parts to livestock.

Replanting: Soybeans may be replanted within 4 months. Any crop except root crops may be planted after 4 months, do not plant root crops within 12 months.

Tank Mixture LINEX 50 DF with Prowl

Product Per Acre	Soil Texture		
	Coarse Sandy Loam	Medium Silt, Loam, Silt Loam	Fine Silt, Clay, Silty Clay Loam, Clay
LINEX 50 DF (lbs.)	2/3 to 1 1/4	1 to 1 2/3	1 1/4 to 2
PLUS			
Prowl (pts.)	1 to 1 1/2	1 1/2 to 2	1 1/2 to 2

Product Per Acre	Soil Texture		
	Coarse Sandy Loam	Medium Silt, Loam, Silt Loam	Fine Silt, Clay, Silty Clay Loam, Clay
LINEX 50 DF (lbs.)	1 1/4 to 2	1 2/3 to 2 1/2	2 to 3
PLUS			
Prowl (pts.)	1 1/2	1 1/2 to 2	2 to 2 1/2

Replanting: Soybeans or field corn may be replanted within 4 months. Other crops may be planted

BEST AVAILABLE COPY

the following year.

Tank Mixture LINEX 50 DF with Amiben®

Product Per Acre	Soil Texture		
	Sandy Loam	1/2 to 3 % Organic Matter Silt Loam	Clay Loam
LINEX 50 DF (lbs.)	2/3 to 1 1/4	1 to 1 2/3	1 1/4 to 2
PLUS			
Amiben (qts.)	3	3	4
3 to 6 % Organic Matter			
LINEX 50 DF (lbs.)	1 1/4 to 2	1 2/3 to 2 1/2	2 to 3
PLUS			
Amiben (qts.)	3	4	5

Replanting: Soybeans or field corn may be replanted within 4 months. Any other crops may be planted after 4 months.

Tank Mixture LINEX 50 DF with Ramrod — Seed Crop Only

Product Per Acre	Soil Texture		
	Sandy Loam	1 to 3 % Organic Matter Silt Loam	Clay Loam
LINEX 50 DF (lbs.)	2/3 to 1 1/4	1 to 1 2/3	1 1/4 to 2
PLUS			
Ramrod (lbs.)	1 to 2	1 1/2 to 2 1/2	2 to 3
3 to 6 % Organic Matter			
LINEX 50 DF (lbs.)	1 1/4 to 2	1 2/3 to 2 1/2	2 to 3
PLUS			
Ramrod (lbs.)	2 to 3	2 1/2 to 4	3 to 4 3/4

NOTE: Do not feed treated plant parts to livestock. Do not graze treated area. Do not use seed for food, feed or oil purposes.

Replanting: Soybeans, sorghum or field corn may be replanted within 4 months. Any crop may be planted after 4 months.

SOYBEANS — Minimum or No-Tillage

LINEX 50 DF preemergence to soybeans may be used with paraquat, Bronco or Roundup in tank mix combinations for preemergence and postemergence control of many broadleaf weeds and grasses and for top kill or suppression of some perennials where soybeans will be planted directly into a preformed bed (stale seed bed), cover crop or in previous crop residues such as corn or small grain stubble. Use as described for Preemergence Application for conventional seed bed preparation in reduced tillage systems. Adding Lasso, Dual SE, Surflan 75W or Surflan AS to the tank mixture improves control of grasses and volunteer small grains. Apply with ground equipment immediately before, during or after planting but before crop emerges. Maintain continuous agitation or spray mixture.

Tank Mixture LINEX 50 DF with Lasso and Paraquat or Roundup®: LINEX 50 DF and Lasso should be thoroughly mixed first according to directions. Add paraquat or Roundup as directed under Paraquat combinations or Roundup combinations below.

Product Per Acre	Soil Texture		
	Sandy Loam	1/2 to 3 % Organic Matter Silt Loam	Clay Loam
LINEX 50 DF (lbs.)	3/4 to 1 1/2	1 1/8 to 2	1 1/4 to 2 1/4
PLUS			
Lasso (qts.)	2	2 1/2	2 1/2
3 to 6 % Organic Matter			
LINEX 50 DF (lbs.)	1 1/2 to 2 2/5	2 to 3	2 1/4 to 3 1/2
PLUS			
Lasso (qts.)	2 1/2	3	3

For control of black nightshade apply:

Product Per Acre	Soil Texture		
	Sandy Loam	1 to 3 % Organic Matter Silt Loam	Clay Loam
LINEX 50 DF (lbs.)	3/4 to 1 1/2	1 to 2	1 1/4 to 2 1/4
PLUS			
Lasso (qts.)	2	2 1/2 to 3	3

Replanting: Soybeans or field corn may be replanted within 4 months. Any crop may be planted after 4 months.

Tank Mixture LINEX 50 DF with Bronco® — Do not add Paraquat or Roundup to this combination.

Product Per Acre	Soil Texture		
	Sandy Loam	1/2 to 5 % Organic Matter Silt Loam	Clay Loam
LINEX 50 DF (lbs.)	3/4 to 2	1 1/8 to 2 1/2	1 1/4 to 3
PLUS			
Bronco (lbs.)	3 1/4 to 5	4 to 5	4 to 5

Do not use on soils with less than 1/2 organic matter or on sand, loamy sand or muck soils as crop injury may result.

Use the high rate of Bronco for dense stubble, heavy crop residue or heavy weed population.

Replanting Soybeans: Treated fields may be replanted to soybeans if initial seeding fails to produce a stand. Soil should not be reworked. Crop injury may result if second application is applied. Do not replant treated areas to any crop other than soybeans within four months after treatment as injury to subsequent crops may result.

Tank Mixture LINEX 50 DF with Surflan 75W or Surflan AS and Paraquat or Roundup: LINEX 50 DF should be thoroughly mixed with Surflan first in the tank, following direction. Then add Paraquat or Roundup as directed under Paraquat Combinations or Roundup Combinations below.

Product Per Acre	Soil Texture		
	Sandy Loam	1/2 to 3 % Organic Matter Silt Loam	Clay Loam
LINEX 50 DF (lbs.)	3/4 to 1 1/2	1 1/8 to 2	1 1/4 to 2 1/4
PLUS			
Surflan 75W (lbs.)	2/3 to 1	1 to 1 1/3	1 1/3 to 1 2/3
OR			
Surflan AS (qts.)	1/2 to 3/4	3/4 to 1	1 to 1 1/4

Do not feed treated plant parts to livestock.

Replanting: Soybeans may be replanted within 4 months. After 4 months any crop may be planted except do not plant root crops within 12 months.

Tank Mix LINEX 50 DF with Paraquat or Roundup: LINEX 50 DF should be thoroughly mixed with

paraquat in spray tank following directions under Paraquat Combinations or Roundup Combinations below.

Product Per Acre	Soil Texture		
	Sandy Loam	1/2 to 3 % Organic Matter Silt Loam	Clay Loam
LINEX 50 DF (lbs.)	1 to 1 2/3	1 1/4 to 2 1/3	1 1/3 to 2 2/3
2 to 5 % Organic Matter			
LINEX 50 DF (lbs.)	1 2/3 to 3	2 1/3 to 4	2 2/3 to 5

(Over 5 % Organic Matter, Use 6 lbs.)

Replanting: Soybeans or field corn may be replanted within 4 months. Any crop may be planted after 4 months.

Paraquat Combinations — Choose and tank mix one of the above treatments in water. Then add 1 to 2 pints Paraquat (2 lbs. active per gal.) per acre for control of emerged weeds. The higher rate should be used for weeds 4" to 6" tall. Add 1/2 pint of surfactant or spreader per 100 gallons of spray mixture as the last ingredient. Maintain continuous agitation. Use 20 to 60 gallons of water per acre. Use the higher gallonage for dense stubble or vegetation.

Roundup Combinations — Choose and tank mix one of the above treatments. Add 1 1/2 quarts of Roundup per acre for control of emerged annual weeds or 2 to 4 quarts per acre for control of emerged perennial annual weeds. Use 20 to 30 gallons of water per acre.

Bronco Combinations — Mix LINEX 50 DF with water in spray tank as directed on the package label and then add Bronco to the dilute mix. Maintain continuous agitation. Use 20 to 30 gallons of water per acre.

SOYBEANS — Directed Postemergence Application: LINEX 50 DF and 2,4-DB can be tank mixed with Premerge 3 as a directed spray to cover weed foliage with minimum contact of the soybean plant. Do not spray higher than 3" on the soybean stem or crop injury may result. Do not spray over top of soybean plants. For broadcast application, use a single flood type spray nozzle. ("K" series or equivalent) per middle mounted on an oiling shoe or gauge wheel. For band treatment use two nozzles per row mounted on an oiling shoe or gauge wheel, one on each side of row. Spray drift may cause crop injury. Do not exceed nozzle pressure of 25 psi nor use nozzle tips smaller than 8002 T-jet (or equivalent) and do not spray under windy conditions. A pint of surfactant may be added for each 25 gallons of spray mixture. Use a preemergence treatment or cultivation to control early weed growth and to increase the differential between height of soybeans and weeds for best results.

LINEX 50 DF with 2,4-DB — Apply 1 lb. LINEX 50 DF plus 1/5 lb. 2,4-DB per acre when soybeans are at least 8" high and when weeds do not exceed 4" in height. Make a second application if needed, but not more than 2 applications per season.

Do not use on soils with less than 1/2% organic matter. Do not apply more than 2 lbs. LINEX 50 DF per acre per season for postemergence treatments. Do not apply within 60 days of harvest. Do not feed soybean forage or hay to livestock from fields treated postemergence. Harvested soybeans may be used for food, feed or oil purposes.

LINEX 50 DF Alone: Apply 1/2 to 1 lb. LINEX 50 DF per acre for soybeans at least 8" high and when weeds do not exceed 2" in height. A second application should be made at the same rate if new flush of weeds occurs.

A single application of 1 to 2 lbs. of LINEX 50 DF per acre should be made for soybeans at least 12" high and when weeds do not exceed 4" high. A split application can be made of 1 lb. per acre followed by a second application at the same rate after 1 week or later.

NONCROP WEED CONTROL

Apply 2 to 6 lbs. LINEX 50 DF per acre in 40 to 100 gallon of water for short term control of annual weeds on noncropland areas such as roadsides and fence rows. Apply shortly before weed growth begins or at early seedling stage of growth for best results. Add 2 qts. of surfactant per 100 gallons of spray mixture for control of established annual weeds. Apply as a thorough coverage spray during periods when daily temperatures exceed 70°F and before weed growth exceeds 8" in height.

WARRANTY

GRIFFIN CORPORATION warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of GRIFFIN. In no case shall GRIFFIN be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. GRIFFIN MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

Amiben is a registered trademark of Union Carbide Agricultural Products Company, Inc.
Bronco is a registered trademark of Monsanto Company
Dual is a registered trademark of Ciba-Geigy Corporation
Lasso is a registered trademark of Monsanto Company
Laxone is a registered trademark of DuPont de Nemours Company
LINEX is a registered trademark of Griffin Corporation
Prowl is a registered trademark of American Cyanamid Company
Ramrod is a registered trademark of Monsanto Company
Roundup is a registered trademark of Monsanto Company
Surflan 75W is a registered trademark of Elanco Products Company
Surflan AS is a registered trademark of Elanco Products Company
Tolban is a registered trademark of Ciba Geigy Corporation

GCN 113080
CPC 027576

BEST AVAILABLE COPY