

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

PM23

1812-328

1/11

APR 28 1995

Ms. Carol Eakins
Griffin Corporation
P.O. Box 1847
Valdosta, GA 31603

Dear Ms. Eakins:

SUBJECT: Label Amendment (Pot in Pot Non-Food Use)
Trilin 10G
EPA Registration No. 1812-328
Your Application Dated January 13, 1995

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable provided you make the following changes:

- Under the use directions for "Pot-in-Pot Nursery Production of Trees and Shrubs":
 - Revise the label to indicate that the "Pot-in-Pot" use directions are for "ornamental trees and shrubs" only. Alternatively, you may add the sentence "Do not apply to pots used for growing plants which will bear edible fruit within one year following application."
 - Revise the first sentence of the second paragraph to read: "Apply Trilin 10G uniformly over the bottom of the socket pot before the growing pot is inserted at the rate corresponding to the container size."
- Include the EPA establishment number, preceded by the phrase "EPA Est. No." on the product label.
- Add the statement "See Additional Precautionary Statements," and a phrase describing the location of the additional precautionary statements to the first page of the label.
- Correct the spelling of the word "explique" in the Spanish language warning on the first page of the label.

RD:STANTON:PM Team 23:Rm. 235:305-5218:Disk #1:S480675.LET

CONCURRENCES

SYMBOL ▶	7505C							
SURNAME ▶	S. Stanton							
DATE ▶	Apr 26, 1995							

2/11
A stamped copy of the label is enclosed for your records.
Please submit (one) final printed copy for the above mentioned
label before releasing the product for shipment.

Sincerely yours,

Joanne I. Miller
Product Manager (23)
Fungicide-Herbicide Branch
Registration Division (7505C)

Enclosure

Griffin



3/11

Trilin[®] 10G

HERBICIDE

Made in the USA

ACCEPTED
with COMMENTS
In EPA Letter Dated
APR 28 1995

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
1812-328

ACTIVE INGREDIENT

Trifluralin a,a,a-trifluoro-2,6-dinitro-N,N-dipropyl-p-toluidine..... 10%

INERT INGREDIENTS..... 90%

TOTAL..... 100%

KEEP OUT OF REACH OF CHILDREN WARNING - AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail).

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush with plenty of water for at least 15 minutes. Get medical attention.

IF ON SKIN: Wash immediately with soap and water.

IF SWALLOWED: Drink 1 to 2 glasses of water. Induce vomiting by placing finger in back of throat. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth to mouth. Get medical attention.

GRIFFIN CORPORATION
VALDOSTA, GEORGIA 31601

Specimen Label

EPA REG. NO. 1812-328

BEST AVAILABLE COPY

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS (AND DOMESTIC ANIMALS)

WARNING

Precautionary Statements: Causes substantial but temporary eye injury. Harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with skin, eyes or clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift or runoff from treatment areas may be hazardous to aquatic organisms in neighboring aquatic sites. Do not contaminate water when disposing of equipment washwaters.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Do not apply this product through any type of irrigation system. Trilin 10G is a herbicide which must be applied prior to the emergence (preemergence) of annual broadleaf and grassy weeds. To assure prolonged control (seasonal) of weeds shown below, soil incorporation is essential.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

Exception: If product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

STORAGE AND DISPOSAL

Do not contaminate water, food, feed, other pesticides, fertilizer or seeds.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag by shaking and tapping sides and bottom to loosen any clinging particles into application equipment. Then dispose of bags in a sanitary landfill or by incineration, or if allowed by State and local authorities by burning. If burned, stay out of smoke.

BROADLEAF AND GRASSY WEEDS CONTROLLED BY TRILIN 10G

BROADLEAF WEEDS

Carpetweed	<u>Mollugo verticillata</u>	Pigweed	<u>Amaranthus spp.</u>
Chickweed	<u>Stellaria media</u>	(Carelessweed)	
Florida Pursley	<u>Richardia scabra</u>	(Prostrate Pigweed)	
(Florida purslane)		(Redroot)	
(Mexican clover)		(Rough Pigweed)	
(Pursley)		(Spiny Pigweed)	
Goosefoot	<u>Chenopodium hybridum</u>	Puncturevines (Western U.S. only)	<u>Trifolium terrestris</u>
Henbit	<u>Lamium amplexicaule</u>	(Caltrop)	
		(Goathead)	
Knotweed	<u>Polygonum aviculare</u>	Purslane	<u>Portulaca oleracea</u>
Lambsquarters	<u>Chenopodium album</u>	Stinging Nettle	<u>Urtica dioica</u>
		(Nettle)	

GRASSES

Annual bluegrass	<u>Poa annua</u>	Johnsongrass	<u>Sorghum halepense</u>
Annual Ryegrass	<u>Lolium multiflorum</u>	(from seed)	
Barnyardgrass	<u>Echinochloa sp.</u>	Junglecane	<u>Echinochloa polystachya</u>
(watergrass)		Sandbur	<u>Cenchrus incertus</u>
Bromegrass	<u>Bromus tectorum</u>	(Burgess)	
(Cheatgrass)		Signalgrass (Brachiaria)	<u>Brachiaria sp.</u>
(Downy Brome)		Sprangletop	<u>Leptochloa latifolia</u>
Cheat	<u>Bromus secalinus</u>	Stinkgrass (Lovegrass)	<u>Eragrostis ciliaris</u>
(Chess)		Wild Cane (Shallercane)	<u>Sorghum bicolor</u>
Crabgrass	<u>Digitaria spp.</u>	Wild Oat	<u>Avena fatua</u>
(Large Crabgrass)		Whoolly cupgrass	<u>Eriochloa villosa</u>
(Smooth Crabgrass)			
Foxtail	<u>Setaria spp.</u>		
(Bottlegrass) (Bristlegrass)			
(Giant Foxtail) (Green Foxtail)			
(Foxtail Millet) (Pigeongrass)			
(Robust Foxtail) (Yellow Foxtail)			

* Applied as a preplant incorporated (PPI) treatment Trilin 10G will provide partial control of wild oats. This claim is for all PPI uses except for fall application for spring seeded cereals at foxtail (pigeongrass) control rates.

Long term and continued use of trifluralin has resulted in the selection of tolerant populations in certain species of weeds. This situation is limited to a few weeds and is generally geographically specific. Weed species known to have some trifluralin tolerant populations are goosegrass, green foxtail (pigeongrass) and Palmer amaranthus (Palmer pigweed). Trilin 10G is not recommended for the control of goosegrass, tolerant green foxtail or Palmer amaranthus. Consult State Agricultural Extension Service or Experiment Station weed specialist for specific recommendations for local weed problems.

SOIL PREPARATION

Soil surface should be smooth enough to enable the proper operation of application and incorporation equipment to insure uniform distribution and incorporation of Trilin 10G. Interference can be caused by ground cover such as crop residue or existing weeds. Soil should be tilled prior to the application of Trilin 10G to allow uniform incorporation into the top 2 to 3 inches of soil. Soil moisture should be such that large clods will be broken up during incorporation process.

Crop Residues or Existing Weeds

Ground cover, such as crop residues or existing weeds, can interfere with the incorporation of Trilin 10G into the soil. A manageable level of such ground cover will allow the Trilin 10G to be uniformly incorporated into the top 2 to 3 inches of soil. If the level of the ground cover is such that this cannot be done, you must till the soil prior to the application of Trilin 10G.

Roughness

The soil surface should be smooth enough so that you can operate application and incorporation equipment properly and at speeds which insure a uniform distribution and incorporation of Trilin 10G.

General Soil Conditions

To assure incorporation of Trilin 10G, soil moisture conditions should be such that large clods can be broken up during the incorporation process.

SOIL TEXTURE

Before application determine soil texture in order to apply the correct rate. Rates given in this booklet refer to the following soil texture groups:

COARSE: Sand, Loamy sand, Sandy loam

MEDIUM: Loam, Silt, Silty Clay loam, Silt loam, Sandy clay loam.

FINE: Clay, Clay loam, Silty clay loam, Silty clay, Sandy clay, Sandy clay loam.

Silty clay loam and sandy clay loam soils are transitional soils and may be classified as either medium or fine textured soils. If silty clay loam or sandy clay loam soils are predominantly sand or silt, they are usually classified as medium textured soils. If they are predominantly clay, they are usually classified as fine textured soils.

A fine textured soil will require a higher application rate than a coarse textured soil. Do not exceed recommended rates.

APPLICATION DIRECTIONS

Aerial/Ground Application

Trilin 10G may be applied with ground or aerial broadcast application equipment. Apply only with equipment capable of accurate calibration and uniform application of herbicide granules. Once properly calibrated the granular distributor should be checked regularly to assure proper operating condition. Do not apply Trilin 10G in narrow bands. Apply at the recommended rate for soil texture class to be treated.

INCORPORATION DIRECTIONS

General Directions

Trilin 10G is a selective preemergence herbicide for control of many annual grasses and broadleaf weeds. Two incorporation passes are required. The first pass should occur within 24 hours after application. For best results, the second incorporation should be delayed 3 to 5 days after the first and be completed prior to planting. A minimum delay of 7 days after the first incorporation is recommended for certain uses in small grains. The second incorporation pass should occur in a different direction than the first. Incorporate Trilin 10G uniformly into the top 2 to 3 inches of the final seedbed. Set equipment so that the second incorporation pass is not deeper than the first as erratic weed control may result if untreated soil is moved to the surface during the second incorporation pass.

After Planting

For directions after planting check label under specific crop.

Bedded Culture

Trilin 10G needs to be incorporated into the top 2 to 3 inches of the final seedbed for effective weed control.

Application Prior to Bedding

Apply Trilin 10G prior to bedding and incorporate it with recommended equipment. The bedding operation serves as the second incorporation. Avoid removal of untreated soil from the seedbed before or during the planting operation. This would expose untreated soil, allowing weeds to germinate in the drill row.

Application After Bedding

Knock off beds to planting height before application and incorporate Trilin 10G with recommended equipment that will conform to the bed shape. Do not leave untreated soil exposed. Avoid removal of untreated soil from the seedbed before or during the planting operation. This would expose untreated soil, allowing weeds to germinate in the drill row.

Cultivation After Planting

Areas treated with Trilin 10G may be shallowly cultivated without loss of weed control activity. Avoid deep cultivation that could bring untreated soil to the soil surface and result in loss of weed control.

Recommended equipment

Any recommended incorporation equipment may be used alone or in combination with any other recommended equipment. Two incorporation passes are necessary unless specifically stated. The second incorporation should not be deeper than the first.

Disc: Set to cut 4 to 6 inches deep and operate at 4 to 6 mph.

Chisel Plow: (For use in northern Great Plains) The chisel plow may be used for the first incorporation pass only. Any other recommended incorporation equipment may be used for the second pass for row crops. The chisel plow may be used for any tillage or incorporation pass in the Summer Fallow program. Operate chisel plow 4 to 5 inches deep at 4 to 6 mph. A chisel plow is defined as having 3 rows or 14 to 16 inch sweeps spaced no more than 12 inches apart. Sweeps should

be staggered so that soil is left unturned. Chisel points should not be used.

Field Cultivator: Defined as implements with sweeps of 3 to 4 rows spaced at intervals of 7 inches or less, staggered so that no soil is left unturned. Set to cut 3 to 4 inches deep, operate at 5 mph or more. Do not use chisel points.

Combination Seedbed Conditioners: Set to cut 3 to 4 inches deep and operate at a speed of at least 5 mph. These are defined as three or more tillage devices combined and used as a single tool. For example 2 to 3 rows of field cultivator C- or S-shaped shanks with an effective sweep spacing of 6 to 9 inches (staggered so that no soil is left unturned), followed by a spike tooth or flexline harrow, followed by a ground driven reel or basket.

Rolling Cultivator: Set to cut 2 to 4 inches deep and operate at a speed of 6 to 8 mph. Rolling cultivators are effective for use on coarse and medium textured soils. The rolling cultivator may be used on fine textured soils when used in sugarcane.

Bed Conditioner (Do-All): Set to cut 2 to 4 inches deep and operate at a speed of 4 to 6 mph. The Do-All is effective when used on coarse or medium textured soils only. Only one incorporation pass is necessary in bedded culture. Two passes with the Do-All are necessary in flat planted culture.

Mulch Treader (other similar disc-type implements): Set Mulch Treader to cut 3 to 4 inches deep and operate at 5 to 8 mph.

P.T.O. Driven Equipment (tillers, cultivators, hoes): Only one incorporation is required. Adjust to incorporate Trilin 10G into the top 2 to 3 inches of the seedbed using rollers spaced to give a clean sweep of the soil. P.T.O. equipment should not be operated at a speed greater than 4 mph.

Other Equipment: Including the flexible tine-tooth harrow (Flexline, Melroe), is also recommended but only for the special usage specified in this label. Use other equipment only as specified herein.

CROP RECOMMENDATIONS

All recommendations are given as the broadcast rates of Trilin 10G per acre. Apply Trilin 10G any time after January 1 when the soil can be worked and is suitable for good incorporation. For fall application see specific crop recommendations for "FALL APPLICATION" heading.

FALL APPLICATION (GENERAL)

These comments apply to areas receiving more than 20 inches average annual rainfall and for crops where Trilin 10G is recommended for preemergence usage, but no specific fall application directions given. If specific fall application rates are not given, use Trilin 10G at rates recommended for spring application. Always see specific crop recommendations before using Trilin 10G. NOTE: Do not apply Trilin 10G prior to planting sugar beets, potatoes and direct-seeded tomatoes in the fall.

In most states apply and incorporate Trilin 10G between October 15 and December 31. In California, Minnesota, North Dakota and South Dakota Trilin 10G may be applied and incorporated between September 1 and December 31.

Apply and incorporate on ground left flat or bedded during the winter. If bedded, knock down to proper height prior to planting moving some treated soil from bed tops into furrows. If treated soil is left flat for winter, exercise caution in bedding operations to avoid turning up treated soil. Any weeds establishing as a result of bedding or other operations should be destroyed before planting. NOTE: Trilin 10G should not be applied in the fall to soils which are wet or subject to prolonged flooding or where rice was grown.

SPECIAL PRECAUTIONS

Under normal growing conditions and if applied according to directions, Trilin 10G will not harm the treated crop. Crop injury or soil residue may result from over application. Erratic weed control or crop injury may result from uneven application or improper soil incorporation of Trilin 10G. Seedling disease, cold weather, deep planting, excessive moisture, high salt concentration or drought may weaken crop seedlings and increase the possibility of damage from Trilin 10G. Under these conditions, delayed crop development or reduce yields may result.

Where specific crop recommendations are not given, use the lower rate for coarser soils or soils with lower organic matter. For soils containing 10% or more organic matter do not use Trilin 10G.

In Arizona, Colorado, Idaho, Montana, Nevada, New Mexico, Oregon, Utah, Washington and Wyoming, after a spring application of Trilin 10G, sugar beets, red beets or spinach should not be planted for 12 months or 14 months after fall application.

Soil should be plowed to a depth of 12 inches prior to planting sugar beets to prevent the possibility of crop injury. After a spring application sorghum (milo), proso millet, corn or oats should not be planted for 14 months or for 16 months after a fall application of Trilin 10G to avoid crop injury. If land has not been irrigated, do not plant any of these crops for 18 months after a spring application or 20 months after a fall application of Trilin 10G.

In all other area sugarbeets, redbeets and spinach should not be planted for 12 months after a spring application or 14 months after a fall application. Before planting sugarbeets, moldboard plow to a depth of 12 inches to reduce the possibility of crop injury.

In Arizona, California, Colorado, Idaho, Montana, Nevada, New Mexico, Oregon, Utah, Washington and Wyoming after a spring application of Trilin 10G, proso millet, corn, sorghum (milo), oats and annual or perennial grass crops or grass mixtures should not be planted for 12 months or 14 months after fall application. If land has not been irrigated, these crops should not be planted for 18 months after a spring application or 20 months after a fall application. Moldboard plowing to depth of 12 inches before planting these crops will reduce the possibility of crop injury.

In Minnesota, North Dakota and South Dakota proso millet, sorghum (milo), oats and annual or perennial grass crops or grass mixtures should not be planted for 12 months after a spring application or 21 months after a fall application of Trilin 10G.

In those areas of Kansas, Nebraska, North Dakota, Oklahoma, South Dakota and Texas where at least 20 inches of rainfall (total) was used to produce the crop, sorghum or oats should not be planted for 12 months after an application of Trilin 10G.

Do not plant sorghum, proso millet, or oats for 18 months after an application of Trilin 10G if less than 20 inches of total water was used to produce the crop. Cool, wet weather conditions during the early stage of growth may increase the possibility of injury to sorghum.

Vegetable crops other than those listed on this label for preplant soil incorporated application of Trilin 10G should not be planted within 5 months following the application of Trilin 10G or other trifluralin products.

ALFALFA Established

Trilin 10G can be applied to established alfalfa during dormancy or throughout the growing season immediately after cutting. Apply Trilin 10G for control of barnyardgrass, bromegrass (cheatgrass, downy brome), canarygrass, cheat (chess), crabgrass, woolly cupgrass, foxtail, junglerice, sandbur and wild barley. Because Trilin 10G does not control established weeds, application must be made prior to the expected time of weed germination.

Apply Trilin 10G to established alfalfa stands prior to weed emergence at a rate of 20 pounds per acre on all soil textures. Trilin 10G may be applied using properly calibrated ground or aerial application equipment. A single rainfall or overhead sprinkler irrigation of 0.5 inches or more or flood irrigation after application is required to activate Trilin 10G. If this does not occur within three days after application, Trilin 10G should be activated using incorporation equipment that will ensure thorough soil mixing with minimum damage to the established alfalfa. If activated using furrow irrigation, the surface of the beds between furrows should be thoroughly wetted.

Do not apply within 21 days of harvest.

Fall Application

Apply Trilin 10G immediately after a cutting between August 1 and October 1. When fall applied Trilin 10G controls bromegrass and cheat in addition to other weeds listed above that germinate after application. Bromegrass and cheat begin to germinate in the fall with the onset of cooler weather.

PRECAUTIONS

Where alfalfa is to be rotated to another crop in the season following application of Trilin 10G, plant only those crops to which Trilin 10G can be applied as a preplant incorporated treatment or injury will result.

ASPARAGUS Established

Apply Trilin 10G to established asparagus as a single or split application for suppression of volunteer seedling asparagus and field bindweed, when recommended rates, application and timing are followed.

Apply in winter or early spring after mature ferns have been removed, but before new spears begin to emerge.

Apply post-harvest applications immediately after harvest in late spring or early summer just before ferns are allowed to develop

Broadcast Rate Per Acre
Trilin 10G (pounds)

	Coarse	Medium	Fine
Single Application Before or After Harvest	10	15	20
Split Application Before and After Harvest	5 + 5	7.5 + 7.5	10 + 10

Do not apply more than 10 pounds per acre on coarse soils, 15 pounds per acre on medium soils or 20 pounds per acre on fine soils during the calendar year.

BEAN Dry

Apply and incorporate Trilin 10G before planting in the spring.

Broadcast Rate Per Acre
Trilin 10G (pounds)

Annual Rainfall	Coarse	Medium	Fine
less than 20"	5	6.25	7.5
greater than 20"	5	7.5	10

On soils with 2-5% organic matter, use 7.5 pounds on coarse and medium soils and 10 pounds on fine soils. On soils with 5-10% organic matter, use 10 pounds on all soil texture types.

Fall Application

Trilin 10G may be applied and incorporated between October 15 and December 31. Apply at a broadcast rate of 5 pounds per acre on coarse soils, 6.25 to 7.5 pounds on medium soils, and 7.5 pounds on fine soils. Use lower rate in areas receiving less than 20" total annual rainfall or irrigation. NOTE: Established weeds must be destroyed prior to application.

BEAN Guar and Mungbean

Apply and incorporate Trilin 10G before planting.

Broadcast Rate Per Acre
Trilin 10G (pounds)

	Coarse	Medium	Fine
	5	7.5	7.5

On soils with 2-5% organic matter, use 7.5 pounds per acre.

BEAN Lima and Snap

Apply and incorporate before planting

Broadcast Rate Per Acre
Trilin 10G (pounds)

	Coarse	Medium	Fine
	5	5	7.5

On soils with 2-5% organic matter, use 7.5 pounds per acre

CARROT

Apply and incorporate Trilin 10G before planting.

Broadcast Rate Per Acre
Trilin 10G (pounds)

Annual Rainfall	Coarse	Medium	Fine
less than 20"	5	6.25	7.5
greater than 20"	5	7.5	10

On soils with 2-5% organic matter, use 7.5 pounds on coarse and medium soils and 10 pounds on fine soils. On soils with 5-10% organic matter, use 10 pounds on all soil texture types.

CASTOR BEAN

Apply and incorporate Trilin 10G before planting.

Broadcast Rate Per Acre
Trilin 10G (pounds)

Annual Rainfall	Coarse	Medium	Fine
less than 20"	5	6.25	7.5
greater than 20"	5	7.5	10

On soils with 2-5% organic matter, use 7.5 pounds on coarse and medium soils and 10 pounds on fine soils. On soils with 5-10% organic matter, use 10 pounds on all soil texture types.

CELERY

Apply as a soil incorporated treatment. Trilin 10G may be applied to direct seeded or transplant celery before planting, at planting or immediately after planting.

Broadcast Rate Per acre
Trilin 10G (pounds)

Annual Rainfall	Coarse	Medium	Fine
less than 20"	5	6.25	7.5
greater than 20"	5	7.5	10

On soils with 2-5% organic matter, use 7.5 pounds on coarse and medium soils and 10 pounds on fine soils. On soils with 5-10% organic matter, use 10 pounds on all soil texture types.

CHICORY/ENDIVE

Apply and incorporate Trilin 10G before planting.

Broadcast Rate Per Acre
Trilin 10G (pounds)

	Coarse	Medium	Fine
	5	7.5	10

On coarse and medium soils with 2-5% organic matter, use 7.5 pounds per acre. On fine soils with 2-5% organic matter, use 10 pounds per acre. On soils with 5-10% organic matter, use 10 to 12.5 pounds per acre on all soil textures.

COLE CROPS Broccoli, Brussels Sprout, Cabbage and Cauliflower

Direct Seeded

Apply and incorporate Trilin 10G before planting.

Broadcast Rate Per Acre
Trilin 10G (pounds)

	Coarse	Medium	Fine
	5	5	7.5

On soils with 2-5% organic matter, use 7.5 pounds per acre

Direct-seeded cole crops have exhibited marginal tolerance to recommended rates of Trilin 10G. Stunting or reduced stands may occur.

Transplant

Apply and incorporate Trilin 10G prior to transplanting

Broadcast Rate Per Acre
Trilin 10G (pounds)

	Coarse	Medium	Fine
	5	6.25	7.5
	5	7.5	10

On soils with 2-5% organic matter, use 7.5 pounds on coarse and medium soils and 10 pounds on fine soils. On soils with 5-10% organic matter, use 10 pounds on all soil texture types.

FIELD CORN, GRAIN SORGHUM and CORN FODDER, FORAGE and SILAGE

Trilin 10G may be applied as a postemergence treatment, allowing the use of a preemergence herbicide. Trilin 10G does not control established weeds. Do not apply Trilin 10G to sweet corn or corn grown for seed. Do not apply Trilin 10G as a preplant or preemergence treatment as crop injury may occur.

Soil Preparation

Cultivate before Trilin 10G application to insure loose, friable soil to remove established weeds and to cover the base of plants with soil. Set cultivation equipment to add approximately one inch of soil to the base of sorghum plants.

Application Directions

Apply and incorporate Trilin 10G as a postemergence treatment at the recommended rates for the soil texture when the crop is well established (8 inches or taller) or immediately after a cultivation, up to a height of 30 inches. Soil incorporation may be accomplished with only one pass of a sweep-type cultivator or a properly adjusted rolling cultivator. The sweep-type cultivator should have 3 to 5 sweeps per row middle and be operated at 6 to 8 mph. Set the middle sweeps so as to avoid exposing untreated soil. Adjust the incorporation tools to prevent crop injury.

Broadcast Rate Per Acre
Trilin 10G (pounds)

Annual Rainfall	Coarse	Medium	Fine
less than 20"	3.75	5	7.5
greater than 20"	5	7.5	10

Use the lower rates when weed pressure is light and the higher rates when weed pressure is heavy.

Apply 5 to 7.5 pounds per acre on coarse soils in Alabama, Florida, Georgia, North Carolina, South Carolina and Virginia to control fall panicum and Texas panicum.

COTTON

Apply and incorporate Trilin 10G before or at planting, immediately after planting and at layby or in the fall. Do not disturb cotton seed when incorporating Trilin 10G after planting (postplant).

Broadcast Rate Per Acre
Trilin 10G (pounds)

Annual Rainfall	Coarse	Medium	Fine
less than 20"	5	6.25	7.5
greater than 20"	5	7.5	10

On soils with 2-5% organic matter, use 7.5 pounds on coarse and medium soils and 10 pounds on fine soils. On soils with 5-10% organic matter, use 10 pounds on all soil texture types.

Layby

Trilin 10G can be applied and incorporated any time up to layby, but not less than 90 days before harvest. Direct layby applications onto the soil between the rows and beneath emerged cotton plants at the same rates as for preplant application. Soil incorporation using one pass of a sweep-type cultivator or properly adjusted rolling cultivator. Operate cultivation equipment at speeds sufficient to provide vigorous soil mixing and exercise care to avoid mechanical injury to the crop.

Fall Application

Trilin 10G may be applied and incorporated to cotton ground any time between October 15 and December 31. The ground may be left flat or bedded-up over winter. Where soil is left flat, take care not to turn up untreated soil from beds into furrows. On bedded grounds, knock down beds to desired heights before planting, moving some treated soil into furrows from the beds. Destroy established weeds during preparation of seedbed. Before planting, destroy weeds which may have become established in furrows due to the uncovering of untreated soil during bedding. Trilin 10G should not be applied to wet soil or soils which are subject to prolonged periods of flooding.

Broadcast Rates Per Acre - Fall Application Only: In Alabama, Arkansas, northern Florida, Georgia, Louisiana, Mississippi, southeastern Missouri, bootheel, North Carolina, New Mexico, Oklahoma, South Carolina, Tennessee and Texas apply and incorporate Trilin 10G at a broadcast rate of 10 pounds per acre on coarse and medium soils and 12.5 pounds on fine soils. In Arizona and Nevada, a broadcast rate of 7.5 pounds Trilin 10G per acre should be used on coarse soil, 10 pounds on medium soils and 12.5 pounds on fine soils.

For other states where cotton may be grown, apply Trilin 10G at a broadcast rate of 5 pounds per acre on coarse soils, 7.5 on medium soils, and 10 pounds on fine soils. For coarse soils with 2-5% organic matter use 7.5 pounds. For fine soils with 2-5% organic matter use 10 pounds. For soils with 5-10% organic matter use 10 to 12.5 pounds. NOTE: Established weeds must be destroyed prior to application.

Preemergence Overlay Application Following Trilin 10G

Apply Trilin 10G as a preplant incorporated treatment. Additional weeds tolerant to Trilin 10G may be controlled using overlay application of products registered for use on cotton. Such applications may be made unless use following a Trilin 10G application is specifically prohibited by the product label. Consult the overlay product label for additional weeds controlled, directions for use, precautions and limitations.

Postemergence Application Following Trilin 10G

Apply Trilin 10G as a preplant incorporated treatment. Additional weeds tolerant to Trilin 10G may be controlled using postemergence application of products registered for use on cotton. Such applications may be made unless use following a trifluralin application is specifically prohibited by the product label. Consult the postemergence product label for additional weeds controlled, directions for use, precautions and limitations.

COTTONWOOD TREES GROWN FOR PULP

New Plantings

Apply and incorporate Trilin 10G before planting

Broadcast Rate Per Acre
Trilin 10G (pounds)

Annual Rainfall	Coarse	Medium	Fine
less than 20"	5	6.25	7.5
greater than 20"	5	7.5	10

All soils with 2-5% organic matter use 7.5 to 10 pounds.
All soils with 5-10% organic matter use 10 pounds.

Established Plantings

Apply Trilin 10G in established plantings as a ground application. A single rainfall or overhead sprinkler irrigation of 0.5 inches or more or flood irrigation after application is required to activate Trilin 10G.

Broadcast Rate Per Acre
Trilin 10G (pounds)

All Soils

10-20

Application rate cannot be adjusted according to weed pressure.

Johnsongrass Suppression

Proper soil preparation before application is necessary for satisfactory results. Use incorporation methods not injurious to the crop.

Broadcast Rate Per Acre
Trilin 10G (pounds)

All Soils
20

Some johnsongrass plants will escape. Timely cultivation with tillage implements or spot spraying with effective postemergence herbicides will improve the level of johnsongrass control.

CUCURBITS

Cantaloupe, Cucumber and Watermelon

Postemergence Only

Apply and incorporate Trilin 10G as a directed application to the soil between the rows and beneath plants which are in the 3 to 4 true leaf stage.

Broadcast Rate Per Acre
Trilin 10G (pounds)

Annual Rainfall	Coarse	Medium	Fine
less than 20"	5	6.25	7.5
greater than 20"	5	7.5	10

On soils with 2-5% organic matter, use 7.5 pounds on coarse and medium soils and 10 pounds on fine soils. On soils with 5-10% organic matter, use 10 pounds on all soil texture types.

FLAX

Apply and incorporate Trilin 10G in the fall from September 1 to December 31 in Minnesota, North Dakota and South Dakota and between October 15 and December 31 in other states.

Broadcast Rates Per Acre

Trilin 10G (pounds)	Coarse	Medium	Fine
	5	7.5	10

Special Instructions

1. Incorporation operations or any other tillages performed in the spring prior to seeding should be relatively shallow so as to maintain a firm seedbed, and the seedbed should be packed just prior to seeding.
2. Seeding should be done with a press drill or hoe drill. Seed into a moist seedbed no more than 1 1/4 inches deep.
3. Flax should not be seeded until the seedbed has warmed up.

FORAGE LEGUMES

Forage Legumes Used as Cover Crops or in the Acreage Conservation Reserve Program

Apply and incorporate Trilin 10G as a preplant treatment.

Broadcast Rate Per Acre
Trilin 10G (pounds)

Annual Rainfall	Coarse	Medium	Fine
less than 20"	5	5	7.5
greater than 20"	5	7.5	7.5

PRECAUTION

If used under the Acreage Conservation Reserve Program, follow the more severe grazing restrictions imposed by the USDA Conservation Use Program. Consult the local ASCS committee or other State Agency to determine the period of the USC grazing restriction. Some crop stand reduction may occur with this use; however, reduced weed competition will allow establishment of a quality stand.

FRUIT, NUTCROPS, VINEYARDS

Apply and incorporate Trilin 10G, for new plantings of almond, apricot, citrus, fig, kiwifruit, peach, pecan, plum, prune and walnut trees, before planting.

Broadcast Rate Per Acre
Trilin 10G (pounds)

Annual Rainfall	Coarse	Medium	Fine
less than 20"	5	3.25 to 7.5	7.5
greater than 20"	5	7.5	10

All soils with 2-5% organic matter use 7.5-10 pounds per acre. All soils with 5-10% organic matter use 10 pounds per acre.

New Plantings of Vineyards

Apply and incorporate Trilin 10G, for new plantings in vineyards, before planting.

Broadcast Rate Per Acre
Trilin 10G (pounds)

Annual Rainfall	Coarse	Medium	Fine
less than 20"	5	7.5	15
greater than 20"	7.5	15	20

All soils with 2-10% organic matter use 15-20 pounds per acre. Do not use more than 10 pounds per acre on heat-treated grape rootings.

Application to Established Non-bearing and Bearing Citrus, Fruit and Nut Crops and Vineyards

Apply Trilin 10G for postplant applications on bearing or non-bearing, established plantings of vineyards and almond, apricot, grapefruit, lemon, nectarine, orange, peach, pecan, plum, prune, tangelo, tangerine and walnut trees. Apply Trilin 10G in established plantings to the soil and use incorporation methods not injurious to the trees or vines. Do not apply to vineyards within 60 days of harvest.

Broadcast Rate Per Acre
Trilin 10G (pounds)

All Soils

10 to 20
pounds

GREENS

Turnip Greens grown for processing;
Collard, Kale and Mustard Greens

Apply and incorporate Trilin 10G as a preplant treatment.

Broadcast Rates Per Acre
Trilin 10G (pounds)

Coarse	Medium	Fine
5	7.5	7.5

HOPS

Apply and incorporate Trilin 10G to established hops during dormancy. Incorporate once using incorporation equipment that will insure thorough soil mixing with minimal damage to crop stand.

Broadcast Rates Per Acre
Trilin 10G (pounds)

Coarse	Medium	Fine
5	6.25 to 7.5	6.25 to 7.5

On coarse and medium soils with 2-5% organic matter, use 7.5 pounds per acre.

MUSTARD

Grown for seed or processing for food in
Minnesota, North Dakota and South Dakota

Apply and incorporate Trilin 10G before planting.

Broadcast Rates Per Acre
Trilin 10G (pounds)

Coarse	Medium	Fine
5	7.5	7.5

OKRA

Apply and incorporate before planting.

Broadcast Rate Per Acre
Trilin 10G (pounds)

Annual Rainfall	Coarse	Medium	Fine
less than 20"	5	6.25	7.5
greater than 20"	5	7.5	10

On soils with 2-5% organic matter, use 7.5 pounds on coarse and medium soils and 10 pounds on fine soils. On soils with 5-10% organic matter, use 10 pounds on all soil texture types.

ONION

Grown for Dry Bulbs Only

A single rainfall or overhead sprinkler irrigation of 0.5 inches or more or flood irrigation after application is required to activate Trilin 10G. Do not apply within 60 days of harvest.

Broadcast Rates Per Acre
Trilin 10G (pounds)

Coarse	Medium
3.75 to 5	5 to 6.25

Use the lower rates where light weed pressure is anticipated.

Soil incorporation may be accomplished by operating sweep-type or rolling cultivators 2 to 4 inches deep at 5 to 8 mph. Incorporation equipment must mix Trilin 10G uniformly in the soil. Incorporate with two passes. The first pass must be within 24 hours of application or erratic weed control may result. Avoid covering exposed onion bulbs with treated soil during incorporation as injury to the crop may occur. Care should be taken to avoid injury to the roots during incorporation.

PRECAUTIONS

Applied according to these directions and under normal growing conditions, Trilin 10G will not adversely affect onion. Diseases, improper incorporation depth, excessive moisture, high salt concentration or drought may weaken the crop and increase the possibility of damage from Trilin 10G. Under these conditions, delayed crop development or reduced yields may result.

PEA

Dry and English

Apply and incorporate Trilin 10G in the spring before planting or in the fall.

Broadcast Rate Per Acre
Trilin 10G (pounds)

	Coarse	Medium	Fine
Spring Application	5	5	7.5
Fall Application*	5	6.25 to 7.5	7.5

* Trilin 10G may be fall applied to Dry and English peas in Idaho, Oregon and Washington.

PEA Southern

Apply and incorporate Trilin 10G before planting.

Broadcast Rate Per Acre
Trilin 10G (pounds)

Annual Rainfall	Coarse	Medium	Fine
less than 20"	5	6.25	7.5
greater than 20"	5	7.5	10

On soils with 2-5% organic matter, use 7.5 pounds on coarse and medium soils and 10 pounds on fine soils. On soils with 5-10% organic matter, use 10 pounds on all soil texture types.

PEANUT

Spanish Peanuts, Florunner and Florigiant
in New Mexico, Oklahoma and Texas

Trilin 10G may be applied and incorporated before or after planting. If applied after planting extreme care must be taken not to disturb the seed during incorporation.

Broadcast Rates Per Acre
Trilin 10G (pounds)

Coarse	Medium
5	7.5

PEPPER

Apply and incorporate Trilin 10G prior to transplanting.

Broadcast Rate Per Acre
Trilin 10G (pounds)

Annual Rainfall	Coarse	Medium	Fine
less than 20"	5	6.25	7.5
greater than 20"	5	7.5	10

On soils with 2-5% organic matter, use 7.5 pounds on coarse and medium soils and 10 pounds on fine soils. On soils with 5-10% organic matter, use 10 pounds on all soil texture types.

POTATOES

Apply and incorporate Trilin 10G after planting, prior to crop emergence, immediately following dragoff or after potato plants have fully emerged.

Set incorporation equipment so that the bed and furrow are uniformly covered with a layer of treated soil. If the layer of treated soil is not uniform and the herbicide is concentrated over the bed, potato emergence may be retarded and stem brittleness can occur. When applying and incorporating Trilin 10G after potato plants have fully emerged, do not completely cover the foliage with treated soil. Do not completely cover foliage during subsequent cultivations. Be careful that incorporation equipment does not damage seed pieces or elongating sprouts.

Broadcast Rate Per acre
Trilin 10G (pounds)

Annual Rainfall	Coarse	Medium	Fine
less than 20"	5	6.25	7.5
greater than 20"	5	7.5	10

On soils with 2-5% organic matter, use 7.5 pounds on coarse and medium soils and 10 pounds on fine soils. On soils with 5-10% organic matter, use 10 pounds on all soil texture types.

RADISH

Apply and incorporate Trilin 10G before planting.

Broadcast Rate Per Acre
Trilin 10G (pounds)

Coarse	Medium	Fine
5	7.5	7.5

RAPSEED (Canola)

Apply and incorporate Trilin 10G in the spring before planting or in the fall after September 1.

Broadcast Rates Per Acre
Trilin 10G (pounds)

Coarse	Medium	Fine
5	7.5	10

SAFFLOWER

Preemergence

Trilin 10G may be applied before planting in the spring or between October 15 and December 31.

Broadcast Rate Per Acre
Trilin 10G (pounds)

Annual Rainfall	Coarse	Medium	Fine
less than 20"	5	6.25 to 7.5	7.5
greater than 20"	5	7.5	10

On soils with 2-5% organic matter, use 7.5 pounds on coarse and medium soils and 10 pounds on fine soils. On soils with 5-10% organic matter, use 10 to 12 pounds on all soil texture types.

Fall Application in Arizona, California, Idaho, Montana, Nevada, Oregon, Utah, Washington and Wyoming.

Trilin 10G may be applied and incorporated between October 15 and December 31 and ground left flat or bedded during the winter. If bedded, knock down to proper height prior to planting mixing some treated soil from bed tops into furrows. If treated soil is left flat

BEST AVAILABLE COPY

for winter, exercise caution in bedding operations to avoid turning up untreated soil. Any weeds establishing as a result of bedding or other operations should be destroyed before planting. NOTE: Trilin 10G should not be applied in the fall to soils which are wet or subject to prolonged flooding.

Broadcast Rate Per Acre
Trilin 10G (pounds)

	Coarse	Medium	Fine
	7.5	10	12.5

SOYBEAN

Preemergence

Trilin 10G may be applied and incorporated before planting.

Broadcast Rate Per Acre
Trilin 10G (pounds)

Annual Rainfall	Coarse	Medium	Fine
less than 20"	5	6.25	7.5
greater than 20"	5	7.5	10

On soils with 2-5% organic matter, use 7.5 pounds on coarse and medium soils and 10 pounds on fine soils. On soils with 5-10% organic matter, use 10 pounds on all soil texture types.

Fall Application

Trilin 10G may be applied and incorporated between October 15 and December 31 and ground left flat or bedded during the winter. If bedded, knock down to proper height prior to planting moving some treated soil from bed tops into furrows. If treated soil is left flat for winter, exercise caution in bedding operations to avoid turning up untreated soil. Any weeds established as a result of bedding or other operations should be destroyed before planting.

Broadcast Rate Per Acre
Trilin 10G (pounds)

	Soil Type	States
10	Coarse and Medium	Alabama, northern Florida, Georgia, Louisiana, Mississippi, Missouri, North Carolina, Oklahoma, South Carolina, Tennessee and Texas
12.5	Fine	Same as for Coarse and Medium
7.5	Coarse and Medium with 2-5% organic matter	All Other States
10	Fine	All Other States
10 to 12.5	5-10% organic matter	All Other States

PRECAUTION

Do not fall apply Trilin 10G to soils which are wet or subject to prolonged periods of flooding, or where rice was grown the previous year.

Preemergence Overlay Applications Following Trilin 10G

Apply and incorporate Trilin 10G as a preplant treatment. Additional weeds tolerant to Trilin 10G may be controlled by using other products registered for use on soybeans. Such treatments may be made, unless use following a trifluralin application is specifically prohibited by the product label. Consult the overlay product label for additional weeds controlled, directions for use, cautions and limitations before use.

Postemergence Treatment Following Trilin 10G

Apply and incorporate Trilin 10G as a preplant treatment. Additional weeds tolerant to Trilin 10G may be controlled using postemergence applications of other products registered for use on soybeans. Such treatments may be made, unless use following a trifluralin application is specifically prohibited by the product label. Consult the overlay or postemergence product label for additional weeds controlled, directions for use, cautions and limitations before use.

SUGAR BEETS

Trilin 10G may be applied over-the-top of 2 to 6 inch tap plants, but only after any exposed sugar beet roots have been covered with soil. Incorporation equipment should be set to avoid injury to the sugar beet tap root while throwing treated soil toward the plants in the row.

Broadcast Rate Per Acre
Trilin 10G (pounds)

Annual Rainfall	Coarse	Medium	Fine
less than 20"	5	6.25	6.25
greater than 20"	5	7.5	7.5

SUGARCANE

Apply and incorporate Trilin 10G twice a year, making the first application in the fall on firmly packed beds immediately after the seed pieces are planted. Make a second application in the spring before or shortly after the cane emerges. Loosen rain-packed beds 2-3 inches deep before spring application. Take care that incorporation equipment does not damage the seed pieces or emerging shoots.

Broadcast Rate Per Acre
Trilin 10G (pounds)

All Soils

10 - 20

Application rate range may be adjusted to weed pressure.

Applications Up to Layby for Plant Cane or Ratoon Cane (for use in Louisiana and Texas)

Apply and incorporate in the spring from shortly before or after cane emergence until layby. Apply after beds have been shaved or false shaved. Loosen rain-packed beds 2-3 inches deep before application. Avoid incorporation equipment damage to seed pieces or emerging shoots. Incorporate with a rolling cultivator or bed chopper for all soil textures. Set rolling cultivator to cut 2 to 4 inches deep and operate at 6 to 8 mph. Set bed chopper to cut 3 to 4 inches deep and operate at 4 to 6 mph. Two incorporation passes are necessary.

Broadcast Rate Per Acre
Trilin 10G (pounds)

All Soils

10 - 20

Application rate range may be adjusted according to weed pressure.

Itchgrass (Raouigrass) Control

Apply and incorporate on plant or ratoon cane. Follow use directions in layby section.

Broadcast Rate Per Acre
Trilin 10G (pounds)

All Soils

10 - 20

Application rate range may be adjusted according to weed pressure.

SUNFLOWER

Trilin 10G may be applied and incorporated in the spring or in the fall between September 15 and December 31 in Minnesota, North Dakota and South Dakota, and between October 15 and December 31 in all other states.

Broadcast Rate Per Acre
Trilin 10G (pounds)

Annual Rainfall	Coarse	Medium	Fine
less than 20"	5	6.25	7.5
greater than 20"	5	7.5	10

On soils with 2-5% organic matter, use 7.5 pounds on coarse and medium soils and 10 pounds on fine soils. On soils with 5-10% organic matter, use 10 to 12 pounds on all soil texture types.

TOMATO

Direct Seeded/Transplanted

For direct seeded tomato apply at blocking or thinning to the soil between rows and beneath plants and incorporate. For transplanted tomato apply and incorporate prior to transplanting. Do not apply after transplanting.

Broadcast Rate Per Acre
Trilin 10G (pounds)

Annual Rainfall	Coarse	Medium	Fine
less than 20"	5	6.25	7.5
greater than 20"	5	7.5	10

On soils with 2-5% organic matter, use 7.5 pounds on coarse and medium soils and 10 pounds on fine soils. On soils with 5-10% organic matter, use 10 to 12 pounds on all soil texture types.

SMALL GRAINS — BARLEY DURUM & WHEAT

Special Precaution For Use of Trilin 10G on Small Grains

Follow carefully the Trilin 10G directions for use on small grains to minimize potential crop stress. Under certain conditions, delayed crop emergence and/or stand reduction may occur when Trilin 10G is applied to small grains. The combined effect of certain cultural practices and unfavorable soil or environmental conditions may cause excessive crop seedling stress resulting in retarded crop growth, stand reduction and possibly reduced yield.

For best results, observe the following cultural practices or precautions:

Use tillage methods that provide a uniformly firm seedbed and time tillage operations to conserve moisture.

Irrigate prior to planting or after germination and emergence. Moisture received between planting and emergence may cause crusting, especially on loose, friable seedbeds.

Do not exceed recommended application rates for Trilin 10G. This is particularly important on coarse textured or low organic matter soils.

Carefully follow incorporation directions. When applying preplant incorporated treatments, operate equipment at recommended depth and speed to place Trilin 10G into the upper 1 to 1.5 inches of soil. If applied after planting, set equipment so as to not disturb planted seed.

Set drills to place seed at the depth specified in use directions. A planting depth greater than 2.5 inches for spring wheat or durum will result in increased seedling stress and decreased emergence.

Where Trilin 10G is to be applied use only high quality seed (avoid use of small seed and low starch reserves).

If seed treatments are used, apply at the correct rate and uniformly across all seeds. Misapplication may result in reduced germination and/or seedling vigor.

Avoid use of seed varieties known to have poor seedling emergence vigor.

Do not apply Trilin 10G in the fall in combinations with any other preplant incorporated herbicide.

Soil characteristics and environmental conditions which may contribute to crop seedling stress that may be accentuated by use of Trilin 10G include:

Soil related: High salinity, eroded knolls/hilltops, loose dry soils and compaction.

Weather related: Cold and/or wet soils, excessively hot soils, excessive moisture, drought and soil crusting from heavy rainfall.

NOTE: Do not apply Trilin 10G on small grains where a dinitroaniline herbicide such as Trilin 10G was applied at a rate recommended for row crops (oil seeds) during the previous growing season.

Incorporation Instructions/Equipment

Special attention must be paid to soil incorporation because unless such is effectively done, Trilin 10G will not provide desired levels of weed control. Use incorporation equipment in the manner described below unless otherwise specified in incorporation instructions. Any of the following implements listed below may be used for the first incorporation of Trilin 10G. Use only the disc or field cultivator for the second incorporation pass and incorporate in a different direction. Poor weed control may result if untreated soil is moved to the surface during the second incorporation pass. To avoid this, the second incorporation should not be deeper than the first.

Tandem Disc: Operate at 3 to 4 inches deep, 4 to 6 mph.

Chisel Plow: A chisel plow is defined to have 3 rows of 1 1/2 to 2 inch sweeps on no greater than 12 inch centers. Sweeps should be staggered so that no soil is left unturned. Operate at 4 to 5 inches deep at 4 to 6 mph. NOTE: Do not use Chisel Plow.

Field Cultivator: A field cultivator is defined to have 3 to 4 rows of sweeps with C- or S-shaped chunks spread 7 inches or less, staggered to leave no soil unturned. Operate at 3 to 4 inches deep at 5 mph or more.

WHEAT, DURUM AND BARLEY

Spring Seeded — Summer Fallow Application

Apply and incorporate Trilin 10G to control labeled weeds during summer fallow as well as for fall (pigeon peas) where grains are seeded the subsequent spring. When applying Trilin 10G soil condition is important, application may be made only where trash levels are manageable, land has been fallow, pre-flooded or left with stubble cover.

BEST AVAILABLE COPY

Two incorporations are essential for weed control. The first incorporation should be made within 24 hours of application. The second and all subsequent tillage activities are to be conducted as required to destroy resistant weeds as present. NOTE: Control of labeled weeds may not occur prior to the second incorporation.

Broadcast Rate Per Acre
Triflin 10G (pounds)

Application Period	Less Than 10 Inches Annual Rainfall	More Than 10 Inches Annual Rainfall
April 15 to 30	8.75	10
May 1 to 31	8.75 to 7.5	10 to 8.75
June 1 to 30	7.5 to 6.25	8.75 to 7.5
July 1 to 31	6.25 to 5.0	7.5 to 6.25
August 1 to 31	5.0	6.25 to 5.0

Where a rate range is shown, use the higher rate per acre during the early part of an application period and the lower rate per acre during the later part of an application period.

Incorporation Instructions

See Incorporation Instructions/Equipment in Small Grains section above.

Seeding Directions

Seed should be planted approximately 2 inches deep.

PRECAUTION

Use of Triflin 10G in this manner may result in a stand reduction; generally slight stand reductions do not affect yield.

BARLEY

Spring Seeded — Fall Application Minnesota, North Dakota and South Dakota (except for special rate programs)

Apply in the fall for general weed control during the following growing season. Incorporate one time within 24 hours. Incorporate a second time before planting to destroy existing weeds and insure uniform distribution of Triflin 10G treated soil. For best control results, the second incorporation should occur at least 7 days after the first.

Broadcast Rate Per Acre
Triflin 10G (pounds)

	Coarse	Medium	Fine
	5	7.5	7.5

Incorporation Instructions

See Incorporation Instructions/Equipment in Small Grains section above.

Seeding Directions

Seed should be planted approximately 2 inches deep.

PRECAUTION

Use of Triflin 10G in this manner may result in a stand reduction; generally slight stand reductions usually do not affect yield.

BARLEY

Spring Seeded — Spring Application Minnesota, North Dakota and South Dakota

Apply in the spring as a preplant incorporated treatment in spring seeded barley for control of foxtail (pigeongrass). Incorporate one time within 24 hours. Incorporate a second time before planting to destroy existing weeds and insure uniform distribution of Triflin 10G in treated soil. For best weed control results, the second incorporation should occur at least 7 days after the first.

Broadcast Rate Per Acre
Triflin 10G (pounds)

	Coarse	Medium	Fine
	5	5	5

Incorporation Instructions

See Incorporation Instructions/Equipment in Small Grains section above.

Seeding Directions

Seed should be planted approximately 2 inches deep.

PRECAUTION

Use of Triflin 10G in this manner may result in a stand reduction; generally slight stand reductions usually do not affect yield.

BARLEY

Spring Seeded — Spring Application Used as a Cover Crop or in the Acreage Conservation Reserve Program Minnesota, North Dakota and South Dakota

Apply in the spring as a preplant incorporated treatment in spring seeded barley used as a cover crop or in acreage in the conservation reserve program for control of foxtail/pigeongrass. Incorporate one time within 24 hours. Incorporate a second time before planting to destroy existing weeds and insure uniform distribution of Triflin 10G in treated soil. For best weed control results, the second incorporation should occur at least 7 days after the first.

Broadcast Rate Per Acre
Triflin 10G (pounds)

	Coarse	Medium	Fine
	5	7.5	7.5

Incorporation Instructions

See Incorporation Instructions/Equipment in Small Grains section above.

Seeding Directions

Seed should be planted approximately 2 inches deep.

PRECAUTION

Use of Triflin 10G in this manner may result in a stand reduction; generally slight stand reductions usually do not affect yield. Follow the most severe grazing restrictions imposed by either the label for Triflin 10G or the USDA Acreage Conservation Reserve Program. Consult the local ARCS office or other state agency to determine the period of the USDA grazing restriction.

DURUM

Spring Preplant Incorporated North Dakota West of the Red River Valley and South Dakota

Apply Triflin 10G in the spring as a preplant incorporated treatment for suppression of green and yellow foxtail (pigeongrass) in spring seeded durum.

Application Rate

Apply at a broadcast rate of 4 pounds per acre for all soil types.

Incorporation Instructions

See instructions for chisel plow and field cultivator in the Small Grains section above.

Seeding Directions

Seed should be planted approximately 1.5 to 2 inches deep.

PRECAUTIONS

Use of Triflin 10G in this manner may result in a stand reduction; generally slight stand reductions usually do not affect yield.

WHEAT AND DURUM

Spring Seeded — Fall Application Foxtail (Pigeongrass) Control

Apply in the fall for foxtail (pigeongrass) control during the following growing season. Incorporate one time within 24 hours. Incorporate a second time before planting to destroy existing weeds and insure a uniform distribution of Triflin 10G in treated soil.

Broadcast Rate Per Acre
Triflin 10G (pounds)

	Coarse	Medium	Fine
	5	5	7.5

Incorporation Instructions

See Incorporation Instructions/Equipment in Small Grains section above.

Seeding Directions

Seed should be planted approximately 2 inches deep.

PRECAUTION

Use of Triflin 10G in this manner may result in a stand reduction; generally slight stand reductions usually do not affect yield.

WINTER WHEAT

Preplant Incorporated Colorado, Idaho, Kansas, Nebraska, Oregon, Washington and Wyoming

Apply Triflin 10G as a preplant incorporated treatment up to three weeks before planting to control labeled weeds including cheatgrass.

Broadcast Rate Per Acre
Triflin 10G (pounds)

	Coarse	Medium	Fine
	7.5	7.5	10

Incorporation Instructions

Triflin 10G should be shallowly incorporated with a flexible tine-tooth harrow set to cut one to two inches deep and operated at 3 to 6 mph. Two incorporation operations are required, the first within 24 hours after application and the second at least five days after the first. The second incorporation should be at a different direction to the first and prior to planting. NOTE: Do not work the soil with a disc between Triflin 10G incorporation with a flexible tine harrow and harvest of crop for which application is being made.

Seeding Instructions

Drill with deep furrow or semi-deep furrow equipment which places seed below soil zones into which Triflin 10G has been incorporated.

PRECAUTION

Do not plant wheat directly into the zone of soil treated with Triflin 10G as crop injury may occur. Delayed emergence or slight stand loss do not usually affect yield.

WINTER WHEAT

Fallow Soil Application Colorado, Idaho, Kansas, Nebraska, Oregon, Washington and Wyoming

Apply and incorporate Triflin 10G up to four months before planting to control labeled weeds including cheatgrass during the fallow period and during the following growing season. Apply any time from May to September prior to fall planting of winter wheat.

Broadcast Rate Per Acre
Triflin 10G

	Coarse	Medium	Fine
	7.5	7.5	10

Incorporation Instructions

Triflin 10G should be shallowly incorporated with a flexible tine-tooth harrow set to cut 1 to 2 inches deep and operated at 3 to 6 mph. Two incorporation operations are required, the first within 24 hours after application and the second at least five days after the first. The second incorporation should be at a different direction to the first and prior to planting. NOTE: Do not work the soil with a disc between Triflin 10G incorporation with a flexible tine harrow and harvest of crop for which application is being made.

Seeding Instructions

Drill with deep furrow or semi-deep furrow equipment which places seed below soil zones into which Triflin 10G has been incorporated.

PRECAUTION

Do not plant wheat directly into the zone of soil treated with Triflin 10G as crop injury may occur. Delayed emergence or slight stand loss do not usually affect yield.

WARRANTY STATEMENT

GRIFFIN CORPORATION warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, reduced liveliness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of GRIFFIN. In no case shall GRIFFIN be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. GRIFFIN MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

Triflin is a registered trademark of Griffin.

GRIFFIN
1000 000 000

11/11

Pot-in-Pot Nursery Production of Trees and Shrubs

For the control of excess rooting between the socket pot and the growing pot. Trilin 10G will reduce the rooting out from the growing container.

Apply Trilin 10G in the **socket** pot before the growing pot is inserted at the rate corresponding to the container size. Use higher rates for aggressive rooting species.

Pot Size (Nursery Gallons)	Grams Trilin 10G/Pot	Teaspoons/Pot
5	1.0 - 1.5	1/2
7	1.0 - 2.0	3/4
10	1.5 - 3.0	1
15	1.6 - 3.2	1 - 1.5
20	2.4 - 4.8	1 - 2
25	3.0 - 5.8	2 - 3
30	3.3 - 6.5	3 - 4