

NOV 1 1993

James Yowell
Griffin Corporation
P.O. Box 1847
Valdosta, GA 31603

Dear Mr. Yowell:

Subject: Increased Application Rate and New Crops
Kocide DF
EPA Registration No. 1812-334 ✓
Your Submission Dated October 12, 1993

The amendment referred to above, submitted in connection with registration under section 3(c)(7)(A) of the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), is acceptable provided that you:

1. Submit/cite all data required for registration/-reregistration of your product under FIFRA section 3(c)(5) or 4(a) when the Agency requires all registrants of similar products to submit such data.

2. Submit one (1) copy of your final printed labeling before you release the product for shipment. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

20727

-2-

This labeling does not bear the required Worker Protection statements as outlined in PR Notices 93-7 and 93-11. We note that you have submitted labels in accordance with the instructions in those Notices. The new protective language in those submissions acceptance on October 13, 1993 must be incorporated into this label for any shipment released after the deadlines that have been established. Note finished printed labels must also be submitted in accordance with the instructions of the WPS group.

Sincerely yours,

/s/

Cynthia Giles-Parker
Product Manager (22)
Fungicide-Herbicide Branch
Registration Division (7505C)

Enclosure

Jacket

30751

wps

10/06/93

KOCIDE DF

DRY FLOWABLE

FUNGICIDE/BACTERICIDE

Active Ingredient

Copper Hydroxide 61.4%

Inert Ingredients 38.6%

Total. 100.0%

ACCEPTED
with COMMENTS
in EPA Letter Dated:

(Metallic Copper Equivalent 40%)

NOV 1 1993

Under the Federal Insecticide
Fungicide, and Rodenticide Act
a product, for the pesticide
registered under EPA Reg. No.

DANGER - PELIGRO

**KEEP OUT OF REACH OF CHILDREN
FOR ADDITIONAL INFORMATION SEE
STATEMENT OF PRACTICAL TREATMENT**

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta haya sido explicado ampliamente.

Griffin Corporation
Valdosta, GA 31601

EPA Reg. No. 1812-334
EPA Est. No. 8901-TX-1

BEST AVAILABLE COPY

40431

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER - PELIGRO

Corrosive. Causes irreversible eye damage. Wear goggles, face shield or safety glasses. Harmful if swallowed, absorbed through the skin or inhaled. May cause skin sensitization reactions in certain individuals. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Protective clothing, including goggles, should be worn. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Hold eyelids open and flush with water for 15 minutes. Get medical attention.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

IF SWALLOWED: Drink promptly a large quantity of milk, egg white, gelatin solution, or if these are not available, large quantities of water. Avoid alcohol. Get medical attention.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate use of gastric lavage.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic organisms. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to fish and aquatic organisms in adjacent aquatic sites. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

RE-ENTRY STATEMENT

Do not enter treated areas without protective clothing until sprays have dried. Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. (Indicate specific oral warnings which inform workers of areas or fields that may not be entered without specific protective clothing, period of time field must be vacated and appropriate actions to take in case of accidental exposure). When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: DANGER. Area treated with KOCIDE DF on (date of application). Do not enter without appropriate protective clothing until spray has dried. In case of accidental exposure see Statement of Practical Treatment.

STORAGE AND DISPOSAL

Store in a cool, dry place.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in sanitary landfill, or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

100107

GENERAL INSTRUCTIONS

Use KOCIDE DF as noted below. KOCIDE DF is adaptable to spraying from aircraft and ground spraying equipment. Depending upon the equipment used and the specific crop, the volume applied per acre will differ. Refer to recommended volume table below.

Minimum Recommended Spray Volume (Gallons) Per Acre When Applying Kocide DF

	Aerial	Dilute	Concentrate
Vegetables	3	20	---
Field Crops	3	20	---
Small Fruits	5	150	50
Vines	5	150	50
Tree Crops	10	400	50
Citrus	10	800	100
			(50 Florida)
Miscellaneous	10	150	50
Turf (Algae control)			

NOTE: - Agricultural chemicals may perform in an unpredictable manner when tank mixed, especially where several products are involved. Reduced effect on pests or crop injury may occur. Unless recommended on this label or by a state/local expert, it is advisable to test for compatibility and tolerance to crop injury prior to full scale commercial utilization of a new tank mix or tank mixing should not be undertaken.

GENERAL CHEMIGATION INSTRUCTIONS

Apply this product only through one or more of the following types of systems: sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move irrigation system(s). Do not apply this product through any other type of irrigation system.

Crop injury or lack of effectiveness can result from nonuniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety device for public water

systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Posting of areas to be chemigated is required when 1) any part of a treated area is within 300 feet of sensitive areas such as residential areas, labor camps, businesses, day care centers, hospitals, in-patient clinics, nursing homes or any public areas such as schools, parks, playgrounds, or other public facilities not including public roads, or 2) when the chemigated area is open to the public such as golf courses or retail greenhouses.

) Posting must conform to the following requirements. Treated areas shall be posted with signs at all usual points of entry and along likely routes of approach from the listed sensitive areas. When there are no usual points of entry, signs must be posted in the corners of the treated areas and in any location affording maximum visibility to sensitive areas. The printed side of the sign should face away from the treated area towards the sensitive area. The signs shall be printed in English. Signs must be posted prior to application and must remain until foliage has dried and soil surface water has disappeared. Signs may remain in place indefinitely as long as they are composed of materials to prevent deterioration and maintain legibility for the duration of the posting period.

) All words shall consist of letters at least 2 1/2 inches tall, and all letters and the symbol shall be a color which sharply contrasts with their immediate background. At the top of the sign shall be the words KEEP OUT, followed by an octagonal stop sign symbol at least 8 inches in diameter containing the word STOP. Below the symbol shall be the words PESTICIDES IN IRRIGATION WATER. ✓

CHEMIGATION SYSTEMS CONNECTED TO PUBLIC WATER SYSTEMS

) Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into the reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

7-23

The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

) Do not apply when wind speed favors drift beyond the area intended for treatment.

) When mixing, fill nurse tank half full with water. Add KOCIDE DF slowly to tank while hydraulic or mechanical agitation is operating and continue filling with water. Stickers, spreaders, insecticides, nutrients, etc. should be added last, DO NOT PRE-MIX OR SLURRY. If compatibility is in question, use the compatibility jar test before mixing a whole tank. Because of the wide variety of possible combinations which can be encountered, observe all cautions and limitations on the label of all products used in mixtures.

KOCIDE DF should be added through a traveling irrigation system continuously or at the last 30 minutes of solid set or hand moved irrigation systems. Agitation is recommended.

SPRINKLER CHEMIGATION ✓

) The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

) The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution

907-11

is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

When mixing, fill nurse tank half full with water. Add KOCIDE DF slowly to tank while hydraulic or mechanical agitation is operating and continue filling with water. Do not premix Kocide DF in a slurry before adding to the nurse tank. Stickers, spreaders, insecticides, nutrients, etc. should be added last, DO NOT PRE-MIX OR SLURRY. If compatibility is in question, use the compatibility jar test before mixing a whole tank. Because of the wide variety of possible combinations which can be encountered, observe all cautions and limitations on the label of all products used in mixtures.

KOCIDE DF should be added through a traveling irrigation system continuously or at the last 30 minutes of solid set or hand moved irrigation systems. Agitation is recommended.

CROP CLASSIFICATION

CITRUS: Grapefruit, Kumquat, Lemon, Lime, Orange, Tangelo, and Tangerine.

FIELD CROPS: Alfalfa, Barley, Oats, Peanut, Potato, Sugar Beet, and Wheat.

SMALL FRUITS: Blackberry, Blueberry, Cranberry, Currant, Gooseberry, Raspberry, and Strawberry.

TREE CROPS: Almond, Apple, Apricot, Avocado, Banana, Cacao, Cherry, Coffee, Filbert, Mango, Nectarine, Olive, Peach, Pear, Pecan, Pistachio, Plum, Prune, Quince, and Walnut.

See 612-356

VEGETABLES: Beans, Broccoli, Brussels Sprouts, Cabbage, Cantaloupe, Carrot, Cauliflower, Celery, Cucumber, Eggplant, Endive, Escarole, Greens (Beet, Collard, Mustard and Turnip), Honeydew, Lettuce, Muskmelon, Onion, Pea, Pepper, Pumpkin, Spinach, Squash, Table Beet, Tomato, Watercress, and Watermelon.

VINES: Grape, Hops, and Kiwi.

MISCELLANEOUS: Atemoya, Carambola, Chives, Douglas Fir, Ginseng, Guava, Litchi, Live Oaks, Macadamia, Mamey Sapote, Papaya, Parsley, Passion Fruit, Sugar Apple, and Sycamore.

100-311

TURF: Algae control.

ORNAMENTALS: As listed.

KOCIDE DF may be applied as an aerial, ground dilute or ground concentrate spray unless specifically directed otherwise by crop.

When selecting a use rate for KOCIDE DF do not apply less than the label recommended minimum amount. Under heavy disease pressure or when conditions favor such, use the higher rate and shorter spray intervals specified for each crop. In addition, use the higher rates for large mature tree crops.

The per acre use rate of KOCIDE DF is applicable for both dilute and concentrate spraying. Consult the KOCIDE DF label for specific rates and timing of application by crop.

Complete spray coverage is essential to assure optimum performance from KOCIDE DF. When treating on a concentrate basis or by aerial application, unless you have had specific previous experience, it is advisable to test for compatibility and tolerance to crop injury prior to full scale commercial utilization.

While volume is important in obtaining full spray coverage, often factors such as foliage density, environmental conditions and spray calibrations, have a greater impact. Always be sure that sprayers are calibrated to spray equipment manufacturer's specifications and environmental conditions are within those recommended by State and local regulatory authorities.

When mixing, fill spray tank one-half full with water. Add KOCIDE DF slowly to tank while hydraulic or mechanical agitation is operating and continue filling with water. Spreaders, stickers (cleared for application to growing crops), nutrients, etc. should be added last. If compatibility is in question, use the compatibility jar test before mixing a whole tank.

NOTE: KOCIDE DF should not be applied in a spray solution having a pH of less than 6.5 as phytotoxicity may occur.

Environmental conditions such as extended periods of wet weather, acid rain, etc. which alter the pH of the leaf surface may affect the performance of KOCIDE DF resulting in possible phytotoxicity or loss of effectiveness.

The following specific instructions are based on general application procedures. The recommendations of the State Agricultural Extension Service should be closely followed as to timing, frequency, and number of sprays per season.

FROST INJURY PROTECTION

BACTERIAL ICE NUCLEATION INHIBITOR

11/28/51

Application of KOCIDE DF made to all crops listed on this label at rates and stages of growth indicated on this label, at least 24 hours prior to anticipated frost conditions, will afford control of ice nucleating bacteria (Pseudomonas syringae, Erwinia herbicola, and Pseudomonas fluorescens) and may therefore provide some protection against light frost. Not recommended for those geographical areas where weather conditions favor severe frost.

CITRUS

NOTE: Adding foliar nutritionals to spray mixtures containing Kocide DF or other products and applying to citrus during the post bloom period when young fruit is present may result in spray burn. Do not use KOCIDE DF on citrus seedlings grown in greenhouses or shade houses.

Disease	Rate/Acre	Use Instructions
Melanose, Scab,	4-12 lbs.	Apply as pre-bloom and post-bloom sprays.
Greasy Spot, Pink Pitting	2-6 lbs.	Apply in summer on expanded new flush. Repeat on subsequent flushes where disease pressure is severe. Use higher rates when conditions favor disease.
Brown Rot	4-8 lbs.	Begin application in fall and continue as needed. Apply to skirts of trees to a height of at least 4 feet. Apply also to bare ground one foot beyond skirt. Use higher rates when conditions favor disease. Make foliar applications to protect developing fruit in orchards with previous disease history or when conditions favor disease development.
NOTE: In California, in areas subject to copper injury, add 1/3 to 1 pound of high quality lime per pound of KOCIDE DF.		
Phytophthora	1 lb.	Mix with 1 gallon of water and paint trunks of trees from the soil surface to the lowest scaffold limbs. Apply in May prior to summer rains and/or in the fall prior to wrapping trees for freeze protection. Treatment serves for protection for up to one year, but does not cure existing infections.

Citrus Can .er 12 lbs.
(Suppression Only)

Spray flushes 7-14 days after shoots begin to grow. Young fruit may require an additional application. Number and timing of applications will be dependent upon disease pressure. Under heavy pressure, each flush of new growth should be sprayed.

FIELD CROPS

Crop	Disease	Rate/Acre	Use Instructions
Alfalfa	Cercospora and Lep- tosphaerulina Leaf Spots	2 lbs.	Apply 10-14 days before each harvest or earlier if disease threatens.
			NOTE: Spray injury may occur with sensitive varieties such as Lathontan.
Peanut	Cercospora Leaf Spot	1.5-3 lbs.	Begin spraying at 35-40 days after planting or when disease symptoms first appear and repeat at 10-14 day intervals as needed. Reduce sprays to 7 day intervals during humid weather. Use higher rates when conditions favor disease.
Potato	Early and Late Blight	1-4 lbs.	Apply 1 - 1.5 lbs. at 7-10 day intervals starting when plants are 6 inches high until 2 weeks before harvest in locations where disease is light and up to 3-4 pounds per acre where disease is more severe.
Sugar Beet	Cercospora Leaf Spot	2-5 lbs.	Begin applications when conditions first favor disease development and repeat at 10-14 day intervals as needed. Use the higher rate when disease is severe. Addition of a suitable agricultural spray oil is

Wheat, Oats
and Barley

Septoria Leaf
Blotch,
Helminthosporium
Spot Blotch

1.5-2 lbs.

recommended.

Make first application at early heading and follow with second spray 10 days later. Use the higher rates when conditions favor disease.

SMALL FRUITS

Crop	Disease	Rate/Acre	Use Instructions
Blackberry (Santiams, Logans, Boysens, Marions, Auroras, Cascades, Chehalems & Thornless Evergreens)	Leaf and Cane Spot	4 lbs.	Apply delayed dormant spray after training in the spring. Make fall application after harvest. Add 1 quart of crop oil per acre.
Blueberry	Bacterial Canker	6 lbs.	Make first application before fall rains and a second application four weeks later.
Cranberry	Fruit Rot	8 lbs.	Make first application in late bloom. One or two additional applications at 10-14 day intervals may be required depending upon disease severity.
	Bacterial stem canker	8 lbs.	Apply post harvest and again in spring before bud burst. One or two additional applications at 10-14 day intervals may be required depending upon disease severity.

1002

C u r r a n t , Gooseberry	Leaf Spot	10 lbs.	Make three applications starting after harvest followed by application before bloom and after petal fall.
-------------------------------	-----------	---------	---

Strawberry	leaf Spot and Leaf Blight	2-3 lbs.	Begin application when plants are established and continue on a weekly schedule throughout season
------------	---------------------------	----------	---

NOTE: Discontinue applications if signs of crop injury appear.

Raspberry	Leaf and Cane Spot	4 lbs	Apply delayed dormant spray after training in the spring. Make fall application after harvest. Add 1 quart of crop oil per acre.
-----------	--------------------	-------	--

TREE CROPS

Crop	Disease	Rate/Acre	Use Instructions
Almond	Coryneum Blight, Blossom Brown Rot	8-12 lbs.	Dormant application: Apply before foliage buds begin to swell. Use higher rates when rainfall is heavy and disease pressure is high.
		6-8 lbs.	Early bloom (popcorn) application: Apply before full bloom. Use higher rates when rainfall is heavy and disease pressure is high.
			NOTE: To avoid plant injury, do not use above rate after full bloom.

	Bacterial Blast (Pseudomonas)	12-16 lbs.	Apply at dormant to early pink bud. For control in sprinkler irrigated orchards or where disease is severe, apply 1 pound per acre at 2 week post-bloom intervals or just before sprinkling.
			NOTE: Injury may occur from post-bloom sprays, especially on Nephelium varieties.
Apple	Anthraxnose, European Canker, Pseudomonas	12-16 lbs.	Apply before fall rains. Use higher rates under severe disease conditions.
			NOTE: Use on yellow varieties may cause discoloration. To avoid, pick before spraying.
	Fireblight	8-16 lbs.	Make application between silver-tip and green-tip. Apply as a full cover spray.
			NOTE: Crop injury may occur from late application; discontinue use when green tip reaches 1/2 inch. ✓
	Crown or Collar Rot	4 lbs.	Mix in 100 gallons of water. Apply 4 gallons of suspension as a drench on the lower trunk area of each tree. Apply either in early spring or in late fall after harvest.
			NOTE: Do not use if soil pH is below 5.5 since copper toxicity may result.

227-1

Apricot	Coryneum Blight (Shot Hole), Blossom Brown Rot	8-12 lbs.	Apply at dormant to full bloom and use higher rates when conditions favor disease.
			NOTE: Applications applied after bloom will result in crop injury.
Avocado	Scab	8-12 lbs.	Apply when bloom buds begin to swell and continue application at monthly intervals for 5-6 applications. Use higher rate when conditions favor disease.
Banana	Sigatoka	2 lbs.	Apply by air in 3 gallons of water combining 0.5 gallons of agricultural oil. Apply on a 14 day schedule throughout the wet season. Apply at 21 day intervals during dry periods.
	Black Pitting	4 lbs.	Mix in 100 gallons of water (4 pounds per acre) directly to the fruit stem and include the basal portion of the leaf crown. Apply during the first and second weeks after fruit emergence.
Cacao	Black Pod	2-8.5 lbs.	Begin applications at the start of the rainy season and continue while infection conditions persist. Apply 2 - 4.5 lbs. as often as 14 to 21 days in high rainfall areas at varying rates depending on disease severity. For drier areas, where 2 to 4 applications are recommended during critical infection periods and at long intervals, use 8.5 pounds per acre, according to disease incidence and planting density.

7/27

Cherry	Dead Bud (<i>Pseudomonas syringae</i>), Coryneum Blight	8-12 lbs.	Make first application in fall before heavy rains and a second at late dormant. In orchards where the disease is severe, a spray should also be applied shortly after harvest. Add 1 pint of superior-type oil per 100 gallons of water as a dilute spray.
	Brown Rot, Blossom Blight	8-12 lbs.	Apply a full cover spray at popcorn stage and a second application at full bloom.
Coffee	Coffee Berry Disease (<i>Colletotrichum coffeanum</i>)	6-8 lbs.	Apply first spray after flowering and before onset of long rains and then at 21-28 day intervals until picking. Use higher rates when rainfall is heavy and disease pressure is high.
	Bacterial Blight (<i>Pseudomonas syringae</i>)	6-8 lbs.	Begin spray program before the onset of the long rains and continue throughout the rainy season at 14-21 day intervals. The critical time of spraying to control this disease is just before, during and after flowering(s) especially when coinciding with wet weather. Use higher rates when rainfall is heavy and disease pressure is high. ✓
	Leaf Rust (<i>Hemileia vastatrix</i>)	2-4 lbs.	Apply before the onset of rain and then at 21 day intervals while the rains continue. Use higher rates when rainfall is heavy and disease pressure is high.

12/27

	Iron Spot (<u>Cercospora coffeicola</u>), Pink Disease (<u>Corticium salmonicolor</u>)	2 lbs.	Use concentrate or dilute spray. Begin treatment at the start of wet season and continue at monthly intervals for three applications.
Filbert	Bacterial Blight	16-24 lbs.	Apply as a post harvest spray. In seasons of heavy rainfall apply a second spray when three- fourths of the leaves have dropped. Add 1 pint of superior- type oil per 100 gallons of water.
	Eastern Filbert Blight	16-24 lbs.	Apply as a dilute spray in adequate water for thorough coverage. Make initial application after harvest in October before heavy winter rains begin. The next application should be made in late February to early March followed by an other application one month later. If desired, add 1 pint of a sticking agent or superior type oil per 100 gallons of water.
Mango	Anthraxnose	8 lbs.	Apply monthly after fruit set until harvest.
Olive (California)	Peacock Spot Olive Knot	8-12 lbs.	Make first application before winter rains fall. A second application in early spring should be made if disease is severe. Apply the high rate for heavy disease pressure or when conditions favor disease development. ✓
Peach, Nectarine	Leaf Curl, Coryneum Blight (Shot Hole)	8-16 lbs.	Apply at leaf fall to late dormant. Use the highest rate when rainfall is very heavy and disease pressure is high. May be used with agricultural spray oil.

Brown Rot, Blossom Blight	8-12 lbs.	Full cover spray at pink bud. Application at this time affords some control of Leaf Curl and Coryneum Blight.
---------------------------	-----------	---

Bacterial Spot	1 lb.	Post bloom application applied at first and second cover sprays.
----------------	-------	--

	8 lbs.	Apply as a dormant spray.
--	--------	---------------------------

NOTE: Do not spray 3 weeks prior to harvest. Use only recommended rates. Spotting of leaves and defoliation may occur from use in cover sprays.

) Pear

Fire Blight	1 lb.
-------------	-------

Apply at 5 day intervals throughout the bloom period.

Pseudomonas Blight	12-16 lbs.
--------------------	------------

Apply before fall rains and again during dormancy before spring growth starts. Use the higher rate when disease pressure is high or when conditions favor disease development. ✓

NOTE: Do not apply to copper sensitive varieties. Excessive dosages may cause fruit russet.

20.43

Pecan	Shuck & Kernel Rot (<u>Phytophthora cactorum</u>) and Zonate Leafspot (<u>Cristulariella pyramidalis</u>)	2-4 lbs.	For suppression, apply in sufficient water to ensure complete spray coverage at 2 to 4 week intervals starting at kernel growth and continuing until shucks open. Use the higher rate and shorter interval if frequent rainfall occurs.
Plum & Prune	Coryneum Blight (Shot Hole)	8-16 lbs.	Apply as a dormant spray. Use the higher rate when rainfall is heavy and/or disease pressure is high.
	Brown Rot, Blossom Blight	8-12 lbs.	Apply full cover application at pink, red or early white bud stage. Use the higher rate when disease pressure is heavy or conditions favor disease development. ✓
Pistachio	Botrytis Blight, Botryosphaeria panicle and shoot blight, Septoria leaf blight, late blight (<u>Alternaria alternata</u>)	4 - 8 lbs.	Make initial application at bud swell and repeat on a 14 - 28 day schedule as dictated by disease conditions. If disease conditions are severe, use the high rate and short spray interval.
Quince	Fire Blight	1 lb.	Apply at 5 day intervals throughout the bloom period. †

Walnut

Walnut Blight

8-12.5 lbs.

Apply first application at delayed dormant or at early pre-bloom prior to catkins partial expansion. Make additional applications during bloom and early nutlet stage or as needed when free moisture occurs.

NOTE: Will not control copper tolerant strains of Xanthomonas bacteria.

VEGETABLES

This product can be mixed with 1.1 - 1.6 quarts/acre MANEX™ or other maneb (Manganese Ethylenebisdithiocarbamate) product to provide improved disease control on peppers for bacterial spot, and Tomato (field or green house) for bacterial spot, anthracnose, gray leaf mold, and septoria leaf spot. These mixes must be used in accordance with the most restrictive label limitations and precautions. No label dosage rates should be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing.

Crop	Disease	Rate/Acre	Use Instructions
Beans	Bacterial Blight (Halo & Common), Brown Spot	1-3 lbs.	Use the higher rate for more severe disease. For protective sprays, make first application when plants are 6 inches high; repeat on a 7-14 day schedule depending upon environmental conditions. ✓
Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Greens, (eg. Beet Collard, Mustard, & Turnip)	Black Rot (Xanthomonas) & Black Leaf Spot (Alternaria) Downy Mildew	1 - 2 lbs.	Apply at 7-10 day intervals. For control of disease of these crops, begin application after transplants are set in the field, or shortly after emergence of field seeded crops or when conditions favor disease development. Use shorter intervals and higher rates when conditions favor disease.

BEST AVAILABLE COPY

20457

NOTE: Reddening of older leaves may occur on Broccoli and a flecking of wrapper leaves may occur on Cabbage.

Lettuce, Endive, Escarole	Downy Mildew	1 - 2 lb.	Begin treatment when disease first appears or when conditions favor disease development and repeat every 7 - 10 days as needed to suppress disease.
---------------------------------	--------------	-----------	---

NOTE: Flecking and/or yellowing of leaves may occur under certain environmental conditions such as extended periods of moist weather.

Cantaloupe, Honeydew, Muskmelon	Downy Mildew	2 lbs.	Begin application when conditions are favorable for disease development and repeat at 5-7 day intervals as needed depending on disease severity.
---------------------------------------	--------------	--------	--

Carrot	Cercospora Leaf Spot	2 lbs.	Begin applications when disease first threatens and repeat at 7-14 day intervals as needed depending on disease severity.
--------	----------------------	--------	---

Celery	Cercospora Early Blight, Septoria Late Blight, & Bacterial Blight	2 lbs.	Begin applications as soon as plants are first established in the field, repeating at 5-7 day intervals depending on disease severity and environmental conditions.
--------	---	--------	---

Cucumber	Angular Leaf Spot and Downy Mildew	1.5-2 lbs.	Apply weekly when plants begin to vine.
----------	------------------------------------	------------	---

BEST AVAILABLE COPY

Box 37

Eggplant	Alternaria Blight, Anthracnose, & Phomopsis	2 lbs.	Begin applications prior to development of disease symptoms. Repeat sprays at 7-10 day intervals or as needed depending on disease severity.
Onion	Purple Blotch, Downy Mildew &	2 lbs.	Begin when plants are 4-6 inches high and repeat at 7-10 day intervals as needed depending upon disease pressure. Can cause phytotoxicity to leaves.
Pea	Powdery Mildew	1.5-3 lbs.	Begin applications when disease symptoms first appear and repeat at weekly intervals as needed. Use higher rate for more severe disease.
Pepper	Bacterial Spot	2-3 lbs.	Begin applications when conditions first favor disease development and repeat at 5-10 day intervals as needed depending on disease severity. Use higher rates for severe disease.

NOTE: Disease control will be improved by adding 1.1 - 1.6 quarts/acre MANEX™ or other maneb product in accordance with the instructions provided above. The maximum application rate WEST of the Mississippi River is 1.6 lb. ai. With a maximum of six applications. East of the Mississippi maximum application rate is 2.4 lb. ai, and not more than six applications per season. When MANEX™ is used in tank mixture with KOCIDE DF, do not apply within 7 days of harvest.

Pumpkin Squash	&	Powdery Mildew	1.5-3 lbs.	Begin applications when plants are 3 weeks old or when disease symptoms first appear and repeat at 7 day intervals as needed to maintain control. Use the higher rates if disease is heavy or conditions favor disease development.
-------------------	---	----------------	------------	---

Spinach		Anthracnose, White Rust, Blue Mold, Cercospora leaf spot	2 - 3 lbs.	Begin application when disease first appears or when conditions favor disease development. Repeat at 7-10 day intervals as needed.
---------	--	--	------------	--

NOTE: Flecking may occur on Spinach leaves.

Table Beet		Cercospora Leaf Spot	2-5 lbs.	Begin applications when conditions first favor disease development and repeat at 10-14 day intervals as needed. Use the higher rate when disease is severe. Addition of a suitable agricultural spray oil is recommended.
------------	--	----------------------	----------	---

Tomato		Early & Late Blight	2-3 lbs.	Begin when disease first threatens and repeat at 7-10 day intervals or as needed depending on disease severity. Use higher rate for severe disease.
--------	--	---------------------	----------	---

		Bacterial Speck	2 lbs.	Begin applications when disease first threatens and repeat at 7-10 day intervals or as needed depending on disease severity.
--	--	-----------------	--------	--

		Bacterial Spot, Anthracnose, Gray	2-4 lbs.	Begin applications when disease first threatens and repeat at 7-10
--	--	-----------------------------------	----------	--

Leaf Mold, Septoria
Leaf Spot

25071

day intervals or as needed depending on disease severity. Use higher rate for severe disease.

NOTE: Disease control will be improved by adding 1.1 - 1.6 quarts/acre of MANEX or other maneb product in accordance with the instructions provided above. The maximum application rate WEST of the Mississippi River is 1.6 lb. ai. With a maximum of four applications. East of the Mississippi maximum application rate is 2.4 lb. ai, and not more than 7 applications per season. When MANEX™ is used in tank mixture with KOCIDE DF, do not apply within 5 days of harvest.

Watercress Cercospora leafspot 2 lbs.

Begin application when plants are first established in the field; repeating at 7 - 14 day intervals depending on disease severity and environmental conditions. Do not exceed 4 applications per crop. Apply using ground spray equipment at no less than 50 gallons of spray solution per acre.

Watermelon Anthracnose,
Bacterial Fruit Bloch
(suppression) &
Downy Mildew 2 lbs.

Apply as soon as plants become established in the field and repeat at weekly intervals as needed depending upon disease severity. Use higher rates when conditions favor disease.

VINES

Crops	Disease	Rate/Acre	Use Instructions
Grape	Black Rot, Powdery Mildew, Downy Mildew	2 lbs.	Begin applications at late dormant up to bud break with subsequent applications throughout the season depending upon disease severity. NOTE: Foliage injury may occur on copper sensitive varieties such as Concord, Delaware, Niagara, and Rosetts. Either test for sensitivity or add 1-3 pounds of hydrated lime per pound of KOCIDE DF.
Hops	Downy Mildew	2 lbs.	Make crown treatment after pruning, but before training. After training, additional treatments are needed at about 10 day intervals. ✓
			NOTE: Discontinue use 2 weeks before harvest.
Kiwi	<u>Pseudomonas syringae</u> , <u>Erwinia herbicola</u> , <u>Pseudomonas fluorescens</u>	8 lbs.	Apply in 200 gallons of water per acre. Make applications on a monthly basis. A maximum of 3 applications may be made.

MISCELLANEOUS

Crop	Disease	Rate/Acre	Use Instructions
------	---------	-----------	------------------

2437

Atemoya	Anthracnose	3 lbs.	Make initial application just before flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage.
Carambola	Anthracnose	6 lbs.	Make initial application just before flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage.
Chives	Downy Mildew	2 lbs	Begin applications when plants are established in the field. Repeat applications every 7-10 days as dictated by disease conditions. If disease pressure is high, use the shorter spray interval.
Douglas Fir	Rhabdoclone needlecast	2 lbs	Begin applications at bud break and repeat at 3-4 week intervals. Apply in a tank mix with another registered pesticide if moderate to severe disease pressure is present. ✓
Ginseng	Alternaria Leaf and Stem Blight	2.6 lbs.	Use as a tank mix with 2 pounds Rovral® 50W in 100 gallons of water. Begin KOCIDE-Rovral applications as soon as plants have emerged in spring.

BEST AVAILABLE COPY

58737

Applications should be repeated every 7 days until plants become dormant in fall. If schedule application is to be made before a rain shower, apply fungicides at least 8 hours before the rain, giving the fungicides time to dry on the plants. Use of a spreader-sticker or sticker is advised.

NOTE: Alternaria Leaf and Stem Blight is most severe in humid conditions such as those found in the dense canopies to 2-, 3- 4-year old Ginseng. It is very important that the stems be thoroughly covered with fungicide; therefore, use a spray apparatus which distributes the fungicide throughout the canopy.

Guava	Anthracnose, Red Algae	3 lbs.	Make initial application just before flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage.
Litchi	Anthracnose	3 lbs.	Make initial application just before flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage.
Live Oak, Pecan (Texas & Florida)	Ball Moss	6 lbs.*	Apply 6 lbs per 100 gallons* of water, in the spring when ball moss is actively growing, using 1.5 gallons of spray per foot of tree height. Make sure to wet ball moss tufts thoroughly. A second application may be required after 12 months.

NOTE: KOCIDE DF may be

BEST AVAILABLE COPY

injurious to ornamentals grown under Live Oaks. This product may be reactive on metal and masonry surfaces such as galvanized roofing. Avoid contact with metal surfaces. Do not spray on cars, houses, lawn furniture, etc.

Macadamia	Anthracnose	6 lbs.	Initiate sprays at first sign of flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage.
	Phytophthora Blight (<u>P. capsici</u>), Raceme Blight (<u>Botrytis cinerea</u>)	4.5-6 lbs.	Apply during raceme development and bloom periods. Apply in sufficient water for thorough coverage.
Mamey Sapote	Anthracnose, Algal Leaf Spot	6-8 lbs.	Apply when conditions favor disease development. Repeat on 14-30 day schedule as disease severity and environmental conditions dictate. ✓
Papaya	Anthracnose	4-10 lbs.	Apply before disease appears. Apply at 10-14 day intervals under light disease pressure and at 5-7 day intervals under heavy disease pressure. The addition of an approved spreader is desirable.
Parsley	Bacterial blight (<u>Pseudomonas</u> sp.)	3 lbs.	Begin applications when plants are first established in the field and repeat at 5-7 days intervals depending upon disease severity and environmental conditions.

35437

Passion Fruit	Anthracnose	6 lbs.	Make initial application just before flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage.
Sugar Apple (Annona)	Anthracnose	12 lbs.	Make initial application just before flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage.
Sycamore	Anthracnose	2-3 lbs.	Apply as a full cover spray. Apply in 100 gallons of water or sufficient volume for thorough coverage. Make first application at bud crack and second application 7-10 days later at 10% leaf expansion.

GREENHOUSE AND SHADE HOUSE CROPS

Notice to User: KOCIDE DF may be used in greenhouses and shade houses to control diseases on some crops which appear on this label. The grower should bear in mind that the sensitivity of crops grown in greenhouses and shade houses differ greatly from crops grown under field conditions. Neither the manufacturer nor seller has determined whether or not KOCIDE DF can be used safely on all greenhouse and shade house-grown crops. The user should determine if KOCIDE DF can be used safely prior to commercial use. In a small area, apply the recommended rates to the plants in question, i.e. foliage, fruit, etc., and observe for 7-10 days for symptoms of phytotoxicity prior to commercial use.

Apply KOCIDE DF according to specific rates given for those crops in pounds per acre or pounds per 100 gallons. One level tablespoons of KOCIDE DF per 1000 square feet is equivalent to 1 pound per acre. One half tablespoon of KOCIDE DF per gallon of water is equivalent to 1 pound per 100 gallons. KOCIDE DF should be applied in adequate water for thorough coverage of plant parts. Begin application at first sign of disease and repeat at 7-14 day intervals as needed; use shorter interval during periods when severe disease conditions persist.

210837

<u>Crop</u>	<u>Disease</u>	<u>Rate Per 1000 Sq Ft</u>	<u>Use Instructions</u>
Eggplant	Alternaria Blight, Anthracnose, & Phomopsis	2 TBSP	Begin applications prior to development of disease symptoms. Repeat sprays at 7-10 day intervals or as disease pressure dictates.
Pepper	Bacterial Spot	2-3 TBSP	Begin applications when conditions first favor disease development and repeat at 5-10 day intervals as needed depending on disease severity. Use higher rates for severe disease.
Cucumber	Angular Leaf Spot and Downy Mildew	1.5-2 TBSP	Apply weekly when plants begin to vine.
Tomato	Early & Late Blight	2-3 TBSP	Begin when disease first threatens and repeat at 7-10 day intervals or as needed depending on disease severity. Use higher rate for severe disease.
	Bacterial Speck	2 TBSP	Begin applications when disease first threatens and repeat at 7-10 day intervals or as needed depending on disease severity.
	Bacterial Spot, Anthracnose, Gray Leaf Mold, Septoria Leaf Spot	2-4 TBSP	Begin applications when disease first threatens and repeat at 7-10 day intervals or as needed depending on disease severity. Use higher rate for severe disease.

NOTE: Disease control will be improved by adding MANEX or

320431

other maneb product in accordance with the instructions provided above. When MANEX™ is used in tank mixture with KOCIDE DF, do not apply within 5 days of harvest.

TURFGRASS

To control algae in turfgrass, apply 1½ pounds KOCIDE DF per 1,000 square feet in 5 gallons of water. KOCIDE DF may be used alone or in combination with other registered fungicides as a maintenance spray. Observe all precautions and limitations on the label of each product used in tank mixes.

NOTE: **Phytotoxicity** may occur depending upon varietal differences. Apply the recommended rate to a small area and observe for 7-10 days for signs of injury. If phytotoxicity occurs, discontinue use. Do not apply in spray solutions with a pH of less than 6.5.

ORNAMENTALS

Notice to User: Plant sensitivities to KOCIDE DF have been found to be acceptable in specific genera and species listed on this label, however, phytotoxicity may occur. Due to the large number of species and varieties of ornamentals and nursery plants, it is impossible to test every one for sensitivity to KOCIDE DF. Neither the manufacturer nor seller has determined whether or not KOCIDE DF can be safely used on ornamental or nursery plants not listed on this label. The user should determine if KOCIDE DF can be used safely prior to commercial use. In a small area, apply the recommended rates to the plants in question, i.e. bedding plants, foliage, etc., and observe for 7-10 days for symptoms of phytotoxicity prior to commercial use.

Use KOCIDE DF on container, bench or bed-grown ornamentals in greenhouses or outdoor nurseries, for professional use on ornamentals grown for indoor and outdoor landscaping, and for control of bacterial and fungal diseases of foliage, flowers and stems.

One-half tablespoon of KOCIDE DF per gallon of water is equivalent to 1 pound per 100 gallons.

Apply as a thorough coverage spray using 1 pound KOCIDE DF per 100 gallons of water. Begin application at first sign of disease and repeat at 7-14 day intervals as needed; use shorter interval during periods of frequent rains or when severe disease conditions persist.

KOCIDE DF may be used alone or in combination with other fungicides as a maintenance spray. Observe all precautions and limitations on the label of each product used in tank mixes.

Crop	LatinName	Disease
Althea (Rose of Sharon)	<u>Hibiscus syriacus</u>	Bacterial Leaf Spot
Aralia	<u>Dizygotheca elegantissima</u>	Xanthomonas & Cercospora Leaf Spots, Alternaria
Arborvitae	<u>Thuja sp.</u>	Alternaria Twig Blight, Cercospora Leaf Blight
Azalea 1/	<u>Rhododendron sp.</u>	Cercospora Leaf Spot, Botrytis Blight, Phytophthora Dieback, Powdery Mildew
Begonia	<u>Begonia semperflorens</u>	Bacterial Leaf Spot (Xanthomonas sp., Erwinia sp., Pseudomonas sp.)
Bougainvillea	<u>Bougainvillea spectabilis</u>	Anthracnose, Bacterial Leaf Spot
Bulbs (Tulip, Gladiolus)	Miscellaneous	Anthracnose, Botrytis Blight
Camphor Tree	<u>Cinnamomum camphora</u>	Pseudomonas Leaf Spot
Carnation 1/	<u>Dianthus sp.</u>	Alternaria Blight, Pseudomonas Leaf Spot, Botrytis Blight
Camelia	<u>Camellia japonica</u> , <u>C. sasangua</u>	Anthracnose, Bacterial Leaf Spot
Canna	<u>Canna sp</u>	Pseudomonas Leaf Spot

3407-11

Chinese Tallow Tree	<u>Sapium sebiferum</u>	Bacterial Leaf Spot, (<u>Xanthomonas</u> sp., <u>Pseudomonas</u> sp.)
Chrysanthemum 1/	<u>Chrysanthemum</u> <u>morifolium</u>	Septoria Leaf Spot, Botrytis Blight
Cotoneaster	<u>Cotoneaster</u> sp.	Botrytis Blight
Dahlia	<u>Dahlia pinnata</u>	Alternaria Leaf Spot, Botrytis Gray Mold, Cercospora Leaf Spot
Date Palm	<u>Phoenix canariensis</u>	Pestalotia Leaf Spot
Dianthus	<u>Dianthus</u> sp.	Bacterial Spot, Bacterial Soft Rot
Dogwood	<u>Cornus florida</u>	Anthracnose
Dusty Miller	<u>Senecio cineraria</u>	Bacterial Leaf Spot (<u>Pseudomonas cichorii</u>)
Easter Lily 2/	<u>Lilium longiflorum</u>	Botrytis Blight
Echinacea	<u>Echinacea</u> sp.	Bacterial Leaf Spot (<u>Pseudomonas cichorii</u>)
Elm "Drake"	<u>Ulmus parvifolia</u>	Xanthomonas Leaf Spot
Euonymus	<u>Euonymus</u> sp.	Botrytis Blight, Anthracnose
European Fan Palm	<u>Champaerops numilis</u>	Pestalotia Leaf Spot
Gardenia	<u>Gardenia jasminoides</u>	Alternaria Leaf Spot, Botrytis Bud Rot, Cercospora Leaf Spot
Geranium	<u>Pelargonium</u> sp.	Alternaria Leaf Spot, Botrytis Gray Mold, Cercospora Leaf Spot
Gladiolus	<u>Gladiolus</u> sp.	Alternaria Leaf Spot

2031

		Botrytis Gray Mold, Bacterial Leaf Blight ;
Golden Rain Tree	<u>Koelreuteria paniculata</u>	Bacterial Leaf Spot
Hibiscus	<u>Hibiscus rosa-sinensis</u>	Bacterial Leaf Spot
Holly Fern	<u>Cyrtomium falcatum</u>	Pseudomonas Leaf Spot
Impatiens	<u>Impatiens salierana</u>	Bacterial Leaf Spot
India Hawthorn 3/	<u>Raphiolepis indica</u>	Anthracnose, Entomosporium Leafspot
Ivy (English, Algerian) 1/	<u>Hendera helix</u> , <u>H. canariensis</u>	Xanthomonas Leaf Spot
Ixora	<u>Ixora coccinea</u>	Xanthomonas Leaf Spot
Juniper (Eastern Red Cedar)	<u>Juniperus virginiana</u>	Anthracnose
Lantana	<u>Lantana camara</u>	Bacterial Leaf Spot
Lilac	<u>Syringa</u> sp.	Cercospora Leaf Spot
Loblolly Bay	<u>Gordonia lasianthus</u>	Anthracnose
Loquat	<u>Eriobotrya japonica</u>	<u>Entomosporium maculata</u> , <u>Colletotrichum</u> sp.
Mandevillas	<u>Mandevilla</u> sp.	Anthracnose
Magnolia (Southern)	<u>Magnolia grandiflora</u>	Algal Leaf Spot, Anthracnose, Bacterial Leaf Spot
Magnolia (Sweet Bay)	<u>Magnolia virginiana</u>	Anthracnose
Magnolia	<u>Magnolia soulangeana</u>	Bacterial Leaf Spot
Marigold	<u>Taraxacum</u> sp.	Alternaria Leaf Spot, Botrytis Leaf and Flower

Page 11

Mulberry, Weeping	<u>Morus alba</u>	Rot, Cercospora Leaf Spot
Oleander	<u>Nerium oleander</u>	Bacterial Leaf Spot Bacterial Leaf Spot, Fungal Leaf Spot
Oak, Laurel	<u>Quercus laurifolia</u>	Algal Leaf Spot (<u>Cephaleuros virescens</u>)
Pachysandra	<u>Pachysandra procumbens</u>	Volutella Leaf Blight
Pansy	<u>Viola Sp.</u>	Downy mildew
Pear (Flowering)	<u>Pyrus calleryana</u>	Fireblight, Leaf Spot
Peony	<u>Paeonia spp.</u>	Botrytis blight
Pentas (Egyptian Star)	<u>Pentas spp.</u>	Bacterial Leaf Spot (<u>Xanthomomas sp.</u>)
Periwinkle	<u>Catharanchus roseus</u> , <u>Vinca sp.</u>	Phomopsis Stem Blight ✓
Phlox	<u>Phlox sp.</u>	Alternaria Leaf Spot
Pistachio	<u>Pistacia chinensis</u>	Anthracnose
Plantain lily	<u>Hosta sp.</u>	Bacterial Leaf Spot
Powder Puff Plant	<u>Callindra sp.</u>	Bacterial Leaf Spot
Philodendron	<u>Philodendron selloum</u>	Bacterial Leaf Spot
Photinia ("Red Top", "Red Leaf")	<u>Photinia fraserii</u> , <u>P. glabra</u>	Anthracnose, Entomosporium
Pyracantha	<u>Pyracantha sp.</u>	Fireblight, Scab
Queen Palm	<u>Arecastrum</u> <u>romanzoffianum</u>	Exosporium Leaf Spot; Phytophthora Bud Rot
Rhododendron	<u>Rhododendron sp.</u>	Alternaria Flower Spot

3-8-83

Rose 1/	<u>Rosa</u> sp.	Powdery Mildew, Black Spot
Verbena	<u>Verbena</u> sp	Xanthomonas Leafspot
Viburnum	<u>Viburnum odoratissimum</u> , <u>V. suspensum</u>	Anthracnose
Washingtonia Palm	<u>Washingtonia robusta</u>	Pestalotia Leaf Spot
Weeping Willow	<u>Salix babylonica</u>	Anthracnose
Yucca (Adam's needle)	<u>Yucca</u> sp.	Cercospora and Septoria Leaf Spot

(1) Discoloration of foliage and/or blooms have been noted on some varieties. To prevent residues on commercial plants, do not spray just before selling season.

(2) Apply 3-5 pounds of Kocide DF in 20 - 100 gallons of water per acre.

(3) For Indian Hawthorn use 2 to 4 lbs. per 100 gallons or 1 to 2 level tablespoons per gallon.

NOTICE OF WARRANTY

GRIFFIN warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of GRIFFIN. In no case shall GRIFFIN be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. GRIFFIN MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

Rovral® is a registered trademark of Rhone-Poulenc.
Manex™ is a trademark of Griffin Corporation.