

1812-350

4/24/2003

1/2



RESTRICTED USE PESTICIDE

Because of the high acute toxicity of triphenyltin hydroxide and its potential for affecting fetal development. For retail sale to and use by Certified Applicators or persons under their direct supervision, and only for those uses covered by the Certified Applicators certification.

SUPPLEMENTAL LABELING FOR CONTROL OF VARIOUS DISEASES FOR PECAN IN ALL STATES EXCEPT CALIFORNIA AND ARIZONA

**Super Tin 80WP®
Fungicide AGPAK® Water Soluble Pack
EPA Reg. No. 1812-350**

Active Ingredient:

Triphenyltin hydroxide.....	80.0%
Inert Ingredients.....	<u>20.0%</u>
Total:.....	100.0%

Contains 6.0 ounces triphenyltin hydroxide per 7.5 ounce pack.
Contains 20.0 ounces of triphenyltin hydroxide per 25.0 ounce pack.
Contains 1.87 pounds triphenyltin hydroxide per 2.34 pound pack.

**KEEP OUT OF REACH OF CHILDREN
DANGER - PELIGRO**



(Skull on red background)

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

IMPORTANT: All application directions, restrictions, and precautions on the Super Tin 80WP package label and the package label for any tank mix partner must be followed. This labeling contains supplemental instructions and restrictions for use of this product which may not appear on the package label. Follow instructions carefully when using this product.

NOTE: This label must be in the possession of the applicator at the time of application.

EXPIRATION DATE: This supplemental label expires on December 31, 2004. These directions for Super Tin 80WP will appear on package labels at the next product label reprinting.

©2002
SuperTin and AGPAK is a trademark of Griffin Corporation.
Griffin and Design are a registered trademark of Griffin Corporation.

Griffin L.L.C.
P.O. Box 1847 • Valdosta, GA 31603-1847
2509 Rocky Ford Rd. • Valdosta, GA 31601
Telephone: (229) 242-8635
Facsimile: (229) 244-5813

ACCEPTED

APR 24 2003

Under the Federal Insecticide
Fungicide, and Rodenticide
as amended, for the pesticide
registered under
EPA Reg. No. 1812-350

Instructions:

Crop	Disease	Dry oz./Acre	Use Directions																
Pecan	Scab, Brown Leaf Spot, Downy Spot, Powdery Mildew, Liver Spot, Sooty Mold, Leaf Blotch	5.0-7.5 (7.5-5.0 acres/2.34 lb pack or 1.5-1.0 acres/7.5 oz. pack or 5.0-3.3 acres/25 oz. pack)	<p>Ground (Enclosed Cabs Only): Apply in sufficient water to provide for full coverage.</p> <p>Aerial: Apply in a minimum of 20 gallons of water.</p> <p>Diluted spray should be directed to all parts of the tree. Application should begin at pre-pollination stages when young leaves are unfolding, and a second application made when the small nuts are forming. Apply a maximum of nine treatments during a single growing season at 2 to 4 week intervals as needed to maintain control. Use the lower rate for the first two applications or until the disease becomes severe or during dry weather. Use the higher rate during wet weather or during severe scab, powdery mildew or other disease infections. Do not apply within 30 days of harvest. Apply a maximum seasonal use rate of 30 ounces of Super Tin 80WP for pecans grown west of Interstate 35. Apply a maximum seasonal use rate of 45 ounces of Super Tin 80WP for pecans grown east of Interstate 35.</p>																
	Scab control	<p>3.75 ounces of Super Tin + 16 ounces of Dodine 65WP</p> <p>Conversions for combining Super Tin packages and Dodine:</p> <table border="1" data-bbox="516 1081 938 1312"> <thead> <tr> <th>Number of Super Tin Packages</th> <th>Pkg Size</th> <th>Ounces of Dodine</th> <th>Acres Treated</th> </tr> </thead> <tbody> <tr> <td>1</td> <td>7.5 oz</td> <td>32 (2 lb)</td> <td>2</td> </tr> <tr> <td>1</td> <td>2.34 lb.</td> <td>160 (10 lb)</td> <td>10</td> </tr> <tr> <td>1</td> <td>25.0 oz.</td> <td>106 (6.6 lb)</td> <td>6.6</td> </tr> </tbody> </table>	Number of Super Tin Packages	Pkg Size	Ounces of Dodine	Acres Treated	1	7.5 oz	32 (2 lb)	2	1	2.34 lb.	160 (10 lb)	10	1	25.0 oz.	106 (6.6 lb)	6.6	<p>Do NOT divide individual water-soluble packages.</p> <p>Use intact water soluble packages of Super Tin. During agitation, add the package(s) of Super Tin and sufficient Dodine 65WP to maintain a ratio of 3.75 ounces of Super Tin to 16 ounces of Dodine 65WP per acre treated. Apply at 10 to 21 day intervals across the growing season to control scab.</p> <p>Ground (Enclosed Cabs Only): Apply in sufficient water to provide for full coverage.</p> <p>Aerial: Apply in a minimum of 20 gallons of water.</p> <p>Diluted spray should be directed to all parts of the tree.</p> <p>Observe all label prohibitions concerning tank-mixing of Super Tin and Dodine with other materials.</p> <p>Do not apply after shucks have started to open or within 30 days of harvest, whichever occurs first. Apply a maximum seasonal use rate of 30 ounces of Super Tin 80WP for pecans grown west of Interstate 35. Apply a maximum seasonal use rate of 45 ounces of Super Tin 80WP for pecans grown east of Interstate 35.</p>
Number of Super Tin Packages	Pkg Size	Ounces of Dodine	Acres Treated																
1	7.5 oz	32 (2 lb)	2																
1	2.34 lb.	160 (10 lb)	10																
1	25.0 oz.	106 (6.6 lb)	6.6																

GS1012203
Page 2 of 2