

harvest. Apply in sufficient water for thorough coverage.

Pecan, Live Ball Moss 6 lbs.*

Apply 6.0 lbs per 100 gallons* of water, in the spring when ball moss is actively growing, using 1.5 gallons of spray per foot of tree height. Make sure to wet ball moss tufts thoroughly. A second application may be required after 12 months.

NOTE: KOCIDE GX306 may be injurious to ornamentals grown under Live Oaks. This product may be reactive on metal and masonry surfaces such as galvanized roofing. Avoid contact with metal surfaces. Do not spray on cars, houses, lawn furniture, etc.

Macadamia Anthracnose 4.5 lbs.

Initiate sprays at first sign of flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage.

Phytophthora Blight (P. capsici), Raceme Blight (Botrytis cinerea)

3.5-4.5 lbs

Apply during raceme development and bloom periods. Apply in sufficient water for thorough coverage. Use higher rates when conditions favor disease.

Mamey Sapote	Anthrachnose, Algal Leaf Spot	4.5-6 lbs.	Apply when conditions favor disease development. Repeat on 14-30 day schedule as disease severity and environmental conditions dictate. Use higher rates when conditions favor disease.
Papaya	Anthracnose	3-7.5 lbs.	Apply before disease appears. Apply at 10-14 day intervals under light disease pressure and at 5-7 day intervals under heavy disease pressure. The addition of an approved spreader is desirable. Use higher rates when conditions favor disease.
Parsley	Bacterial Blight (<u>Pseudomonas</u> sp.)	2.25 lbs.	Begin applications when plants are first established in the field and repeat at 5-7 days intervals depending upon disease severity and environmental conditions.
Passion Fruit	Anthrachnose	4.5 lbs.	Make initial application just before flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage.
Sugar Apple (<u>Annona</u>)	Anthrachnose	9 lbs.	Make initial application just before flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage.
Sycamore	Anthrachnose	1.5-2.25 lbs.	Apply as a full cover spray. Apply in 100 gallons of water or sufficient volume for thorough coverage. Make first application at bud crack and second application 7-10 days later at 10% leaf expansion. Use higher rates when conditions favor disease.

TURFGRASS

To control algae in turfgrass, apply 1.0 pound KOCIDE GX 306 per 1,000 square feet in 5 gallons of water. KOCIDE GX 306 may be used alone or in combination with other registered fungicides as a maintenance spray. Observe all precautions and limitations on the label of each product used in tank mixes.

NOTE: **Phytotoxicity** may occur depending upon varietal differences. Apply the recommended rate to a small area and observe for 7-10 days for signs of injury. If phytotoxicity occurs, discontinue use. Do not apply in spray solutions with a pH of less than 6.5.

GREENHOUSE AND SHADE HOUSE CROPS

Notice to User: KOCIDE GX306 may be used in greenhouses and shade houses to control diseases on some crops which appear on this label; specific instructions have been included for certain crops. The grower should bear in mind that the sensitivity of crops grown in greenhouses and shade houses differ greatly from crops grown under field conditions. Neither the manufacturer nor seller has determined whether or not KOCIDE GX306 can be used safely on all greenhouse and shade house-grown crops. The user should determine if KOCIDE GX306 can be used safely prior to commercial use. In a small area, apply the recommended rates to the plants in question, i.e. foliage, fruit, etc., and observe for 7-10 days for symptoms of phytotoxicity prior to commercial use.

Apply KOCIDE GX306 according to specific rates given for those crops in pounds per acre or pounds per 100 gallons. **One level tablespoon of KOCIDE GX306 per 1000 square feet is equivalent to 1 pound per acre.** KOCIDE GX306 should be applied in adequate water for thorough coverage of plant parts. Begin application at first sign of disease and repeat at 7-14 day intervals as needed; use shorter interval during periods when severe disease conditions persist

NOTE: Do not use GX306 on citrus seedlings grown in greenhouses or shadehouses.

<u>Crop</u>	<u>Disease</u>	<u>Rate Per 1000 Sq Ft</u>	<u>Use Instructions</u>
Cucumber	Angular Leaf Spot and Downy Mildew	1.25-1.5 TBSP	Apply weekly when plants begin to vine. Use higher rates when conditions favor disease.
Eggplant	Alternaria Blight, Anthracnose, & Phomopsis	1.5 TBSP	Begin applications prior to development of disease symptoms. Repeat sprays at 7-10 day intervals or as disease pressure dictates.

Pepper

Bacterial Spot

1.5-2.25
TBSP

Begin applications when conditions first favor disease development and repeat at 5-10 day intervals as needed depending on disease severity. Use higher rates for severe disease.

Tomato

Anthracnose,
Bacterial Speck,
Bacterial Spot, Early
Blight, Grey Leaf
Mold, Late Blight,
Septoria Leaf Spot

1.5-2.25
TBSP

Begin when disease first threatens and repeat at 7-10 day intervals or as needed depending on disease severity. Use higher rate for severe disease.

ORNAMENTALS

Notice to User: Plant sensitivities to KOCIDE GX306 have been found to be acceptable in specific genera and species listed on this label, however, phytotoxicity may occur. Due to the large number of species, widely varying growth conditions, and varieties of ornamentals and nursery plants, it is impossible to test every one for sensitivity to KOCIDE GX306. Neither the manufacturer nor seller has determined whether or not KOCIDE GX306 can be safely used on ornamental or nursery plants not listed on this label. The user should determine if KOCIDE GX306 can be used safely prior to commercial use. In a small area, apply the recommended rates to the plants in question, i.e. bedding plants, foliage, etc., and observe for 7-10 days for symptoms of phytotoxicity prior to commercial use.

Use KOCIDE GX306 on container, bench or bed-grown ornamentals in greenhouses or outdoor nurseries, for professional use on ornamentals grown for indoor and outdoor landscaping, and for control of bacterial and fungal diseases of foliage, flowers and stems.

One half tablespoon of KOCIDE GX306 per gallon of water is equivalent to 1.0 pound per 100 gallons.

Apply as a thorough coverage spray using 0.75 pound KOCIDE GX306 per 100 gallons of water. Begin application at first sign of disease and repeat at 7-14 day intervals as needed; use shorter interval during periods of frequent rains or when severe disease conditions persist.

KOCIDE GX306 may be used alone or in combination with other fungicides as a maintenance spray. Observe all precautions and limitations on the label of each product used in tank mixes.

NOTE: Do not tank mix Kocide GX306 with Aliette® fungicide unless appropriate precautions have been taken to buffer the spray solution. Severe phytotoxicity may result if adequate precautions are not taken.

Crop	Latin Name	Disease
Althea (Rose of Sharon)	<u>Hibiscus syriacus</u>	Bacterial Leaf Spot
Aralia	<u>Dizygotheca elegantissima</u>	Xanthomonas & Cercospora Leaf Spots, Alternaria
Arborvitae	<u>Thuja</u> sp.	Alternaria Twig Blight, Cercospora Leaf Blight
Azalea 1/	<u>Rhododendron</u> sp.	Cercospora Leaf Spot, Botrytis Blight, Phytophthora Dieback, Powdery Mildew (
Begonia	<u>Begonia semperflorens</u>	Bacterial Leaf Spot (Xanthomonas sp., Erwinia sp., Pseudomonas sp.)
Bougainvillea	<u>Bougainvillea spectabilis</u>	Anthracnose, Bacterial Leaf Spot
Bulbs (Tulip, Gladiolus)	Miscellaneous	Anthracnose, Botrytis Blight /
Camphor Tree	<u>Cinnamomum camphora</u>	Pseudomonas Leaf Spot (
Carnation 1/	<u>Dianthus</u> sp	Alternaria Blight, Pseudomonas Leaf Spot, Botrytis Blight
Camelia	<u>Camellia japonica</u> , <u>C. sasangua</u>	Anthracnose, Bacterial Leaf Spot
Canna	<u>Canna</u> sp.	Pseudomonas Leaf Spot
Chinese Tallow Tree	<u>Sapium sebiferum</u>	Bacterial Leaf Spot (Xanthomonas sp., Pseudomonas sp.)



Chrysanthemum <u>1/</u>	<u>Chrysanthemum morifolium</u>	Septoria Leaf Spot, Botrytis Blight
Cotoneaster	<u>Cotoneaster sp.</u>	Botrytis Blight
Dahlia	<u>Dahlia pinnata</u>	Alternaria Leaf Spot, Botrytis Gray Mold, Cercospora Leaf Spot
Date Palm	<u>Phoenix canariensis</u>	Pestalotia Leaf Spot
Dianthus	<u>Dianthus sp.</u>	Bacterial Spot, Bacterial Soft Rot
Dogwood	<u>Cornus florida</u>	Anthracnose
Dusty Miller	<u>Senecio cineraria</u>	Bacterial Leaf Spot (<u>Pseudomonas cichorii</u>)
Easter Lily <u>2/</u>	<u>Lilium longiflorum</u>	Botrytis Blight
Echinacea	<u>Echinacea sp.</u>	Bacterial Leaf Spot (<u>Pseudomonas cichorii</u>)
Elm "Drake"	<u>Ulmus parvifolia</u>	Xanthomonas Leaf Spot
Euonymus	<u>Euonymus sp.</u>	Botrytis Blight, Anthracnose
European Fan Palm	<u>Champaerops numilis</u>	Pestalotia Leaf Spot
Gardenia	<u>Gardenia jasminoides</u>	Alternaria Leaf Spot, Botrytis Bud Rot, Cercospora Leaf Spot
Geranium	<u>Pelargonium sp.</u>	Alternaria Leaf Spot, Botrytis Gray Mold, Cercospora Leaf Spot
Gladiolus	<u>Gladiolus sp.</u>	Alternaria Leaf Spot, Botrytis Gray Mold, Bacterial Leaf Blight
Golden Rain Tree	<u>Koelreuteria paniculata</u>	Bacterial Leaf Spot

34 9 36


Hibiscus	<u>Hibiscus rosa-sinensis</u>	Bacterial Leaf Spot
Holly Fern	<u>Cyrtomium falcatum</u>	Pseudomonas Leaf Spot
Impatiens	<u>Impatiens sallerana</u>	Bacterial Leaf Spot
India Hawthorn 3/	<u>Raphiolepis indica</u>	Anthracnose , Entomosporium Leafspot
Ivy (English, Algerian) 1/	<u>Hendera helix</u> , <u>H. canariensis</u>	Xanthomonas Leaf Spot
Ixora	<u>Ixora coccinea</u>	Xanthomonas Leaf Spot
Juniper (Eastern Red Cedar)	<u>Juniperus virginiana</u>	Anthracnose
Lantana	<u>Lantana camara</u>	Bacterial Leaf Spot
Lilac	<u>Syringa</u> sp.	Cercospora Leaf Spot
Loblolly Bay	<u>Gordonia lasianthus</u>	Anthracnose
Loquat	<u>Eriobotrya japonica</u>	<u>Entomosporium maculata</u> , <u>Colletotrichum</u> sp.
Mandevillas	<u>Mandevilla</u> sp.	Anthracnose
Magnolia (Southern)	<u>Magnolia grandiflora</u>	Algal Leaf Spot, Anthracnose, Bacterial Leaf Spot
Magnolia (Sweet Bay)	<u>Magnolia virginiana</u>	Anthracnose
Magnolia	<u>Magnolia soulangiana</u>	Bacterial Leaf Spot
Marigold	<u>Tagetes</u> sp.	Alternaria Leaf Spot, Botrytis Leaf and Rot, Cercospora Leaf Spot
Mulberry, Weeping	<u>Morus alba</u>	Bacterial Leaf Spot

Oleander	<u>Nerium oleander</u>	Bacterial Leaf Spot, Fungal Leaf Spot
Oak, Laurel	<u>Quercus laurifolia</u>	Algal Leaf Spot (<u>Cephaleuros virescens</u>)
Pachysandra	<u>Pachysandra procumbens</u>	Volutella Leaf Blight
Pansy	<u>Viola Sp.</u>	Downy mildew
Pear (Flowering)	<u>Pyrus calleryana</u>	Fireblight, Leaf Spot
Peony	<u>Paeonia sp.</u>	Botrytis blight
) Pentas (Egyptian Star)	<u>Pentas sp.</u>	Bacterial Leaf Spot (<u>Xanthomonas sp.</u>)
Periwinkle	<u>Catharanthus roseus</u> , <u>Vinca sp.</u>	Phomopsis Stem Blight
Phlox	<u>Phlox sp.</u>	Alternaria Leaf Spot
Pistachio	<u>Pistacia chinensis</u>	Anthracnose
Plantain lily	<u>Hosta sp.</u>	Bacterial Leaf Spot
Powder Puff Plant	<u>Callindra sp.</u>	Bacterial Leaf Spot
, Philodendron	<u>Philodendron selloum</u>	Bacterial Leaf Spot
Photinia ("Red Top", "Red Leaf")	<u>Photinia fraserii</u> , <u>P. glabra</u>	A n t h r a c n o s e , Entomosporium
Pyracantha	<u>Pyracantha sp.</u>	Fireblight, Scab
Queen Palm	<u>A r e c a s t r u m</u> <u>romanzoffianum</u>	Exosporium Leaf Spot, Phytophthora Bud Rot
Rhododendron	<u>Rhododendron sp.</u>	Alternaria Flower Spot
Rose 1/	<u>Rosa sp.</u>	Powdery Mildew, Black Spot

ACCEPTED

MAR 24 1995

11682-35



P.S.T. 6

ACTIVE INGREDIENTS: A coordination product of zinc ion

manganese ethylene bisdithiocarbamate..... 6.00%

In which the ingredients are:

Manganese++.....1.20%

Zinc ++.....0.15%

Ethylene bisdithiocarbamate ion (C₂H₄N₂S₄).....4.65%

INERT INGREDIENTS:.....94.00%

TOTAL.....100.00%

EPA Reg. No. 11682-35

EPA Est. No. 8917-ID-1

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

IF ON SKIN wash with soap and water. If irritation persists, get medical attention.

IF IN EYES flush with water for at least 15 minutes. Get medical attention.

IF INHALED remove victim to fresh air. Get medical attention.

PRECAUTIONARY STATEMENTS

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS
HARMFUL IF TAKEN INTERNALLY**

AVOID INHALING DUST

AVOID CONTACT WITH SKIN, EYES AND CLOTHING

May be harmful if swallowed, inhaled or absorbed through skin. May cause irritation of eyes, nose, throat and skin. Do not rub eyes or mouth with hands.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear: coveralls over long-sleeved shirt and long pants, waterproof gloves, shoes plus socks, and protective eyewear. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not contaminate water by cleaning of equipment or disposal of wastes. Cover or incorporate spilled treated seed.

Do not allow this product to drift onto neighboring crops or use in a manner different from the label directions. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

STORAGE: Store in a cool, dry place. Avoid contact with floor and moisture.

DISPOSAL: Completely empty bag into application equipment. Then dispose of in a sanitary landfill, or by incineration, or if allowed by State and local authorities, by burning. If burned stay out of smoke.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls over long-sleeved shirt and long pants, waterproof gloves, shoes plus socks and protective eyewear.

POTATO SEED-PIECE TREATMENT

P.S.T. 6 Potato Seed Treater aids in the suppression of fusarium seed-piece decay, the most common seed and soilborne fungus disease. The use of high quality disease free seed carefully selected and handled, is the first consideration in obtaining best seed performance.

APPLICATION RATE: Use 1 lb. of P.S.T. 6 Potato Seed Treater per 100 lbs. of cut seed. Apply dust so that the entire seed-piece surface is thoroughly covered by the correct amount of fungicide dust. Prevent inhalation of dust by use of exhaust fans during treatment. Subject the seed potatoes to temperatures of 50 degrees to 55 degrees for 10 to 15 days before handling or cutting.

Before starting the cutting operation, thoroughly clean and sanitize all equipment which will contact the seed-pieces such as: Cutting Equipment including: cutting machines, knives, tables and all other associated cutting equipment; Planting Equipment including: planters, truck beds and all other associated planting equipment.

Plant treated seed immediately after cutting into soil sufficiently warm and moist to promote good wound healing. Storage of cut seed can potentially increase the incidence of decay. Consult local University or Agricultural Extension Agents for specific recommendations.

NOTICE: P.S.T. 6 Potato Seed Treater will not control bacterial decay such as blackleg and soft rot, or diseases present within the seed.

DO NOT USE TREATED SEED FOR FEED OR FOOD

DISCLAIMER: Seller warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the DIRECTIONS FOR USE. It is impossible to eliminate all risks associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of factors such as weather conditions, presence of other materials, soil conditions, or the manner of use or application, all of which are beyond the control of seller. All such risks shall be assumed by the user. Any damages arising from a breach of this warranty shall be limited to direct damages and shall not include consequential commercial damages such as loss of profits or values or any other special or indirect damages. Seller makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY.

JRS 3107 (10/94)

Simplex

J.R. SIMPLOT COMPANY

P.O. Box 198
Lathrop, CA 95330

P.O. Box 15057
Boise, ID 83715

Reg # 1812-358

PM-22 Jacket 1936



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg.
Number:

1812-358

Date of Issuance:

MAR 29 1995

NOTICE OF PESTICIDE:

☒ Registration
☐ Reregistration

(under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

Kocide GX306

Name and Address of Registrant (include ZIP Code):

Griffin Corporation
P.O. Box 1847
Valdosta, GA 31603-1847

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.

2. Make the following label changes listed below before you release the product for shipment:

- a. Add the phrase, "EPA Reg. No. 1812-358".
- b. In the ingredient declaration correct Inert Ingredients to read 46.2%.
- c. Update the Environmental Hazards Statement to include the following:

Do not contaminate water by disposal of equipment washwaters.

Signature of Approving Official:

/s/

Date:

MAR 29 1995

1436

22 36
page 2

EPA Reg. No. 1812-358

d. Delete the "Non-Agricultural Use Requirements" box or add statements that are protective for the re-entry of non-agricultural workers.

3. Submit one (1) copy of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Cynthia Giles-Parker
Product Manager (22)
Fungicide-Herbicide Branch
Registration Division (7505C)

Enclosure

3936
✓

ACCEPTED
WITH COMMENTS
In EPA Letter Dated:

02/22/94

KOCIDE GX306

DRY FLOWABLE

FUNGICIDE/BACTERICIDE

Under the Federal Insecticide
Fungicide and Rodenticide Act
Registration No. 1812-358

Active Ingredient

Copper Hydroxide 53.8%

Inert Ingredients 42.2%

Total 100.0%

(Metallic Copper Equivalent 35%)

**KEEP OUT OF REACH OF CHILDREN
DANGER - PELIGRO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle
(If you do not understand the label, find someone to explain it to you in detail).

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Hold eyelids open and flush with water for 15 minutes. Get medical attention.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

IF SWALLOWED: Drink promptly a large quantity of milk, egg white, gelatin solution, or if these are not available, large quantities of water. Avoid alcohol. Get medical attention.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate use of gastric lavage.

Griffin Corporation
Valdosta, GA 31601

EPA Reg No 1812-358
EPA Est. No 8901-TX;1

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS (AND DOMESTIC ANIMALS)

DANGER - PELIGRO

Corrosive. Causes irreversible eye damage. Harmful if swallowed, absorbed through the skin or inhaled. May cause skin sensitization reactions in certain individuals. Avoid contact with skin, eyes, or clothing. Avoid breathing dust.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic organisms. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean water mark. Drift and runoff from treated areas may be hazardous to fish and aquatic organisms in adjacent aquatic sites. Do not allow rinsate from cleaning of equipment or disposed material to enter surface or ground water.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS:

Users Should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

5936

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours without required PPE.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides 40 CFR part 170. The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses. Non-crop weed control is not within the scope of the Worker Protection Standard.

STORAGE AND DISPOSAL

Store in a cool, dry place.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in sanitary landfill, or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

GENERAL INSTRUCTIONS

Use KOCIDE GX306 as noted below. KOCIDE GX306 may be applied as an aerial, ground dilute or ground concentrate spray unless specifically directed otherwise by specific crop instructions. Depending upon the equipment used and the specific crop, the volume applied per acre will differ. Refer to recommended volume table below.

This product may be reactive on metal and masonry surfaces such as galvanized roofing. Avoid contact with metal surfaces. Do not spray on cars, houses, lawn furniture, etc.

Minimum Recommended Spray Volume (Gallons) Per Acre When Applying Kocide GX306

	Aerial	Dilute	Concentrate	
Vegetables	3	20	---	✓
Field Crops	3	20	---	
Small Fruits	5	150	50	(Florida)
Vines	5	150	50	
Tree Crops	10	400	50	
Citrus	10	200	100	
Miscellaneous	10	150	50	
Turf (Algae control) 1.5 pounds in 5 gallons of water per 1,000 square feet.				

* Pesticide application equipment such as Curtec® or other similar sprayers which are capable of obtaining thorough coverage at low volumes may be used at as low as 20 gpa of spray volume

It must be determined in the selection process if proper application equipment is available and if the waste associated with its use can be properly handled. Materials used in the construction of application equipment is also an important factor as agricultural chemicals are often reactive with soft metals such as aluminum and even some synthetic materials such as plastics, rubbers, etc. Therefore it is necessary when working with equipment containing these materials that they are thoroughly flushed with clean water after each days use.

NOTE - Agricultural chemicals may perform in an unpredictable manner when tank mixed especially where several products are involved. Reduced effect on pests or crop injury may occur. Unless recommended on this label or by a state/local expert, or the user has small scale direct experience, tank mixing should not be undertaken.

NOTE: Do not tank mix Kocide GX306 with Aliette® fungicide unless appropriate precautions have been taken to buffer the spray solution or severe phytotoxicity may result.

GENERAL CHEMIGATION INSTRUCTIONS

Apply this product only through one or more of the following types of sprinkler irrigation systems: center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Posting of areas to be chemigated is required when 1) any part of a treated area is within 300 feet of sensitive areas such as residential areas, labor camps, businesses, day care centers, hospitals, in-patient clinics, nursing homes or any public areas such as schools, parks, playgrounds, or other public facilities not including public roads, or 2) when the chemigated area is open to the public such as golf courses or retail greenhouses.

Posting must conform to the following requirements. Treated areas shall be posted with signs at all usual points of entry and along likely routes of approach from the listed sensitive areas. When there are no usual points of entry, signs must be posted in the corners of the treated areas.

and in any location affording maximum visibility to sensitive areas. The printed side of the sign should face away from the treated area towards the sensitive area. The signs shall be printed in English. Signs must be posted prior to application and must remain until foliage has dried and soil surface water has disappeared. Signs may remain in place indefinitely as long as they are composed of materials to prevent deterioration and maintain legibility for the duration of the posting period.

All words shall consist of letters at least 2 1/2 inches tall, and all letters and the symbol shall be a color which sharply contrasts with their immediate background. At the top of the sign shall be the words KEEP OUT, followed by an octagonal stop sign symbol at least 8 inches in diameter containing the word STOP. Below the symbol shall be the words PESTICIDES IN IRRIGATION WATER.

This sign is in addition to any sign posted to comply with the Worker Protection Standard.

CHEMIGATION SYSTEMS CONNECTED TO PUBLIC WATER SYSTEMS

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into the reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

When mixing, fill nurse tank half full with water. Add KOCIDE GX306 slowly to tank while hydraulic or mechanical agitation is operating and continue filling with water. **DO NOT PRE-MIX OR SLURRY KOCIDE GX306.** Stickers, spreaders (cleared for use on growing crops), nutrients, etc. should be added last. If compatibility is in question, use the Compatibility Jar Test before mixing a whole tank. Because of the wide variety of possible combinations which can be encountered, observe all cautions and limitations on the label of all products used in mixtures.

KOCIDE GX306 should be added through a traveling irrigation system continuously or at the last 30 minutes of solid set or hand moved irrigation systems. Agitation of the mixture in the nurse tank is recommended.

SPRINKLER CHEMIGATION

) The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

) The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. ✓

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

When mixing, fill nurse tank half full with water. Add KOCIDE GX306 slowly to tank while hydraulic or mechanical agitation is operating and continue filling with water. **DO NOT PRE-MIX OR SLURRY Kocide GX 306** before adding to the nurse tank. Stickers, spreaders (cleared

10 36
Y

for use on growing crops), nutrients, etc. should be added last. If compatibility is in question, use the Compatibility Jar Test before mixing a whole tank. Because of the wide variety of possible combinations which can be encountered, observe all cautions and limitations on the label of all products used in mixtures.

KOCIDE GX306 should be added through a traveling irrigation system continuously or at the last 30 minutes of solid set or hand moved irrigation systems. Agitation of the mixture in the nurse tank is recommended.

Shut off injection equipment after treatment and continue to operate irrigation system until Kocide GX306 has been cleared from the last sprinkler head.

CROP CLASSIFICATION

CITRUS: Grapefruit, Kumquat, Lemon, Lime, Orange, Tangelo, and Tangerine.

FIELD CROPS: Alfalfa, Barley, Oats, Peanut, Potato, Sugar Beet, and Wheat.

SMALL FRUITS: Blackberry, Blueberry, Cranberry, Currant, Gooseberry, Raspberry and Strawberry.

TREE CROPS: Almond, Apple, Apricot, Avocado, Banana, Cacao, Cherry, Coffee, Filbert, Mango, Nectarine, Olive, Peach, Pear, Pecan, Pistachio, Plum, Prune, Quince, and Walnut

VEGETABLES: Bean, Broccoli, Brussels Sprouts, Cabbage, Cantaloupe, Carrot, Cauliflower, Celeriac, Celery, Cucumber, Eggplant, Endive, Escarole, Greens (Beet, Collard, Mustard and Turnip), Honeydew, Lettuce, Muskmelon, Onion, Pea, Pepper, Pumpkin, Spinach, Squash, Table Beet, Tomato, Watercress and Watermelon.

VINES: Grape, Hops, and Kiwi.

MISCELLANEOUS: Atemoya, Carambola, Chives, Dill, Douglas Fir, Ginseng, Guava, Litchi, Live Oak, Macadamia, Mamey Sapote, Papaya, Parsley, Passion Fruit, Sugar Apple and Sycamore.

GREENHOUSE AND SHADE HOUSE CROPS: While specific directions are presented for Citrus Cucumber, Eggplant, Pepper, and Tomato; general use may occur for any crop on this label where physiology allows Greenhouse or Shade house culture.

TURFGRASS (ALGAE CONTROL)

ORNAMENTALS: Species as listed

When selecting a use rate for KOCIDE GX306 do not apply less than the label recommended

...

10

minimum amount. Under heavy disease pressure or when conditions favor such, use the higher rate and shorter spray intervals specified for each crop. In addition, use the higher rates for large mature tree crops.

The per acre use rate of KOCIDE GX306 is applicable for both dilute and concentrate spraying. Consult the KOCIDE GX306 label for specific rates and timing of application by crop.

Complete spray coverage is essential to assure optimum performance from KOCIDE GX306. When treating on a concentrate basis or by aerial application, unless you have had specific previous experience, it is advisable to test for compatibility and tolerance to crop injury prior to full scale commercial utilization.

While volume is important in obtaining full spray coverage, often factors such as foliage density, environmental conditions and spray calibrations, have a greater impact. Always be sure that sprayers are calibrated to spray equipment manufacturer's specifications and environmental conditions are within those recommended by State and local regulatory authorities.

When mixing, fill spray tank one-half full with water. Add KOCIDE GX306 slowly to tank while hydraulic or mechanical agitation is operating and continue filling with water. DO NOT PRE-MIX OR SLURRY KOCIDE GX306. Spreaders, stickers (cleared for use on growing crops), nutrients, etc. should be added last. If compatibility is in question, use the Compatibility Jar Test before mixing a whole tank. Because of the wide variety of possible combinations which can be encountered, observe all cautions and limitations on the label of all products used in mixtures.

NOTE: KOCIDE GX306 should not be applied in a spray solution having a pH of less than 6.5 as phytotoxicity may occur.

Environmental conditions such as extended periods of wet weather, acid rain, etc. which alter the pH of the leaf surface may affect the performance of KOCIDE GX306 resulting in possible phytotoxicity or loss of effectiveness.

The following specific instructions are based on general application procedures. The recommendations of the State Agricultural Extension Service should be closely followed as to timing, frequency, and number of sprays per season.

FROST INJURY PROTECTION

BACTERIAL ICE NUCLEATION INHIBITOR

Application of KOCIDE GX306 made to all crops listed on this label at rates and stages of growth indicated on this label, at least 24 hours prior to anticipated frost conditions, will afford control of ice nucleating bacteria (*Pseudomonas syringae*, *Erwinia herbicola*, and *Pseudomonas*).

12936

fluorescens) and may therefore provide some protection against light frost. Not recommended for those geographical areas where weather conditions favor severe frost.

CITRUS

KOCIDE GX306 may be mixed with dry foliar nutritionals (micronutrients) to create "Shot Bag" mixes to meet the various nutritional requirements of citrus and provide disease protection as described on this label. KOCIDE GX306 per acre rates in these mixes must not exceed the maximum recommended labeled rates for disease control.

NOTE: Adding foliar nutritionals or other products to spray mixtures containing Kocide GX306 and applying to citrus during the post bloom period when young fruit is present may result in spray burn.

Disease	Rate/Acre	Use Instructions
Melanose, Scab, Algal Spot	3-9 lbs.	Apply as pre-bloom and post-bloom sprays. Use higher rates when conditions favor disease.
Greasy Spot, Pink Pitting	1.5-4.5 lbs.	Apply in summer on expanded new flush. Repeat on subsequent flushes where disease pressure is severe. Use higher rates when conditions favor disease.
Alternaria Brown Spot (suppression)	6-7.5 lbs	On susceptible varieties apply when the first spring flush appears and each flush thereafter. Application to the fruiting bodies should start after two thirds of the petals have fallen and be repeated on a 21-day schedule.
Phytophthora Brown Rot, Septoria Spot	3-6 lbs	Begin application in fall before or just after the first rain and continue as needed. Apply to entire tree. Apply also to bare ground one foot beyond skirt. Use higher rates when conditions favor disease.

NOTE: In California, in areas subject to copper injury, add 1/3 to 1 pound of high quality lime per pound of KOCIDE GX306

Handwritten notes and stamps at the bottom of the page, including a circular stamp with the number 10.

Phytophthora Foot Rot	1 lb.	Mix with 1 gallon of water and paint trunks of trees from the soil surface to the lowest scaffold limbs. Apply in May prior to summer rains and/or in the fall prior to wrapping trees for freeze protection. Treatment serves as protection for up to one year, but does not cure existing infections.
-----------------------	-------	---

Citrus Canker (Suppression Only)	9 lbs.	Spray flushes 7-14 days after shoots begin to grow. Young fruit may require an additional application. Number and timing of applications will be dependent upon disease pressure. Under heavy pressure, each flush of new growth should be sprayed.
-------------------------------------	--------	---

NOTE: Do not use KOCIDE GX306 on citrus seedlings grown in greenhouses or shade houses.

CITRUS Field Nursery Grown


To control melanose, scab, pink pitting, greasy spot, brown rot and for citrus canker (suppression), apply 1.5 pounds of Kocide GX306 per 100 gallons of water (3-6 lbs/acre). Apply Kocide GX306 at 28 day intervals or as needed depending on disease severity.

FIELD CROPS

Crop	Disease	Rate/Acre	Use Instructions
Alfalfa	Cercospora and Lep- tosphaerulina Leaf Spots	1.5 lbs.	Apply 10-14 days before each harvest or earlier if disease threatens

NOTE: Spray injury may occur with sensitive varieties such as Lathontan.

Peanut	Cercospora Leaf Spot	1.25-2.25 lbs.	One to two quarts of SUPER SIX® or equivalent flowable sulfur per acre may be added. Begin spraying at 35-40 days after planting or when disease symptoms first appear and repeat at 10-14 day intervals as needed. Reduce sprays to 7 day intervals during humid weather. Use higher rates when conditions favor disease
Potato	Early and Late Blight	0.75-3 lbs.	Apply 0.75 - 1.25 lbs. at 7-10 day intervals starting when plants are 6 inches high in locations where disease is light and up to 2.25-3 pounds per acre where disease is more severe. Under conditions of severe disease control with GX306 will be improved by tank mixing with other compatible fungicides registered for use on potatoes. Read and follow all label instructions of tank mix partners.
Sugar Beet	Cercospora Leaf Spot	1.5-3.75 lbs.	Begin applications when conditions first favor disease development and repeat at 10-14 day intervals as needed. Use the higher rate when disease is severe. Addition of a suitable agricultural spray oil is recommended
Wheat, Oats and Barley	Septoria Leaf Blotch, Helminthosporium Spot Blotch	1.25-1.5 lbs	Make first application at early heading and follow with second spray 10 days later. Use the higher rates when conditions favor disease.



Crop	Disease	Rate/Acre	Use Instructions
Blackberry (Santiams, Logans, Boysens, Marions, Auroras, Cascades, Chehalems & Thornless Evergreens)	Leaf and Cane Spot, Purple Blotch, Anthracnose, Yellow Rust, Pseudomonas Blight	3 lbs.	Make fall application after harvest. Apply delayed dormant spray after pruning/training in the spring. Add 1 quart of crop oil per acre.
	Leaf and Cane Spot, Purple Blotch, Anthracnose, Yellow Rust	1.5 lbs.	Apply when leaf buds begin to open and repeat when flower buds show white. Add 1 quart of crop oil per acre.
<p>NOTE: Crop injury may occur if applied to foliage under certain environmental conditions such as hot or prolonged moist periods. Discontinue applications if signs of crop injury appear.</p>			
Blueberry	Bacterial Canker	4.5 lbs.	Make first application before fall rains and a second application four weeks later.
Cranberry	Fruit Rot	6 lbs.	Make first application in late bloom. One or two additional applications at 10-14 day intervals may be required depending upon disease severity.
	Rose Bloom	6 lbs.	Apply three sprays on 10 - 14 day schedule as soon as symptoms are observed.
	Bacterial Stem Canker	6 lbs.	Apply post harvest and again in spring before bud burst. One or two additional applications at 10-14 day intervals may be required depending upon disease severity.
	Tip Blight (Monilinia), Stem and Leaf Blight, Red	6 lbs.	Apply delayed dormant spray in the Spring. Repeat at 10-14 day intervals as needed through pre-

	Leaf Spot			bloom.
Current, Gooseberry	Anthracnose, Leaf Spot	7.5 lbs.		Make initial application after first leaves have expanded. Continue on a 10 - 14 day schedule during wet conditions in the Spring. Make an additional application after harvest.
Raspberry	Leaf and Cane Spot, Purple Blotch, Anthracnose, Yellow Rust, Pseudomonas Blight	3 lbs.		Make fall application after harvest. Apply delayed dormant spray after training in the spring. Add 1 quart of crop oil per acre.
	Leaf and Cane Spot, Purple Blotch, Anthracnose, Yellow Rust	1.5 lbs.		Apply when leaf buds begin to open and repeat when flower buds show white. Add 1 quart of crop oil per acre.
				NOTE: Crop injury may occur if applied to foliage under certain environmental conditions such as hot or prolonged moist periods. Discontinue applications if signs of crop injury appear.
Strawberry	Leaf Spot and Leaf Blight	1.5-2.75 lbs		Begin application when plants are established and continue on a weekly schedule throughout season. Apply in at least 20 gallons of water. Use the higher rates when conditions favor disease.
				NOTE: Discontinue applications if signs of crop injury appear.

TREE CROPS

Crop	Disease		Rate/Acre	Use Instructions
Almond	Coryneum Blight		6-9 lbs	Make first application before fall
			14	

Apricot, Cherry,
Plum, Prune

(Shot Hole), Bacteria
Canker and Blast
(Pseudomona)

rains and a second at late dormant.
Use higher rates when rainfall is
heavy and disease pressure is high.
One pint of superior-type oil per
100 gallons of water may be
added.

For Cherries, where disease is
severe, an additional application at
leaf-fall may be required.

Almond only: For bacterial blast
control in sprinkler irrigated
orchards or where disease is
severe, apply 0.75 lbs. per acre
post-bloom, at 2 week intervals or
just before sprinkling.

NOTE: Injury may occur from
post-bloom sprays on almonds,
especially on Neplus varieties.

Coryneum Blight
(Shot Hole), Blossom
Brown Rot

4.5-6 lbs.
(almonds)
6-7.5 lbs.
(all others)

Early bloom (popcorn)
application: Apply before full
bloom. Use higher rates when
rainfall is heavy and disease
pressure is high.

NOTE: To avoid plant injury, do
not use above rate after full
bloom.

Apple

Anthracoise,
European Canker,
Blossom and Shoot
Blast (Pseudomonas)

9-12 lbs.

Apply before fall rains. Use
higher rates under severe disease
conditions.

NOTE: Use on yellow varieties
may cause discoloration. To avoid
discoloration, pick before
spraying.

	Fireblight	6-12 lbs.	Make application between silver-tip and green-tip. Apply as a full cover spray. NOTE: Crop injury may occur from late application; discontinue use when green tip reaches 1/2 inch.
	Crown or Collar Rot	3 lbs.	Mix in 100 gallons of water. Apply 4 gallons of suspension as a drench on the lower trunk area of each tree. Apply either in early spring or in late fall after harvest. NOTE: Do not use if soil pH is below 5.5 since copper toxicity may result.
Avocado	Anthrachnose, Blotch, Scab	6-9 lbs.	Apply when bloom buds begin to swell and continue application at monthly intervals for 5-6 applications. Use higher rate when conditions favor disease.
Banana	Sigatoka	1.5 lbs.	Apply by air in 10 gallons of water combining 0.5 gallons of agricultural oil. Apply on a 14 day schedule throughout the wet season. Apply at 21 day intervals during dry periods.
	Black Pitting	3 lbs	Mix in 100 gallons of water (4.0 pounds per acre) directly to the fruit stem and include the basal portion of the leaf crown. Apply during the first and second weeks after fruit emergence.

Cacao	Black Pod	1.5-6.5 lbs.	Begin applications at the start of the rainy season and continue while infection conditions persist. Apply 1.5 - 3.5 lbs. as often as 14 to 21 days in high rainfall areas at varying rates depending on disease severity. For drier areas, where 2 to 4 applications are recommended during critical infection periods and at long intervals, use 6.5 pounds per acre, according to disease incidence and planting density.
Coffee	Coffee Berry Disease (<u>Colletotrichum coffeanum</u>)	4.5-6 lbs.	Apply first spray after flowering and before onset of long rains and then at 21-28 day intervals until picking. Use higher rates when rainfall is heavy and disease pressure is high.
	Bacterial Blight (<u>Pseudomonas syringae</u>)	4.5-6 lbs.	Begin spray program before the onset of the long rains and continue throughout the rainy season at 14-21 day intervals. The critical time of spraying to control this disease is just before, during and after flowering(s) especially when coinciding with wet weather. Use higher rates when rainfall is heavy and disease pressure is high.
	Leaf Rust (<u>Hemileia vastatrix</u>)	1.5-3 lbs.	Apply before the onset of rain and then at 21 day intervals while the rains continue. Use higher rates when rainfall is heavy and disease pressure is high.
	Iron Spot (<u>Cercospora coffeicola</u>), Pink Disease (<u>Corticium salmonicolor</u>)	1.5 lbs.	Use concentrate or dilute spray. Begin treatment at the start of wet season and continue at monthly intervals for three applications.

Filbert	Bacterial Blight	12-18 lbs.	Apply as a post harvest spray. In seasons of heavy rainfall apply a second spray when three-fourths of the leaves have dropped. Add 1 pint of superior-type oil per 100 gallons of water. Use higher rates when rainfall is heavy and disease pressure is high.
	Eastern Filbert Blight	12-18 lbs.	Apply as a dilute spray in adequate water for thorough coverage. Make initial application after harvest in October before heavy winter rains begin. The next application should be made in late February to early March followed by another application one month later. If desired, add 1 pint of a sticking agent or superior type oil per 100 gallons of water. Use higher rates when rainfall is heavy and disease pressure is high.
Mango	Anthracnose	6 lbs	Apply monthly after fruit set until harvest. Use higher rates when rainfall is heavy and disease pressure is high.
Olive	Peacock Spot Olive Knot	6-9 lbs	Make first application before winter rains fall. A second application in early spring should be made if disease is severe. Apply the high rate for heavy disease pressure or when conditions favor disease development.
P e a c h , Nectarine	Leaf Curl, Coryneum Blight (Shot Hole), Bacterial Canker and Blast (<u>Pseudomonas</u>), Bacterial Spot (<u>Xanthomonas</u>)	6-12 lbs	Apply at leaf fall. Use the highest rate when rainfall is very heavy and disease pressure is high. May be used with agricultural spray oil.



	Blossom Brown Rot, Leaf Curl, Coryneum Blight, (Shot Hole)	6-9 lbs.	Full cover spray at pink bud.
	Bacterial Spot	0.75 lb.	Post-bloom application applied at first and second cover sprays.
			NOTE: Do not spray 3 weeks prior to harvest. Use only recommended rates. Spotting of leaves and defoliation may occur from use in cover sprays. Post- bloom application applied at first and second cover sprays.
) Pear	Fire Blight	0.75 lb.	Apply at 5 day intervals throughout the bloom period.
	Blossom Blast (<u>Pseudomonas</u>)	9-12 lbs.	Apply before fall rains and again during dormancy before spring growth starts. Use the higher rate when disease pressure is high or when conditions favor disease development.
			NOTE: Do not apply to D-Anjou pears. Excessive dosages may cause fruit russet.
) Pecan	Shuck & Kernel Rot (<u>Phytophthora cactorum</u>) and Zonate Leafspot (<u>Cristulariella pyramidalis</u>)	1 5-3 lbs.	For suppression, apply in sufficient water to ensure complete spray coverage at 2 to 4 week intervals starting at kernel growth and continuing until shucks open. Use the higher rate and shorter interval if frequent rainfall occurs.
Pistachio	Botrytis Blight, Botryosphaeria panicle and shoot blight, Septoria leaf blight, late blight (<u>Alternaria alternata</u>)	3-6 lbs.	Make initial application at bud swell and repeat on a 14-28 day schedule as dictated by disease conditions. If disease conditions are severe, use the high rate and short spray interval.

Quince	Fire Blight	0.75 lb.	Apply at 5 day intervals throughout the bloom period. Apply in adequate water for thorough coverage.
--------	-------------	----------	--

Walnut	Walnut Blight	6-9.5 lbs.	Apply first spray at early pre-bloom prior to or when catkins are partially expanded. Make additional applications during bloom and early nutlet stage or as needed if frequent rainfall occurs. Thorough coverage of catkins, leaves and nutlets is essential for effective control. When applied as a dilute spray, 1 pint of summer oil emulsion may be added per 100 gallons of spray.
--------	---------------	------------	--

NOTE: Adequate control may not be obtained when copper tolerant species of Xanthomonas bacteria are present.

VEGETABLES

Crop	Disease	Rate/Acre	Use Instructions
Bean (dry & green)	Brown Spot, Halo Blight, Common Blight	0.75-2.25 lbs	Use the higher rate for more severe disease. For protective sprays, make first application when plants are 6 inches high; repeat on a 7-14 day schedule depending upon environmental conditions.
Carrot	Alternaria and Cercospora Leaf Spot	1.5 lbs	Begin applications when disease first threatens and repeat at 7-14 day intervals as needed depending on disease severity.
Celery, Celeriac	Cercospora Early Blight, Septoria Late Blight, & Bacterial Blight	1.5 lbs	Begin applications as soon as plants are first established in the field, repeating at 5-7 day intervals depending on disease severity and

environmental conditions.

Crucifers
[Broccoli,
Brussels
Sprouts,
Cabbage,
Cauliflower,
Greens (Beet,
Collard,
Mustard and
Turnip)]

Black Rot
(Xanthomonas) &
Black Leaf Spot
(Alternaria) Downy
Mild ✓

0.75-1.5 lbs.

Apply at 7-10 day intervals.
Begin application after transplants
are set in the field, or shortly after
emergence of field seeded crops or
when conditions favor disease
development. Use higher rates
when conditions favor disease.

NOTE: Reddening of older
leaves may occur on Broccoli and
a flecking of wrapper leaves may
occur on Cabbage.

Cucurbits
(Cucumber,
Cantaloupe,
Honeydew,
Muskmelon,
Watermelon,
Pumpkin, and
Squash)

Alternaria Leaf Spot,
Angular Leaf Spot,
Anthracnose, Downy
and Powdery
Mildew, Gummy
Stem Blight, and
Watermelon Bacterial
Fruit Blotch
(suppression)

1.5 lbs.

Begin application when conditions
are favorable for disease develop-
ment and prior to development of
disease symptoms. Repeat at 5-7
day intervals. Use shorter
intervals and higher rates when
disease is present and conditions
are favorable for rapid expansion.

NOTE: Crop injury may occur
from application at higher rates
and shorter intervals. Discontinue
use if injury occurs.

Eggplant

Alternaria Blight,
Anthracnose, &
Phomopsis

1.5 lbs.

Begin applications prior to
development of disease symptoms.
Repeat sprays at 7-10 day
intervals or as needed depending
on disease severity.

Lettuce, Endive, Escarole	Downy Mildew	1.5 lbs.	Begin treatment when disease first appears and repeat every 7 - 10 days as needed to suppress disease.
			NOTE: Flecking and/or yellowing of leaves will occur under certain environmental conditions such as extended periods of moist weather, acid rains, or other conditions favoring reduced pH on leaf surfaces. Injury may be severe enough to reduce crop value.
Onion	Purple Blotch, Downy Mildew & Bacterial Blight	1.5 lbs. 0.75 - 1.25 lbs.	Begin when plants are 4-6 inches high and repeat at 7-10 day intervals as needed depending upon disease pressure. Can cause phytotoxicity to leaves.
Peas	Powdery Mildew	1.25 - 2.25 lbs.	Begin applications when disease symptoms first appear and repeat at weekly intervals as needed. Use higher rate for more severe disease.
Pepper	Bacterial Spot	1.5-2.25 lbs.	Begin applications when conditions first favor disease development and repeat at 5-10 day intervals as needed depending on disease severity. Use higher rates for severe disease.
Spinach	Anthracnose, White Rust, Blue Mold, Cercospora Leaf Spot	1.5-2.25 lbs	Begin application when disease first appears or when conditions favor disease development. Repeat at 7-10 day intervals as needed. Use higher rates when conditions favor disease.
			NOTE: Flecking may occur on Spinach leaves



Table Beet	Cercospora Leaf Spot	1.5-3.75 lbs.	Begin applications when conditions first favor disease development and repeat at 10-14 day intervals as needed. Use the higher rate when disease is severe. Addition of a suitable agricultural spray oil is recommended.
Tomato	Anthrachnose, Bacterial Speck, Bacterial Spot, Early Blight, Gray Leaf Mold, Late Blight, Septoria Leaf Spot	1.5-3 lbs.	Begin when disease first threatens and repeat at 7-10 day intervals or as needed depending on disease severity. Use higher rate for severe disease.
Watercress	Cercospora Leafspot	1.5 lbs.	Begin application when plants are first established in the field, repeating at 7 - 14 day intervals depending on disease severity and environmental conditions. Do not exceed 4 applications per crop. Apply using ground spray equipment at no less than 50 gallons of spray solution per acre. ✓

VINES

Crops	Disease	Rate/Acre	Use Instructions
Grape	Black Rot, Powdery Mildew, Downy Mildew	1.5 lbs.	Begin applications at late dormant up to bud break with subsequent applications throughout the season depending upon disease severity.

NOTE: Foliage injury may occur on copper sensitive varieties such as Concord, Delaware, Niagara, and Rosetts. Either test for sensitivity or add 1 to 3 pounds of hydrated lime per pound of KOCIDE GX306.

Hops	Downy Mildew	1.5 lbs.	Make crown treatment after pruning, but before training. After training, additional treatments are needed at about 10 day intervals.
------	--------------	----------	--

NOTE: Discontinue use 2 weeks before harvest.

Kiwi	<u>Pseudomonas syringae</u> , <u>Erwinia herbicola</u> , <u>Pseudomonas fluorescens</u>	6 lbs.	Apply in 200 gallons of water per acre. Make applications on a monthly basis. A maximum of 3 applications may be made.
------	---	--------	--

MISCELLANEOUS

Crop	Disease	Rate/Acre	Use Instructions
Atemoya	Anthracnose	2.25 lbs.	Make initial application just before flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage.
Carambola	Anthracnose	4.5 lbs.	Make initial application just before flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage.
Chives	Downy Mildew	1.5 lbs.	Begin applications when plants are established in the field. Repeat applications every 7-10 days as dictated by disease conditions. If disease pressure is high, use the shorter spray interval.
Dill	Phoma Leaf Spot and Rhizoctonia Foliage Blight	2 lbs.	Begin applications when plants are first established in the field and repeat at 7-10 day intervals depending upon disease severity and environmental conditions. If disease pressure is high, use the shorter spray interval and the

Douglas Fir	Rhabdoclone needlecast	1.5 lbs	higher rate. Begin applications at bud break and repeat at 3-4 week intervals. Apply in a tank mix with another registered pesticide if moderate to severe disease pressure is present.
Ginseng	Alternaria Leaf and Stem Blight	2 lbs.	Use as a tank mix with 2 pounds Rovral® 50W in 100 gallons of water. Begin KOCIDE-Rovral applications as soon as plants have emerged in spring. Applications should be repeated every 7 days until plants become dormant in fall. If schedule application is to be made before a rain shower, apply fungicides at least 8 hours before the rain, giving the fungicides time to dry on the plants. Use of a spreader-sticker or sticker is advised. NOTE: Alternaria Leaf and Stem Blight is most severe in humid conditions such as those found in the dense canopies of 2-4-year old Ginseng. It is very important that the stems be thoroughly covered with fungicide; therefore, use a spray apparatus which distributes the fungicide throughout the canopy.
Guava	Anthraxnose, Red Algae	2.25 lbs	Make initial application just before flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage.
Litchi	Anthraxnose	2.25 lbs	Make initial application just before flowering and repeat on a weekly schedule until just before