

DEC 31 1996

Angela Tessier  
Griffin Corporation  
P.O. Box 1847  
Valdosta, GA 31603

Dear Ms. Tessier:

Subject: Label Revision  
Kocide 2000  
EPA Registration No. 1812-358  
Your Submission Dated December 6, 1996

The amendment referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended is acceptable provided that you:

1. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling:

- Portions of the text were missing apparently due to holes punched in the master label. Assure that appropriate text appears on the printed labeling and for future submissions submit a complete copy of the label text.

2. Submit one (1) copy of your final printed labeling before you release the product for shipment.

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,



tu Philip V. Errico  
Acting Product Manager (22)  
Fungicide-Herbicide Branch  
Registration Division (7505C)

Enclosure

# Kocide® 2000

FUNGICIDE / BACTERICIDE

Made in the U.S.A.

2/13  
**DRY FLOWABLE**

**ACTIVE INGREDIENT**

Copper Hydroxide .....	53.8%
<b>INERT INGREDIENTS</b> .....	<del>46.2%</del>
<b>TOTAL</b> .....	100.0%

(Metallic Copper Equivalent 35%)

**KEEP OUT OF REACH OF CHILDREN**

**DANGER – PELIGRO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail).

**STATEMENT OF PRACTICAL TREATMENT**

**IF IN EYES:** Hold eyelids open and flush with water for 15 minutes. Get medical attention.

**IF ON SKIN:** Wash with plenty of soap and water. Get medical attention if irritation persists.

**IF SWALLOWED:** Drink promptly a large quantity of milk, egg white, gelatin solution, or if these are not available, large quantities of water. Do not induce vomiting or give anything by mouth to an unconscious person. Avoid alcohol. Get medical attention.

**IF INHALED:** Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate use of gastric lavage.

See Label for Additional Precautions and Directions for Use.

**GRIFFIN CORPORATION**  
VALDOSTA, GEORGIA 31601

**Specimen Label**

EPA REG. NO. 1812-358

3/13

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS (AND DOMESTIC ANIMALS)**  
**DANGER – PELIGRO**

Corrosive. Causes irreversible eye damage. Harmful if swallowed, absorbed through the skin or inhaled. May cause skin sensitization reactions in certain individuals. Avoid contact with skin, eyes or clothing. Avoid breathing dust.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**USER SAFETY RECOMMENDATIONS**

Users Should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

**ENVIRONMENTAL HAZARDS**

This pesticide is toxic to fish and aquatic organisms. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to fish and aquatic organisms in adjacent aquatic sites. Do not contaminate water by disposal of equipment washwaters.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours without required PPE.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

**STORAGE AND DISPOSAL**

Store in a cool, dry place.

**PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**INSTRUCTIONS**

Kocide 2000 may be applied as an aerial, ground dilute or ground concentrate spray unless specifically directed otherwise in the specific crop use directions.

The per acre use rate of Kocide 2000 is applicable for both dilute and concentrate spraying. Depending upon the equipment used and the specific crop, the spray volume applied per acre will differ. Refer to Recommended Minimum Spray Volume Table below. Complete spray coverage is essential to assure optimum performance from Kocide 2000. When treating by aerial application, or with low volume application equipment, unless you have had specific previous experience, it is advisable to test for compatibility and tolerance to crop injury prior to full scale commercial utilization.

While volume is important in obtaining full spray coverage, often factors such as foliage density, environmental conditions and sprayer calibrations, have a greater impact. Always be sure that sprayers are calibrated to spray equipment manufacturer's specifications and environmental conditions are within those recommended by state and local regulatory authorities.

Consult the Kocide 2000 label for specific rates and timing of application by crop. When selecting a Kocide 2000 use rate do not apply less than the label recommended minimum amount. Where application rates are provided in a range (6 to 12 pounds), the higher rates are recommended when rainfall is heavy and/or disease pressure high. Under heavy disease pressure or when conditions favor such, use the higher rate and shorter spray intervals specified for each crop. In addition, use the higher rates for large mature tree crops.

When mixing, fill spray tank 1/2 full with water. Add Kocide 2000 slowly to tank while hydraulic or mechanical agitation is operating and continue filling with water. Spreaders, stickers (cleared for application to growing crops), nutrients, etc. should be added last. If compatibility is in question, use the Compatibility Jar Test before mixing a whole tank.

**NOTE:** Kocide 2000 should not be applied in a spray solution having a pH of less than 6.5 as phytotoxicity may occur.

**NOTE:** Do not tank mix Kocide 2000 with Aliette® fungicide unless appropriate precautions have been taken to buffer the spray solution or severe phytotoxicity may result.

**NOTE:** This product may be reactive on masonry and metal surfaces such as galvanized roofing. Avoid contact with metal surfaces. Do not spray on cars, houses, lawn furniture, etc.

**NOTE:** Environmental conditions such as extended periods of wet weather, acid rain, etc. which alter the pH of the leaf surface may affect the performance of Kocide 2000 resulting in possible phytotoxicity or loss of effectiveness.

**NOTE:** Agricultural chemicals may perform in an unpredictable manner when tank mixed, especially where several products are involved. Reduced effect on pests or crop injury may occur. Unless recommended on this label or by a state/local expert, or the user has small scale direct experience, tank mixing should not be undertaken.

**NOTE:** It must be determined in the selection process if proper application equipment is available and if the waste associated with its use can be properly handled. Materials used in the construction of application equipment is also an important factor as agricultural chemicals are often reactive with soft metals such as aluminum and even some synthetic materials such as plastics, rubbers, etc. Therefore it is necessary when working with equipment containing these materials that they are thoroughly flushed with clean water after each days use.

## CROP CLASSIFICATION

**CITRUS:** Grapefruit, Kumquat, Lemon, Lime, Orange, Tangelo and Tangerine.

**FIELD CROPS:** Alfalfa, Barley, Oats, Peanut, Potato, Sugarbeet and Wheat.

**SMALL FRUITS:** Blackberry, Blueberry, Cranberry, Currant, Gooseberry, Raspberry and Strawberry.

**TREE CROPS:** Almond, Apple, Apricot, Avocado, Banana, Cacao, Cherry, Coffee, Filbert, Mango, Nectarine, Olive, Peach, Pear, Pecan, Pistachio, Plum, Prune, Quince and Walnut.

**VEGETABLES:** Bean, Beet Greens, Broccoli, Brussels Sprout, Cabbage, Cantaloupe, Carrot, Cauliflower, Celeriac, Celery, Cucumber, Eggplant, Greens (Collard, Mustard and Turnip), Honeydew, Muskmelon, Onion, Pea, Pepper, Pumpkin, Spinach, Squash, Table Beet, Tomato, Watercress and Watermelon.

**VINES:** Grape, Hops and Kiwi.

**GREENHOUSE AND SHADEHOUSE CROPS:** While specific directions are presented for Citrus, Cucumber, Eggplant, Pepper and Tomato; general use may occur for any crop on this label where physiology allows greenhouse or shadehouse culture.

**TURFGRASS:** Algae control

**MISCELLANEOUS:** Atemoya, Carambola, Chives, Dill, Douglas Fir, Ginseng, Guava, Litchi, Live Oak, Macadamia, Mamey Sapote, Papaya, Parsley, Passion Fruit, Pecan, Sugar Apple and Sycamore.

**ORNAMENTALS:** Species as listed.

### Minimum Recommended Spray Volume (Gallons) Per Acre When Applying Kocide 2000

	Aerial	Dilute	Concentrate
Vegetables	3	20	
Field Crops	3	20	
Small Fruits	5	150	50
Vines	5	150	50
Tree Crops	10	400	50
Citrus	10	800	100
			(Florida)*
Miscellaneous	10	150	50
Turf (Algae control) and Ornamentals			

\*Pesticide application equipment such as Curtec® or other similar sprayers which are capable of obtaining thorough coverage at low volumes may be used at as low as 20 gpa of spray volume.

The following specific instructions are based on general application procedures. The recommendations of the State Agricultural Extension Service should be closely followed as to timing, frequency and number of sprays per season.

## FROST INJURY PROTECTION BACTERIAL ICE NUCLEATION INHIBITOR

Application of Kocide 2000 made to all crops listed on this label at rates and stages of growth indicated on this label, at least 24 hours prior to anticipated frost conditions, will afford control of ice nucleating bacteria (*Pseudomonas syringae*, *Erwinia herbicola*, and *Pseudomonas fluorescens*) and may therefore provide some protection against light frost. Not recommended for those geographical areas where weather conditions favor severe frost.

## CITRUS

Kocide 2000 may be mixed with dry foliar nutritionals (micronutrients) to create "Shot Bag" mixes to meet the various nutritional requirements of citrus and provide disease protection as described on this label. Kocide 2000 per acre rates in these mixes must not exceed the maximum recommended labeled rates for disease control.

**NOTE:** Adding foliar nutritionals or other products to spray mixtures containing Kocide 2000 and applying to citrus during the post bloom period when young fruit is present may result in spray burn.

Disease	Rate / Acre	Use Instructions
Melanose, Scab, Algal Spot	3-9 lbs.	Apply as pre-bloom and post-bloom sprays. Use higher rates when conditions favor disease.
Greasy Spot, Pink Pitting	1.5-4.5 lbs.	Apply in summer on expanded new flush. Repeat on subsequent flushes where disease pressure is severe. Use higher rates when conditions favor disease.
Alternaria Brown Spot (Suppression)	6-7.5 lbs	On susceptible varieties apply when the first spring flush appears and each flush thereafter. Application to the fruit should start after two thirds of the petals have fallen and be repeated on a 21 day schedule.

CITRUS *Cont'd.*

5/13

Disease	Rate / Acre	Use Instructions
Phytophthora Brown Rot, Septoria	3-6 lbs.	Begin application in fall before or just after the first rain and continue as needed. Apply to entire tree. Apply also to bare ground 1 foot beyond skirt. Use higher rates when conditions favor disease. NOTE: In California, in areas subject to copper injury, add 1/3 to 1 pound of high quality lime per pound of Kocide 2000.
Phytophthora Foot Rot	1 lb.	Mix with 1 gallon of water and paint trunks of trees from the soil surface to the lowest scaffold limbs. Apply in May prior to summer rains and/or in the fall prior to wrapping trees for freeze protection. Treatment serves as protection for up to 1 year, but does not cure existing infections.
Citrus Canker (Suppression Only)	9 lbs.	Spray flushes 7 to 14 days after shoots begin to grow. Young fruit may require an additional application. Number and timing of applications will be dependent upon disease pressure. Under heavy pressure, each flush of new growth should be sprayed.

NOTE: Do not use Kocide 2000 on citrus seedlings grown in greenhouses or shadehouses.

### CITRUS Field Nursery Grown

To control melanose, scab, pink pitting, greasy spot, brown rot and for citrus canker (Suppression), apply 1.5 pounds of Kocide 2000 per 100 gallons of water (3 to 6 pounds/acre). Apply Kocide 2000 at 28 day intervals or as needed depending on disease severity.

### FIELD CROPS

Crop	Disease	Rate / Acre	Use Instructions
Alfalfa	Cercospora and Leptosphaerulina Leaf Spots	1.5 lbs.	Apply 10 to 14 days before each harvest or earlier if disease threatens. NOTE: Spray injury may occur with sensitive varieties such as Lathontan.
Peanut	Cercospora Leaf Spot	1.25-2.25 lbs.	One to two quarts of flowable sulfur per acre may be added. Begin spraying at 35 to 40 days after planting or when disease symptoms first appear and repeat at 10 to 14 day intervals as needed. Reduce sprays to 7 day intervals during humid weather. Use higher rates when conditions favor disease.
Potato	Early and Late Blight	0.75-3 lbs.	Apply 0.75 to 1.25 pounds at 7 to 10 day intervals starting when plants are 6 inches high in locations where disease is light and up to 2.25 to 3 pounds per acre where disease is more severe. Under conditions of severe disease, control with Kocide 2000 will be improved by tank mixing with other compatible fungicides registered for use on potatoes. Read and follow all label instructions of tank mix partners.
Sugarbeet	Cercospora Leaf Spot	1.5-3.75 lbs.	Begin applications when conditions first favor disease development and repeat at 10 to 14 day intervals as needed. Use the higher rate when disease is severe. Addition of a suitable agricultural spray oil is recommended.
Wheat, Oats and Barley	Septoria Leaf Blotch, Helminthosporium Spot Blotch	1.25-1.5 lbs.	Make first application at early heading and follow with second spray 10 days later. Use the higher rates when conditions favor disease.

### SMALL FRUITS

Crop	Disease	Rate / Acre	Use Instructions
Blackberry (Santiams, Logans, Boysens, Marions, Auroras, Cascades, Chehalms & Thornless Evergreens)	Leaf and Cane Spot, Purple Blotch, Anthracnose, Yellow Rust, Pseudomonas Blight	3 lbs.	Make fall application after harvest. Apply delayed dormant spray after pruning/training in the spring. Add 1 quart of crop oil per acre.
	Leaf and Cane Spot, Purple Blotch, Anthracnose, Yellow Rust	1.5 lbs.	Apply when leaf buds begin to open and repeat when flower buds show white. Add 1 quart of crop oil per acre.  NOTE: Crop injury may occur if applied to foliage under certain environmental conditions such as hot or prolonged moist periods. Discontinue applications if signs of crop injury appear.
Blueberry	Bacterial Canker	4.5 lbs.	Make first application before fall rains and a second application 4 weeks later.
Raspberry	Fruit Rot	6 lbs.	Make first application in late bloom. One or two additional applications at 10 to 14 day intervals may be required depending upon disease severity.
	Rose Bloom	6 lbs.	Apply three sprays on 10 to 14 day schedule as soon as symptoms are observed.

# SMALL FRUITS *Cont'd.*

6/13

Crop	Disease	Rate / Acre	Use Instructions
Cranberry <i>Cont'd.</i>	Bacterial Stem Canker	6 lbs.	Apply postharvest and again in spring before bud burst. One or two additional applications at 10 to 14 day intervals may be required depending upon disease severity.
	Tip Blight (Monilinia), Stem and Leaf Blight, Red Leaf Spot	6 lbs.	Apply delayed dormant spray in the spring. Repeat at 10 to 14 day intervals as needed through pre-bloom.
Currant, Gooseberry	Anthrachnose, Leaf Spot	7.5 lbs.	Make initial application after first leaves have expanded. Continue on a 10 to 14 day schedule during wet conditions in the spring. Make an additional application after harvest.
Raspberry	Leaf and Cane Spot, Purple Blotch, Anthracnose, Yellow Rust, Pseudomonas Blight	3 lbs.	Make fall application after harvest. Apply delayed dormant spray after training in the spring. Add 1 quart of crop oil per acre.
	Leaf and Cane Spot, Purple Blotch, Anthracnose, Yellow Rust	1.5 lbs.	Apply when leaf buds begin to open and repeat when flower buds show white. Add 1 quart of crop oil per acre.  <b>NOTE:</b> Crop injury may occur if applied to foliage under certain environmental conditions such as hot or prolonged moist periods. Discontinue applications if signs of crop injury appear.
Strawberry	Leaf Spot and Leaf Blight	1.5-2.75 lbs.	Begin application when plants are established and continue on a weekly schedule throughout season. Apply in at least 20 gallons of water. Use the higher rates when conditions favor disease.  <b>NOTE:</b> Discontinue applications if signs of crop injury appear.

## TREE CROPS

Crop	Disease	Rate / Acre	Use Instructions
Almond, Apricot, Cherry, Plum, Prune	Coryneum Blight (Shot Hole), Bacterial Canker and Blast ( <i>Pseudomona</i> )	6-9 lbs.	Make first application before fall rains and a second at late dormant. Use higher rates when rainfall is heavy and disease pressure is high. One pint of superior-type oil per 100 gallons of water may be added.  For cherries, where disease is severe, an additional application at leaf fall may be required.  Almond only: For bacterial blast control in sprinkler irrigated orchards or where disease is severe, apply 0.75 pound per acre post-bloom, at 2 week intervals or just before sprinkling.  <b>NOTE:</b> Injury may occur from post-bloom sprays on almonds, especially on Neplus varieties.
	Coryneum Blight (Shot Hole), Blossom Brown Rot	4.5-6 lbs. (almonds) 6-7.5 lbs. (all others)	Early bloom (popcorn) application: Apply before full bloom. Use higher rates when rainfall is heavy and disease pressure is high.  <b>NOTE:</b> To avoid plant injury, do not use above rate after full bloom.
Apple	Anthrachnose, European Canker, Blossom and Shoot Blast ( <i>Pseudomonas</i> )	9-12 lbs.	Apply before fall rains. Use higher rates under severe disease conditions.  <b>NOTE:</b> Use on yellow varieties may cause discoloration. To avoid discoloration, pick before spraying.
	Fireblight	6-12 lbs.	Make application between silver-tip and green-tip. Apply as a full-cover spray.  <b>NOTE:</b> Crop injury may occur from late application; discontinue use when green tip reaches 1/2 inch.
	Crown or Collar Rot	3 lbs.	Mix in 100 gallons of water. Apply 4 gallons of suspension as a drench on the lower trunk area of each tree. Apply either in early spring or in late fall after harvest.  <b>NOTE:</b> Do not use if soil pH is below 5.5 since copper toxicity may result.
Avocado	Anthrachnose, Blotch, Scab	6-9 lbs.	Apply when bloom buds begin to swell and continue application at monthly intervals for five to six applications. Use higher rate when conditions favor disease.
Banana	Sigatoka	1.5 lbs.	Apply by air in 10 gallons of water combining 0.5 gallon of agricultural oil. Apply on a 14 day schedule throughout the wet season. Apply at 21 day intervals during dry periods.
	Black Pitting	3 lbs.	Mix in 100 gallons of water (3.4 pounds per acre) directly to the fruit stem and include the basal portion of the leaf crown. Apply during the first and second weeks after fruit emergence.

**TREE CROPS *Contra*.**

7/13

Crop	Disease	Rate / Acre	Use Instructions
Cacao	Black Pod	1.5-6.5 lbs.	Begin applications at the start of the rainy season and continue while infection conditions persist. Apply 1.5 to 3.5 pounds as often as 14 to 21 days in high rainfall areas at varying rates depending on disease severity. For drier areas, where two to four applications are recommended during critical infection periods and at long intervals, use 6.5 pounds per acre, according to disease incidence and planting density.
Coffee	Coffee Berry Disease ( <i>Colletotrichum coffeanum</i> )	4.5-6 lbs.	Apply first spray after flowering and before onset of long rains and then at 21 to 28 day intervals until picking. Use higher rates when rainfall is heavy and disease pressure is high.
	Bacterial Blight ( <i>Pseudomonas syringae</i> )	4.5-6 lbs.	Begin spray program before the onset of the long rains and continue throughout the rainy season at 14 to 21 day intervals. The critical time of spraying to control this disease is just before, during and after flowering(s) especially when coinciding with wet weather. Use higher rates when rainfall is heavy and disease pressure is high.
	Leaf Rust ( <i>Hemileia vastatrix</i> )	1.5-3 lbs.	Apply before the onset of rain and then at 21 day intervals while the rains continue. Use higher rates when rainfall is heavy and disease pressure is high.
	Iron Spot ( <i>Cercospora coffeicola</i> ), Pink Disease ( <i>Corticium salmonicolor</i> )	1.5 lbs.	Use concentrate or dilute spray. Begin treatment at the start of wet season and continue at monthly intervals for three applications.
Filbert	Bacterial Blight	12-18 lbs.	Apply as a postharvest spray. In seasons of heavy rainfall apply a second spray when three fourths of the leaves have dropped. Add 1 pint of superior-type oil per 100 gallons of water. Use higher rates when rainfall is heavy and disease pressure is high.
	Eastern Filbert Blight	12-18 lbs.	Apply as a dilute spray in adequate water for thorough coverage. Make initial application after harvest in October before heavy winter rains begin. The next application should be made in late February to early March followed by another application 1 month later. If desired, add 1 pint of a sticking agent or superior type oil per 100 gallons of water. Use higher rates when rainfall is heavy and disease pressure is high.
Mango	Anthrachnose	6 lbs.	Apply monthly after fruit set until harvest.
	Peacock Spot, Olive Knot	6-9 lbs.	Make first application before winter rains fall. A second application in early spring should be made if disease is severe. Apply the high rate for heavy disease pressure or when conditions favor disease development.
Peach, Nectarine	Leaf Curl, Coryneum Blight (Shot Hole), Bacterial Canker and Blast ( <i>Pseudomonas</i> ), Bacterial Spot ( <i>Xanthomonas</i> )	6-12 lbs.	Apply at leaf fall. Use the highest rate when rainfall is very heavy and disease pressure is high. May be used with agricultural spray oil.
	Blossom Brown Rot, Leaf Curl, Coryneum Blight (Shot Hole)	6-9 lbs.	Full cover spray at pink bud.
	Bacterial Spot	0.75 lb.	Post-bloom application applied at first and second cover sprays.  NOTE: Do not spray 3 weeks prior to harvest. Use only recommended rates. Spotting of leaves and defoliation may occur from use in cover sprays. Post-bloom application applied at first and second cover sprays.
Pear	Fire Blight	0.75 lb.	Apply at 5 day intervals throughout the bloom period. NOTE: Do not apply to D-Anjou pears. Excessive dosages may cause fruit russet.
	Blossom Blast ( <i>Pseudomonas</i> )	9-12 lbs.	Apply before fall rains and again during dormancy before spring growth starts. Use the higher rate when disease pressure is high or when conditions favor disease development.
Pecan	Shuck & Kernel Rot ( <i>Phytophthora cactorum</i> ) and Zonate Leafspot ( <i>Cristulariella pyramidalis</i> )	1.5-3 lbs.	For suppression, apply in sufficient water to ensure complete spray coverage at 2 to 4 week intervals starting at kernel growth and continuing until shucks open. Use the higher rate and shorter interval if frequent rainfall occurs.
Pistachio	Botrytis Blight, Botryosphaeria Panicle and Shoot Blight, Septoria Leaf Blight, Late Blight ( <i>Alternaria alternata</i> )	3-6 lbs.	Make initial application at bud swell and repeat on a 14 to 28 day schedule as dictated by disease conditions. If disease conditions are severe, use the high rate and short spray interval.

Crop	Disease	Rate / Acre	Use Instructions
Quince	Fire Blight	0.75 lb.	Apply at 5 day intervals throughout the bloom period. Apply in adequate water for thorough coverage.
Walnut	Walnut Blight	6-9.5 lbs.	Apply first spray at early pre-bloom prior to or when catkins are partially expanded. Make additional applications during bloom and early nutlet stage or as needed if frequent rainfall occurs. Thorough coverage of catkins, leaves and nutlets is essential for effective control. When applied as a dilute spray, 1 pint of summer oil emulsion may be added per 100 gallons of spray.

NOTE: Adequate control may not be obtained when copper tolerant species of *Xanthomonas* bacteria are present.

## VEGETABLES

Crop	Disease	Rate / Acre	Use Instructions
Bean (dry & green)	Brown Spot, Halo Blight, Common Blight	0.75-2.25 lbs.	Use the higher rate for more severe disease. For protective sprays, make first application when plants are 6 inches high; repeat on a 7 to 14 day schedule depending upon environmental conditions.
Carrot	Alternaria and Cercospora Leaf Spot	1.5 lbs.	Begin applications when disease first threatens and repeat at 7 to 14 day intervals as needed depending on disease severity.
Celery, Celeriac	Cercospora Early Blight, Septoria Late Blight, Bacterial Blight	1.5 lbs.	Begin applications as soon as plants are first established in the field, repeating at 5 to 7 day intervals depending on disease severity and environmental conditions.
Crucifers [Broccoli, Brussels Sprout, Cabbage, Cauliflower, Greens (Collard, Mustard and Turnip)]	Black Rot ( <i>Xanthomonas</i> ), Black Leaf Spot ( <i>Alternaria</i> ), Downy Mildew	0.75-1.5 lbs.	Apply at 7 to 10 day intervals. Begin application after transplants are set in the field, or shortly after emergence of field seeded crops or when conditions favor disease development. Use higher rates when conditions favor disease.
			NOTE: Reddening of older leaves may occur on broccoli and a flecking of wrapper leaves may occur on cabbage.
Cucurbits (Cantaloupe, Cucumber, Honeydew, Muskmelon, Pumpkin Squash and Watermelon)	Alternaria Leaf Spot, Angular Leaf Spot, Anthracnose, Downy and Powdery Mildew, Gummy Stem Blight, and Watermelon Bacterial Fruit Blotch (Suppression)	1.5 lbs.	Begin application when conditions are favorable for disease development and prior to development of disease symptoms. Repeat at 5 to 7 day intervals. Use shorter intervals when disease is present and conditions are favorable for rapid expansion.
			NOTE: Crop injury may occur from application at shorter intervals. Discontinue use if injury occurs.
Eggplant	Alternaria Blight, Anthracnose, Phomopsis	1.5 lbs.	Begin applications prior to development of disease symptoms. Repeat sprays at 7 to 10 day intervals or as needed depending on disease severity.
Onion	Purple Blotch, Downy Mildew	1.5 lbs.	Begin when plants are 4 to 6 inches high and repeat at 7 to 10 day intervals as needed depending upon disease pressure.
	Bacterial Blight	0.75-1.25 lbs.	Can cause phytotoxicity to leaves.
Peas	Powdery Mildew	1.25-2.25 lbs.	Begin applications when disease symptoms first appear and repeat at weekly intervals as needed. Use higher rate for more severe disease.
Pepper	Bacterial Spot	1.5-2.25 lbs.	Begin applications when conditions first favor disease development and repeat at 5 to 10 day intervals as needed depending on disease severity. Use higher rates for severe disease.
Spinach	Anthracnose, White Rust, Blue Mold, Cercospora Leaf Spot	1.5-2.25 lbs.	Begin application when disease first appears or when conditions favor disease development. Repeat at 7 to 10 day intervals as needed. Use higher rates when conditions favor disease.
			NOTE: Flecking may occur on spinach leaves.
Table Beet	Cercospora Leaf Spot	1.5-3.75 lbs.	Begin applications when conditions first favor disease development and repeat at 10 to 14 day intervals as needed. Use the higher rate when disease is severe. Addition of a suitable agricultural spray oil is recommended.
Tomato	Anthracnose, Bacterial Speck, Bacterial Spot, Early Blight, Gray Leaf Mold, Late Blight, Septoria Leaf Spot	1.5-3 lbs.	Begin when disease first threatens and repeat at 7 to 10 day intervals or as needed depending on disease severity. Use higher rate for severe disease.
Watercress	Cercospora Leafspot	1.5 lbs.	Begin application when plants are first established in the field, repeating at 7 to 14 day intervals depending on disease severity and environmental conditions. Do not exceed four applications per crop. Apply using ground spray equipment at no less than 50 gallons of spray solution per acre.



## VINES

9/13

Crop	Disease	Rate / Acre	Use Instructions
Grape	Black Rot, Powdery Mildew, Downy Mildew	1.5 lbs.	Begin applications at late dormant up to bud break with subsequent applications throughout the season depending upon disease severity.  NOTE: Foliage injury may occur on copper sensitive varieties such as Concord, Delaware, Niagara and Rosetts. Either test for sensitivity or add 1 to 3 pounds of hydrated lime per pound of Kocide 2000.
Hops	Downy Mildew	1.5 lbs.	Make crown treatment after pruning, but before training. After training, additional treatments are needed at about 10 day intervals.  NOTE: Discontinue use 2 weeks before harvest.
Kiwi	<u>Pseudomonas syringae</u> , <u>Erwinia herbicola</u> , <u>Pseudomonas fluorescens</u>	6 lbs.	Apply in 200 gallons of water per acre. Make applications on a monthly basis. A maximum of three applications may be made.

## MISCELLANEOUS

Crop	Disease	Rate / Acre	Use Instructions
temoya	Anthrachnose	2.25 lbs.	Make initial application just before flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage.
Carambola	Anthrachnose	4.5 lbs.	Make initial application just before flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage.
Chives	Downy Mildew	1.5 lbs	Begin applications when plants are established in the field. Repeat applications every 7 to 10 days as dictated by disease conditions. If disease pressure is high, use the shorter spray interval.
1	Phoma Leaf Spot and Rhizoctonia Foliage Blight	2 lbs.	Begin applications when plants are first established in the field and repeat at 7 to 10 day intervals depending upon disease severity and environmental conditions. If disease pressure is high, use the shorter spray interval and the higher rate.
Douglas Fir	Rhabdocline Needlecast	1.5 lbs	Begin applications at bud break and repeat at 3 to 4 week intervals. Apply in a tank mix with another registered pesticide if moderate to severe disease pressure is present.
Ginseng	Alternaria Leaf and Stem Blight	2 lbs.	Use as a tank mix with 2 pounds Rovral® 50W in 100 gallons of water. Begin Kocide-Rovral applications as soon as plants have emerged in spring.  Applications should be repeated every 7 days until plants become dormant in fall. If schedule application is to be made before a rain shower, apply fungicides at least 8 hours before the rain, giving the fungicides time to dry on the plants. Use of a spreader-sticker or sticker is advised.  NOTE: Alternaria leaf and stem blight is most severe in humid conditions such as those found in the dense canopies of 2 to 4 year old ginseng. It is very important that the stems be thoroughly covered with fungicide; therefore, use a spray apparatus which distributes the fungicide throughout the canopy.
Guava	Anthrachnose, Red Algae	2.25 lbs.	Make initial application just before flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage.
Litchi	Anthrachnose	2.25 lbs.	Make initial application just before flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage.
Pecan, Live Oaks	Ball Moss	6 lbs.	Apply 6 pounds per 100 gallons of water, in the spring when ball moss is actively growing, using 1.5 gallons of spray per foot of tree height. Make sure to wet ball moss tufts thoroughly. A second application may be required after 12 months.  NOTE: Kocide 2000 may be injurious to ornamentals grown under Live Oaks. This product may be reactive on metal and masonry surfaces such as galvanized roofing. Avoid contact with metal surfaces. Do not spray on cars, houses, lawn furniture, etc.

Crop	Disease	Rate / Acre	Use Instructions
Macadamia	Anthraco	4.5 lbs.	Initiate sprays at first sign of flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage.
	Phytophthora Blight ( <i>P. capsici</i> ), Raceme Blight ( <i>Botrytis cinerea</i> )	3.5-4.5 lbs.	Apply during raceme development and bloom periods. Apply in sufficient water for thorough coverage. Use higher rates when conditions favor disease.
Mamey Sapote	Anthraco, Algal Leaf Spot	4.5-6 lbs.	Apply when conditions favor disease development. Repeat on 14 to 30 day schedule as disease severity and environmental conditions dictate. Use higher rates when conditions favor disease.
Papaya	Anthraco	3-7.5 lbs.	Apply before disease appears. Apply at 10 to 14 day intervals under light disease pressure and at 5 to 7 day intervals under heavy disease pressure. The addition of an approved spreader is desirable. Use higher rates when conditions favor disease.
Parsley	Bacterial Blight ( <i>Pseudomonas</i> sp.)	2.25 lbs.	Begin applications when plants are first established in the field and repeat at 5 to 7 day intervals depending upon disease severity and environmental conditions.
Passion Fruit	Anthraco	4.5 lbs.	Make initial application just before flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage.
Sugar Apple ( <i>Annona</i> )	Anthraco	9 lbs.	Make initial application just before flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage.
Sycamore	Anthraco	1.5-2.25 lbs.	Apply as a full cover spray. Apply in 100 gallons of water or sufficient volume for thorough coverage. Make first application at bud crack and second application 7 to 10 days later at 10% leaf expansion. Use higher rates when conditions favor disease.

## TURFGRASS

To control algae in turfgrass, apply 1 pound Kocide 2000 per 1,000 square feet in 5 gallons of water. Kocide 2000 may be used alone or in combination with other registered fungicides as a maintenance spray. Observe all precautions and limitations on the label of each product used in tank mixes.

**NOTE:** Phytotoxicity may occur depending upon varietal differences. Apply the recommended rate to a small area and observe for 7 to 10 days for signs of injury. If phytotoxicity occurs, discontinue use. Do not apply in spray solutions with a pH of less than 6.5.

## GREENHOUSE AND SHADEHOUSE CROPS

**Notice to User:** Kocide 2000 may be used in greenhouses and shadehouses to control diseases on some crops which appear on this label; specific instructions have been included for certain crops. The grower should bear in mind that the sensitivity of crops grown in greenhouses and shadehouses differ greatly from crops grown under field conditions. Neither the manufacturer nor seller has determined whether or not Kocide 2000 can be used safely on all greenhouse and shadehouse grown crops. The user should determine if Kocide 2000 can be used safely prior to commercial use. In a small area, apply the recommended rates to the plants in question, i.e. foliage, fruit, etc., and observe for 7 to 10 days for symptoms of phytotoxicity prior to commercial use.

Apply Kocide 2000 according to specific rates given for those crops in pounds per acre or pounds per 100 gallons. **One level tablespoon of Kocide 2000 per 1000 square feet is equivalent to 1 pound per acre.** Kocide 2000 should be applied in adequate water for thorough coverage of plant parts. Begin application at first sign of disease and repeat at 7 to 14 day intervals as needed; use shorter interval during periods when severe disease conditions persist.

**NOTE:** Do not use Kocide 2000 on citrus seedlings grown in greenhouses or shadehouses.

Crop	Disease	Rate Per 1000 Sq Ft	Use Instructions
Cucumber	Angular Leaf Spot and Downy Mildew	1.25-1.5 TBSP	Apply weekly when plants begin to vine. Use higher rates when conditions favor disease.
Eggplant	Alternaria Blight, Anthracnose, & Phomopsis	1.5 TBSP	Begin applications prior to development of disease symptoms. Repeat sprays at 7 to 10 day intervals or as disease pressure dictates.
Pepper	Bacterial Spot	1.5-2.25 TBSP	Begin applications when conditions first favor disease development and repeat at 5 to 10 day intervals as needed depending on disease severity. Use higher rates for severe disease.
Tomato	Anthraco, Bacterial Speck, Bacterial Spot, Early Blight, Grey Leaf Mold, Late Blight, Septoria Leaf Spot	1.5-2.25 TBSP	Begin when disease first threatens and repeat at 7 to 10 day intervals or as needed depending on disease severity. Use higher rate for severe disease.

# ORNAMENTALS

11/13

**Notice to User:** Plant sensitivities to Kocide 2000 have been found to be acceptable in specific genera and species listed on this label, however, phytotoxicity may occur. Due to the large number of species, widely varying growth conditions, and varieties of ornamentals and nursery plants, it is impossible to test every one for sensitivity to Kocide 2000. Neither the manufacturer nor seller has determined whether or not Kocide 2000 can be safely used on ornamental or nursery plants not listed on this label. The user should determine if Kocide 2000 can be used safely prior to commercial use. In a small area, apply the recommended rates to the plants in question, i.e. bedding plants, foliage, etc., and observe for 10 days for symptoms of phytotoxicity prior to commercial use.

Use Kocide 2000 on container, bench or bed-grown ornamentals in greenhouses or outdoor nurseries, for professional use on ornamentals grown for indoor and outdoor landscaping, and for control of bacterial and fungal diseases of foliage, flowers and stems.

One half tablespoon of Kocide 2000 per gallon of water is equivalent to 1 pound per 100 gallons.

Apply as a thorough coverage spray using 0.75 pound Kocide 2000 per 100 gallons of water. Begin application at first sign of disease and repeat at 7 to 14 day intervals as needed; use shorter interval during periods of frequent rains or when severe disease conditions persist.

Kocide 2000 may be used alone or in combination with other fungicides as a maintenance spray. Observe all precautions and limitations on the label of each product used in tank mixes.

**NOTE:** Do not tank mix Kocide 2000 with Aliette® fungicide unless appropriate precautions have been taken to buffer the spray solution. Severe phytotoxicity may result if adequate precautions are not taken.

Crop	Latin Name	Disease
Althea (Rose of Sharon)	<u>Hibiscus syriacus</u>	Bacterial Leaf Spot
Aralia	<u>Dizygotheca elegantissima</u>	Xanthomonas & Cercospora Leaf Spots, Alternaria
Arborvitae	<u>Thuja</u> sp.	Alternaria Twig Blight, Cercospora Leaf Blight
Azalea	<u>Rhododendron</u> sp.	Cercospora Leaf Spot, Botrytis Blight, Phytophthora Dieback, Powdery Mildew
Begonia	<u>Begonia semperflorens</u>	Bacterial Leaf Spot ( <u>Xanthomonas</u> sp., <u>Erwinia</u> sp., <u>Pseudomonas</u> sp.)
Bougainvillea	<u>Bougainvillea spectabilis</u>	Anthrachnose, Bacterial Leaf Spot
Bulbs (Tulip, Gladiolus)	Miscellaneous	Anthrachnose, Botrytis Bligh
Camphor Tree	<u>Cinnamomum camphora</u>	Pseudomonas Leaf Spot
Carnation	<u>Dianthus</u> sp.	Alternaria Blight, Pseudomonas Leaf Spot, Botrytis Blight
Camelia	<u>Camellia japonica</u> , <u>C. sasangua</u>	Anthrachnose, Bacterial Leaf Spot
Canna	<u>Canna</u> sp.	Pseudomonas Leaf Spot
Chinese Tallow Tree	<u>Sapium sebiferum</u>	Bacterial Leaf Spot ( <u>Xanthomonas</u> sp., <u>Pseudomonas</u> sp.)
Chrysanthemum	<u>Chrysanthemum morifolium</u>	Septoria Leaf Spot, Botrytis Blight
Cotoneaster	<u>Cotoneaster</u> sp.	Botrytis Blight
Dahlia	<u>Dahlia pinnata</u>	Alternaria Leaf Spot, Botrytis Gray Mold, Cercospora Leaf Spot
Date Palm	<u>Phoenix canariensis</u>	Pestalotia Leaf Spot
Dianthus	<u>Dianthus</u> sp.	Bacterial Spot, Bacterial Soft Rot
Dogwood	<u>Cornus florida</u>	Anthrachnose
Dusty Miller	<u>Senecio cineraria</u>	Bacterial Leaf Spot ( <u>Pseudomonas cichorii</u> )
Easter Lily	<u>Lilium longiflorum</u>	Botrytis Blight
Echinacea	<u>Echinacea</u> sp.	Bacterial Leaf Spot ( <u>Pseudomonas cichorii</u> )
Elm "Drake"	<u>Ulmus parvifolia</u>	Xanthomonas Leaf Spot
Euonymus	<u>Euonymus</u> sp.	Botrytis Blight, Anthrachnose
European Fan Palm	<u>Chamaerops numilis</u>	Pestalotia Leaf Spot
Gardenia	<u>Gardenia jasminoides</u>	Alternaria Leaf Spot, Botrytis Bud Rot, Cercospora Leaf Spot
Geranium	<u>Pelargonium</u> sp.	Alternaria Leaf Spot, Botrytis Gray Mold, Cercospora Leaf Spot
Gladiolus	<u>Gladiolus</u> sp.	Alternaria Leaf Spot, Botrytis Gray Mold, Bacterial Leaf Blight
Golden Rain Tree	<u>Koeleruteria paniculata</u>	Bacterial Leaf Spot
Hibiscus	<u>Hibiscus rosa-sinensis</u>	Bacterial Leaf Spot
Holly Fern	<u>Cyrtomium falcatum</u>	Pseudomonas Leaf Spot
Impatiens	<u>Impatiens sallerana</u>	Bacterial Leaf Spot
India Hawthorn	<u>Raphiolepis indica</u>	Anthrachnose, Entomosporium Leafspot
Ivy (English, Algerian)	<u>Hedera helix</u> , <u>H. canariensis</u>	Xanthomonas Leaf Spot
Ixora	<u>Ixora coccinea</u>	Xanthomonas Leaf Spot
Juniper (Eastern Red Cedar)	<u>Juniperus virginiana</u>	Anthrachnose
Lantana	<u>Lantana camara</u>	Bacterial Leaf Spot
Lilac	<u>Syringa</u> sp.	Cercospora Leaf Spot
Loblolly Bay	<u>Gordonia lasianthus</u>	Anthrachnose
Loquat	<u>Eriobotrya japonica</u>	<u>Entomosporium maculata</u> , <u>Colletotrichum</u> sp.
Mandevillas	<u>Mandevilla</u> sp.	Anthrachnose
Magnolia (Southern)	<u>Magnolia grandiflora</u>	Algal Leaf Spot, Anthrachnose, Bacterial Leaf Spot
Magnolia (Sweet Bay)	<u>Magnolia virginiana</u>	Anthrachnose
Magnolia	<u>Magnolia soulangiana</u>	Bacterial Leaf Spot
Marigold	<u>Tagetes</u> sp.	Alternaria Leaf Spot, Botrytis Leaf and Rot, Cercospora Leaf Spot
Mulberry, Weeping	<u>Morus alba</u>	Bacterial Leaf Spot
Oleander	<u>Nerium oleander</u>	Bacterial Leaf Spot, Fungal Leaf Spot
Osage Laurel	<u>Quercus laurifolia</u>	Algal Leaf Spot ( <u>Cephaleuros virescens</u> )
Pachysandra	<u>Pachysandra procumbens</u>	Volutella Leaf Blight
Pansy	<u>Viola</u> sp.	Downy mildew
Pear (Flowering)	<u>Pyrus calleryana</u>	Fireblight, Leaf Spot
Peony	<u>Paeonia</u> sp.	Botrytis blight
Pentas (Egyptian Star)	<u>Pentas</u> sp.	

Crop  
 Periwinkle  
 Phlox  
 Pistachio  
 Plantain lily  
 Powder Puff Plant  
 Ilodendron  
 Photinia ("Red Tip," "Red Leaf")  
 Pyracantha  
 Queen Palm  
 Rhododendron  
 Rose  
 Verbena  
 Viburnum  
 Washingtonia Palm  
 Weeping Willow  
 Yucca (Adam's needle)

Latin Name  
Catharanchus roseus, Vinca sp.  
Phlox sp.  
Pistacia chinensis  
Hosta sp.  
Callindra sp.  
Philodendron selloum  
Photinia fraseri, P. glabra  
Pyracantha sp.  
Arecastrum romanzoffianum  
Rhododendron sp.  
Rosa sp.  
Verbena sp.  
Viburnum odoratissimum, V. suspensum  
Washingtonia robusta  
Salix babylonica  
Yucca sp.

Disease  
 Bacterial Leaf Spot (Xanthomonas sp.)  
 Phomopsis Stem Blight  
 Alternaria Leaf Spot  
 Anthracnose  
 Bacterial Leaf Spot  
 Bacterial Leaf Spot  
 Bacterial Leaf Spot  
 Anthracnose, Entomosporium  
 Fireblight, Scab  
 Exosporium Leaf Spot, Phytophthora Bud Rot  
 Alternaria Flower Spot  
 Powdery Mildew, Black Spot  
 Xanthomonas Leafspot  
 Anthracnose  
 Pestalotia Leaf Spot  
 Anthracnose  
 Cercospora and Septoria Leaf Spot

12/13

- (1) Discoloration of foliage and/or blooms have been noted on some varieties. To prevent residues on commercial plants, do not spray just before selling season.
- (2) Apply 2.25 to 3.75 pounds of Kocide 2000 in 20 to 100 gallons of water per acre.
- (3) For Indian Hawthorn use 1.5 to 3 pounds per 100 gallons or 0.75 to 1.5 level tablespoons per gallon.

## GENERAL CHEMIGATION INSTRUCTIONS

Apply this product only through one or more of the following types of sprinkler irrigation systems: center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set or hand move. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Shut off injection equipment after treatment and continue to operate irrigation system until Kocide 2000 has been cleared from the last sprinkler head.

**NOTE:** It must be determined in the selection process if proper application equipment is available and if the waste associated with its use can be properly handled. Materials used in the construction of application equipment is also an important factor as agricultural chemicals are often reactive with soft metals such as aluminum and even some synthetic materials such as plastics, rubbers, etc. Therefore it is necessary when working with equipment containing these materials that they are thoroughly flushed with clean water after each days use.

Posting of areas to be chemigated is required when 1) any part of a treated area is within 300 feet of sensitive areas such as residential areas, labor camps, businesses, day care centers, hospitals, in-patient clinics, nursing homes or any public areas such as schools, parks, playgrounds or other public facilities not including public roads, or 2) when the chemigated area is open to the public such as golf courses or retail greenhouses.

Posting must conform to the following requirements. Treated areas shall be posted with signs at all usual points of entry and along likely routes of approach from the listed sensitive areas. When there are no usual points of entry, signs must be posted in the corners of the treated areas and in any location affording maximum visibility to sensitive areas. The printed side of the sign should face away from the treated area towards the sensitive area. The signs shall be printed in English. Signs must be posted prior to application and must remain until foliage has dried and soil surface water has disappeared. Signs may remain in place indefinitely as long as they are composed of materials to prevent deterioration and maintain legibility for the

duration of the posting period.

All words shall consist of letters at least 2 1/4 inches tall and all letters and the symbol shall be a color which sharply contrasts with their immediate background. At the top of the sign shall be the words KEEP OUT, followed by an octagonal stop sign symbol at least 8 inches in diameter containing the word STOP. Below the symbol shall be the words PESTICIDES IN IRRIGATION WATER.

This sign is in addition to any sign posted to comply with the Worker Protection Standard.

## CHEMIGATION SYSTEMS CONNECTED TO PUBLIC WATER SYSTEMS

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into the reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

When mixing, fill nurse tank half full with water. Add Kocide 2000 slowly to tank while hydraulic or mechanical agitation is operating and continue filling with water. DO NOT PRE-MIX OR SLURRY Kocide 2000. Stickers, spreaders (cleared for use on growing crops), nutrients, etc. should be added last. If compatibility is in question, use the Compatibility Jar Test before mixing a whole tank. Because of the wide variety of possible combinations which can be encountered, observe all cautions and limitations on the label of all products used in mixtures.

Kocide 2000 should be added through a traveling irrigation system continuously or at the last 30 minutes of solid set or hand moved irrigation systems. Agitation of the mixture in the nurse tank is recommended.

## SPRINKLER CHEMIGATION

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

When mixing, fill nurse tank half full with water. Add Kocide 2000 slowly to tank while hydraulic or mechanical agitation is operating and continue filling with water. DO NOT PRE-MIX OR SLURRY Kocide 2000 before adding to the nurse tank. Stickers, spreaders (cleared for use on growing crops), nutrients, etc. should be added last. If compatibility is in question, use the Compatibility Jar Test before mixing a whole tank. Because of the wide variety of possible combinations which can be encountered, observe all cautions and limitations on the label of all products used in mixtures.

Kocide 2000 should be added through a traveling irrigation system continuously or at the last 30 minutes of solid set or hand moved irrigation systems. Agitation of the mixture in the nurse tank is recommended.

## NOTICE OF WARRANTY

13/13

GRIFFIN warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials or the manner of use or application, all of which are beyond the control of GRIFFIN. In no case shall GRIFFIN be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. GRIFFIN MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

Aliette is a registered trademark of Rhone-Poulenc.

Kocide is a registered trademark of Griffin Corporation.

Rovral is a registered trademark of Rhone-Poulenc.

GCN 121295  
CPC 038438

ACCEPTED  
with COMMENTS  
In EPA Letter Dated

DEC 31 1996  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

1512-358