



U.S. ENVIRONMENTAL PROTECTION AGENCY  
Office of Pesticide Programs  
Registration Division (7505C)  
401 "M" St., S.W.  
Washington, D.C. 20460

EPA Reg. Number:  
**1812-366**

Date of Issuance:  
**NOV - 6 1995**

**NOTICE OF PESTICIDE:**  
XX Registration  
Reregistration

Term of Issuance:  
**CONDITIONAL**

Name of Pesticide Product:  
**Cynex 4L**

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

**Griffin Corporation  
P.O. Box 1847  
Valdosta, GA 31603-1847**

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

- I. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
- II. Make the following label changes:
  - A. On page 1:
    - 1) Revise the EPA Registration Number to read, "EPA Reg. No. 1812-366".
    - 2) Enclose the Restricted Use Pesticide statements in a box per PR Notice 93-1 and 40 CFR 156.10(j).
    - 3) Under **WARNING**, modify the last word in the first sentence from "detaile" to "detalle".
    - 4) Under the **STATEMENT OF PRACTICAL TREATMENT (IF IN EYES:)** section, delete the second "f" in "Flush".
    - 5) You must include the Net Contents as prescribed in paragraph (d) of 40 CFR §156.10.

Continued on page 2

Signature of Approving Official:

Date: **11/6/95**

B. On page 3:

- 1) Move the second paragraph ("When handlers use...") to the top of the page so that it is underneath the heading "Engineering control statements:".
- 2) Under the **DIRECTIONS FOR USE** section, add the sentence "This labeling must be in the possession of the user at the time of pesticide application." after the first sentence. Move the following sentences to this section: "Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation."
- 3) In the **AGRICULTURAL USE REQUIREMENTS** box, delete the second comma after "nurseries," in the second sentence. Add the following bullets to the list of required PPE for early entry to treated areas: "-Shoes plus socks." and "-Protective eyewear."

C. On page 4:

- 1) In the **STORAGE AND DISPOSAL** box, add the heading "**STORAGE**" in front of the first sentence.
- 2) Under the **BEST MANAGEMENT PRACTICES FOR GROUND AND SURFACE WATER PROTECTION** section, correct the spelling of "perennial" in the first sentence. Move the last paragraph and make it the second paragraph.

D. Delete all references to grain sorghum (milo) (except in the rotational crop section) and wheat from the label because they are considered new Me-Too uses since they are not registered for the product cited as being substantially similar to your product. Also, delete all references to the peanut and soybean uses from the label because they are considered new uses and must be supported with additional data.

E. On pages 5 and 6:

- 1) Under the **GENERAL INFORMATION (Weather Effects:)** section, correct the spelling of "hoeing".
- 2) Under **APPLICATION DIRECTIONS (General)** section,
  - a. Under 1., change "7.5" to "10" gallons of water per acre or provide justification as part of an amendment to registration as to why the lower rate should/can be used.

F. On page 7:

- 1) Under 2., add "to corn." to the end of the first sentence.

- 2) Under 8. and 13., change "emulsible oil" to "emulsible crop oil" and "oil concentrate" to "crop oil concentrate".
- 3) Under 1!., correct the spelling of "formulations" in the first sentence.

G. On page 8:

- 1) Delete the high rate (40 gallons of liquid carrier per acre and .06 Teaspoons of Cynex 4L per "pint" (correct spelling) of Liquid Carrier) from the **Jar Test for Cynex 4L Compatibility** chart or submit as an amend-ment, the justification for this higher rate.
- 2) Check periods after all numbers and letters.

H. On page 10:

- 1) Move the period next to "A" and change the last two words to "impregnating fertilizer".
- 2) Delete the extra "4l" in 3.
- 3) Modify B. to read "Fertilizer impregnated with Cynex 4L may be used in tank mixes with other **dry** herbicides."
- 4) Delete the statement "(For more information..., see Du Pont Bulletin...) unless you intend to provide the technical bulletin with your product.

I. On page 11: Modify the first sentence to read "Apply fertilizer immediately after impregnation."

J. On page 13:

- 1) Under the Cynex 4L plus Atrazine section, you must specify by product name all applicable atrazine tank mix products.
- 2) Consolidate all paragraphs relating to Rotational Crops under one section. Clarify which "tank mixtures" are associated with the rotational crops.

K. On page 14: Under the first chart, correct the spelling of "between" after \*\*. In addition, make sure all of the rates are correct for both Cynex 4L alone and with tank mixes throughout the label.

L. On page 18: Under **CONSERVATION TILLAGE WEED CONTROL**, define early preplant (EPP) as "(30 days prior to planting until emergence".

M. On page 30:

- 1) On Table 13, add "height of cotton band 6" or more".

1/2/99

- 2) On Table 14, add "silt and silt loam" to the coarse soil texture part of table and delete from medium part.
  - 3) On Table 15, define under "Product" heading "quarts/acre".
- N. Please note that propazine is not a registered active ingredient for use on sorghum.

- O.
- 1) List specific product names for the 2,4-D LV ester and paraquat tank mix products.
  - 2) Check all tank mix products rates and use directions throughout the label to make sure they are similar to their product label language and that they are compatible with Cynex 4L. According to PR Notice 82-2, the tank mix directions must include statements similar to the following:

"This product can be mixed with \_\_\_\_\_ (chemical name, including percentage of active ingredient and type of formulation, or specific product name, or both) for use on \_\_\_\_\_ (crops/sites) in accordance with the more (most) restrictive of label limitations and precautions. No label dosage rates should be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing.

III. The Griffin Corporation letter dated September 5, 1995 agreed to the terms and conditions in the DuPont/Agency cyanazine phase-out agreement which was approved on August 2, 1995. You must include in all of your cyanazine registrations that you agree to the terms and conditions set forth in sections 3., 4., 5., 6., 7., and 8. of that Cyanazine Phase-Out Agreement. Some of those terms include the following:

- A. The labels of all cyanazine formulated end-use products released for shipment after July 25, 1996, for use in the U.S., must be amended as follows:
- (1) Limit the maximum use rates from the current 6.5 lbs./acre to 5 lbs./acre beginning January 1, 1997; 3 lbs./acre beginning January 1, 1998; and 1 lb./acre beginning January 1, 1999 through December 31, 2002.
  - (2) Specify that closed cab application will be required for applications to be made during or after the 1998 use season.
  - (3) Add the following statements: "This product may not be sold or distributed after September 30, 2002." and "This product may not be used after December 31, 2002."

- B. No cyanazine formulated end use products registered for use in the U.S. shall be released for shipment by a registrant after December 31, 1999.
  - C. Existing stocks of all cyanazine formulated end use products that have been released for shipment by a registrant on or before December 31, 1999, may continue to be distributed and sold in the channels of trade in accordance with their labels through September 30, 2002. The use of such existing stocks may continue in accordance with their labels through December 31, 2002.
  - D. The voluntary cancellation date of December 31, 1999 shall become a part of the terms and conditions of all cyanazine registrations.
- IV. Submit two copies of the revised final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

6/29

**RESTRICTED USE PESTICIDE**

This product is a restricted use herbicide due to reproductive and ground and surface water concerns. Users must read and follow all precautionary statements and instructions for use in order to minimize potential for Cyanazine to reach ground and surface water.

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

**Cynex 4L  
HERBICIDE LIQUID**

**ACTIVE INGREDIENTS:**

Cyanazine: 2[[4-chloro-6-(ethylamino)-s-triazin- 2-yl]amino]-2-methylpropionitrile.....43%

**INERT INGREDIENTS:**.....57%

**TOTAL**.....100%

Contains 4 lbs. active ingredient per gallon.

**KEEP OUT OF REACH OF CHILDREN**

**WARNING -AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

**STATEMENT OF PRACTICAL TREATMENT**

**IF SWALLOWED:** Call a physician or poison control center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

**IF IN EYES:** Flush with plenty of water. If irritation persists, call a physician.

**IF ON SKIN,** wash immediately with plenty of soap and water.

**IF INHALED,** remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth and get medical attention.

**ACCEPTED  
with COMMENTS  
In EPA Letter Dated  
NOV - 6 1995**

**Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.**

1812-366

GRIFFIN CORPORATION  
Valdosta, GA 31601

EPA REG. NO. 1812-

17 99

(

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS (AND DOMESTIC ANIMALS)  
WARNING**

May be fatal if swallowed. Harmful if inhaled or absorbed through the skin. Causes temporary eye injury. This product may be hazardous to your health. This product is classified "Restricted Use" because, at doses which caused serious maternal illness in laboratory animals, birth defects were present. Use of protective clothing and equipment and following the precautions below can reduce risk. Avoid breathing spray mist. Avoid contact with skin, eyes, or clothing. Do not get in eyes or on clothing.

Keep out of reach of domestic animals, particularly cattle. Consumption of this product, spray solutions, or water contaminated with product can result in serious illness or possible death of bovines.

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning equipment or disposal of wastes. Cyanazine, the active ingredient in Cynex 4L has been detected in surface waters that receive run-off from treated areas. To minimize cyanazine run-off, follow the Best Management Practices outlined in the Directions For Use section of this label.

Cyanazine is a chemical which can move (seep or travel) through soil and can contaminate groundwater which may be used as drinking water. Cyanazine has been found in groundwater as a result of agricultural use. Ground water contamination may be reduced by diking and flooring of permanent liquid bulk storage sites with an impermeable material. Users are advised not to apply Cynex 4L where the water table (groundwater) is close to the surface and where the soils are very permeable (i.e., well drained soils such as loamy sands). Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater.

**PERSONAL PROTECTIVE EQUIPMENT**

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or polyvinyl chloride or viton or neoprene rubber
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant apron when cleaning equipment, mixing or loading

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering control statements:

**USER SAFETY RECOMMENDATIONS**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove personal protective equipment immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR part 170.240 (d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls.
- Chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or polyvinyl chloride or viton or neoprene rubber.

Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.



## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Do not use or store around the home environment. Avoid contact with water. In case of spill or leak, soak up with sand, earth or synthetic absorbant (do not use alkaline absorbants) and dispose of wastes in compliance with local, State or Federal regulations. Ground water contamination may be reduced by diking and flooring of permanent liquid bulk storage sites with an impermeable material.

**PESTICIDE DISPOSAL:** Pesticide, spray mixture or rinsate that cannot be used according to label instructions must be disposed of according to applicable Federal, State or local procedures.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

## BEST MANAGEMENT PRACTICES FOR GROUND AND SURFACE WATER PROTECTION

This product may not be mixed or loaded within 50 feet of perennial or intermittent streams and rivers, natural or impounded lakes, and reservoirs. This product may not be mixed, loaded or used within 50 feet of all wells, including abandoned wells, drainage wells, and sinkholes.

Operations that involve mixing, loading, rinsing, or washing of this product into or from pesticide handling or application equipment or containers within 50 feet of any well are prohibited unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or wash-water, and rainwater that may fall on the pad. Surface water shall not be allowed to either flow over or from the pad, which means the pad must be self-contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain, at a minimum, 110% of the capacity of the largest pesticide container or application equipment on the pad. A pad that is covered by a roof or sufficient size to completely exclude precipitation from contact with the pad shall have a minimum containment capacity of 100% of the capacity of the largest pesticide container or application equipment on the pad. Containment capacities as described above shall be maintained at all times. The above-specified minimum containment capacities do not apply to vehicles when delivering pesticide shipments to the mixing/loading site.

States may have in effect additional requirements regarding well-head setbacks and operational area containment.

This product may not be applied aerially or by ground within 66 feet of the points where field surface water run-off enters perennial or intermittent streams and rivers or within 200 feet of natural or impounded lakes and reservoirs. This product may only be applied to highly erodible land if the 66 foot buffer or set-back from run-off points is planted to crop or seeded with grass.

**Cyanazine Rate Limits:** one quart of Cynex 4L contains 1.0 lb cyanazine active ingredient (a.i.). Adhere to the use rate recommendations in this or other label. In addition:

- a. Do not apply more than 6.5 lbs. total cyanazine a.i. (all sources) per acre per year to any land.
- b. On highly erodible land, as defined by the Soil Conservation Service, if plant residue cover is less than 30%, do not apply more than 3.0 pounds total cyanazine a.i. (all sources) per acre per year.

Where there are state/local requirements regarding cyanazine use (including lower maximum rates and/or higher setbacks) which are different from the label, the more restrictive/protective requirements apply.

### GENERAL INFORMATION

Cynex 4L Herbicide is a selective herbicide for the control of annual grasses and broadleaved weeds in field corn, popcorn, sweet corn, cotton and grain sorghum.

Consult your local Agricultural Extension Agent for help in determining soil texture, organic matter content, and the most appropriate herbicide rate for local conditions.

Do not apply this product in irrigation water with any kind of irrigation system.

Do not apply this product with aerial application equipment.

Cynex 4L is not effective when used preemergence on peat or muck soils. DO not use Cynex 4L on sands or loamy sands (soils consisting of more than 70% sand) containing less than 1% organic matter.

**Weather Effects:** As a preemergence herbicide, Cynex 4L is active mainly through the roots, and therefore, its effect on weeds is dependent on adequate rainfall or sprinkler irrigation to move the herbicide into the root zone. Moisture should be sufficient to thoroughly wet the soil throughout the zone where weed seeds may germinate and enough to make soil too wet to cultivate. Rotary hosing or shallow cultivation is recommended for those applications which are not incorporated at the time of treatment, if adequate rainfall or sprinkler irrigation has not occurred within about ten days after application of Cynex 4L.

Heavy rainfall between planting and crop emergence may cause crop injury or stand loss. Rainfall tends to cause excessive concentrations of herbicide in the seed furrow, resulting in possible crop injury. Level deep planter marks or seed furrows before application.

Under conditions which delay weed germination, such as low temperatures, lack of soil surface moisture, or when germination extends over a long period, the effectiveness of the herbicide may be

11/19  
impaired. Rotary hosing, shallow cultivation or a postemergence herbicide treatment may be of benefit under these circumstances.

If the crop is cultivated, tillage should be shallow to minimize herbicide dilution in the soil. Should the crop stand be lost due to adverse weather conditions, insects, etc., the field can be replanted the same season to corn or sorghum.

To enhance weed control in areas of less than 25 inches of rainfall or where long dry periods are common, these treatments may require shallow incorporation with a tool such as a field cultivator operated from 5-7 mph. Incorporation should not be more than three inches deep to keep from burying the herbicide. A spike-toothed harrow, deep tillage disk or rolling basket device is not recommended for incorporating Cynex 4L.

When applied as a post-emergence herbicide, Cynex 4L is also active through foliage as well as through the roots. Yellowing and/or stunting of the crop may result from this treatment, particularly if cold, adverse growing conditions occur after application. Extended or extreme cold and wet conditions may reduce stands. Do not apply Cynex 4L postemergence to a crop that is damaged or growing under stress.

#### **OBSERVE ALL CAUTIONS AND LIMITATIONS ON LABELING OF ALL PRODUCTS USED IN MIXTURES.**

**Triazine Resistant Weeds:** In fields where triazine resistant biotypes of weeds have been identified, Cynex 4L should be used in combination with or in sequence with other registered non-triazine herbicides. (Triazine resistant biotypes of kochia and pigweed have been identified in some fields in the Western Great Plains and triazine resistant biotypes of pigweed and lambquarters have been identified in some fields in various states.) Consult with appropriate state agricultural extension service representatives for specific recommendations.

#### **APPLICATION DIRECTIONS**

##### **General mixing and spraying instructions**

This product may not be applied by means of chemigation or aerial application.

Use sufficient agitation to ensure that the Cynex 4L is completely dispersed and in uniform suspension prior to application or tank mixing with other formulations.

The following general mixing instructions are recommended when using this or any other liquid suspension formulation.

##### **I. General**

1. Unless otherwise specified, use at least 7.5 gallons of water per acre for soil applications and at least 15 gallons of water per acre for foliar applications for all applications with ground equipment. **NOTE:** Sufficient carrier must be used to assure uniform application. Follow label requirement of all products used in tank mix combinations.

2. A nitrogen solution or complete liquid fertilizer may replace all or part of the water as a carrier for preemergence or preplant application. Do not apply fertilizer mixtures after crop emerges, because injury may occur.
3. Always check the tank mix compatibility (TMC) of this or any other formulation before mixing with liquid fertilizer carriers or other formulations. A simple but generally reliable TMC evaluation procedure has been provided for your use in step II of these mixing instructions.
4. Start with thoroughly clean equipment. (See the label of previous compound used for cleaning instructions.)
5. Fill tank 1/2 full with carrier. Start and maintain consistent agitation through all mixing and spraying procedures. Make sure that the agitation system is working properly and creates a rippling or rolling action on the liquid surface.
6. Add the Cynex 4L to the tank with agitation.
7. Fill tank to 75 percent capacity with carrier. Filling and bypass lines should be kept below liquid surface. Increase tank agitation if necessary to maintain surface action.
8. When desired, appropriate emulsible oil, oil concentrate, or other tank mix formulations should be added at this time. Pre-slurry these added ingredients before addition, if the compatibility test shows it to be necessary.
9. Complete filling tank maintaining sufficient agitation at all times to ensure surface action. This applies to both spray and nurse tanks.
10. Tank mixtures should always be applied immediately after preparation. If for any reason this is not possible, assure that sufficient agitation has been provided to re-mix all products and check for complete resuspension prior to application.
11. Empty tank as completely as possible before refilling to prevent buildup of oil or emulsible concentrate residues when tank mixing with these formulations. Always maintain agitation to avoid separation.
12. If an oil or emulsible concentrate film starts to build up after using these formulations, drain and clean the tank with strong detergent solution or appropriate solvent.
13. It is recommended that the sprayer be thoroughly cleaned by flushing with a detergent solution at the end of each work day when any emulsible oil, oil concentrate, or other emulsible formulation has been used either alone or in tank mix combinations with other pesticide formulations, even if no obvious problems have been encountered. This precaution will ensure a clean sprayer and continued trouble-free operation.

**II. Tank Mix Compatibility Evaluation Procedure**

1. Add one pint of carrier liquid to each of two one-quart jars. Mark one quart jar "with" and the other "without".
2. Add 1/4 teaspoon of a suitable tank mix compatibility agent (1/4 teaspoon/pint = 2 pints/100 gallons of carrier) to the jar marked "with," cap the jar, and shake gently for five to ten seconds to mix.
3. Add the appropriate amount of herbicide to both jars, cap each jar, and shake gently for five to ten seconds to mix. If problems are encountered in mixing wettable powder or dry flowable formulations into a liquid fertilizer, then pre-slurry these formulations in water prior to their addition to the liquid fertilizer and proceed with the test. The following chart has been provided to assist you in selecting the appropriate Cynex 4L use rate for this evaluation. If more than one herbicide is to be used in the tank mixture, each should be added separately as follows: water solubles first, wettable powders or dry flowables second, liquid flowables third, and

emulsible concentrate or oil formulations last with each jar capped and gently shaken for five to ten seconds for each addition.

**Jar Test for Cynex 4L Compatibility**

Gallons of Liquid Carrier per Acre	4	7.5	15	20	25	30	40
Teaspoons of Cynex 4L per pint of Liquid Carrier	6.0	3.2	1.6	1.2	1.0	0.8	0.6

This chart is based on one quart of Cynex 4L (1 pound active ingredients) per acre in the indicated carrier volumes. Intended field use rates are achieved by varying the amount of Cynex 4L; i.e., for a field use rate of 2.5 quarts of Cynex 4L in 15 gallons of carrier per acre, add 4 teaspoons of Cynex 4L to the quart jars containing one pint of carrier. Calculation: For 2.5 quarts of Cynex 4L/15 gallons of carrier per acre, multiply 2.5 X 1.6 to get 4 teaspoons of Cynex 4L per pint of carrier.

4. Let each jar stand one-half hour and make observations. If any separation, agglomeration, or precipitation has occurred, shake the jar again for 10 to 15 seconds, and note whether any of the following occur: a. Separated phases do not re-mix uniformly. b. Screen/ nozzle plugging lumps do not disperse. c. Precipitate does not resuspend readily. d. Precipitate sticks tenaciously to glass.
5. If none of the above problems occur in either jar, then the herbicides can, in most cases, be safely used without a compatibility agent.
6. If problems 4.a or 4.b occur in the jar marked "without" but do not occur in the jar marked "with," the compatibility agent should be used.
7. If problems 4.a or 4.b are seen in both jars, then the herbicides and carrier mixture are incompatible and should not be used in the same spray tank. Alternatively, a different tank mix compatibility agent can be evaluated.
8. If problems 4.c or 4.d occur in the jar marked "without" but do not occur in the jar marked "with," the compatibility agent should be used unless constant, thorough agitation can be maintained and immediate clean-out of spray system is performed.
9. If problems 4.c and 4.d are seen in the jar marked "with," the user proceeds with mixing and application at his own risk should agitation in the system be insufficient or curtailed.
10. Those mixtures defined as compatible in this test should then be mixed for use as indicated in steps 1-12 of the general mixing instructions listed above.

If a test such as outlined indicates that components of a proposed mix are compatible, the applicator still has the responsibility of combining materials in sequence to the spray tank in accordance with directions prescribed on the label of the herbicides or pesticides involved.

Tests have indicated that compatibility agents, noted below by the various tank mix combinations, may give improved compatibility in liquid fertilizers.

**Tank Mix Combination**

**Compatibility Agents**

Cynex 4L/LASSO (Liquid Fertilizer Grade)	Probably not needed in 28-0-0, 10-34-0. "Compex" may help in others.
Cynex 4L/SUTAN + 6.7E	Probably not needed in 28-0-0. Incompatible in 10-34-0. "Unite", "Spray-Mate", "Kem-Link", may help in others.
Cynex 4L/DUAL 8E	Probably not needed in 28-0-0. "Unite", "Spraymate", "Ivory Liquid" may help in others.

### III. Application Equipment

1. Use application equipment fitted with nozzles that provide accurate and uniform coverage. Be certain that nozzles are uniformly spaced and the same size. Calibrate sprayer before use and recheck frequently during use whenever possible.
2. Use a pump with capacity to:
  - a. Maintain 35-40 psi at nozzles.
  - b. Provide sufficient agitation in tank to keep mixture in suspension.
  - c. Provide a minimum of 20 percent bypass at all times.
3. Use centrifugal pumps which provide sufficient shear action for dispersing and mixing this product. The pump should provide a minimum of 10 gallons/minute/100-gallon tank size circulated through the jets of a correctly-positioned sparger tube.
4. Use screens to protect the pump and to prevent nozzles from clogging. Screens placed on suction side of pump should be 10 to 16 mesh. do not place a screen in the recirculation line. Use a 40- to 50-mesh screen between the pump and boom and, where required, 50-mesh screens at the nozzles. Check your equipment manufacturer's literature for specific recommendations.

### FERTILIZER IMPREGNATION

Cynex 4L Herbicide may be applied when coated on or impregnated in dry granular fertilizer for early preplant, preemergence or preplant incorporated weed control in field corn. All recommendations, cautions and special precautions on this label must be followed along with state regulations relating to dry bulk fertilizer blending, impregnating and labeling. Also, follow the precautions on the label of any product mixed with this product.

#### General Blending Directions

Cynex 4L may be coated on or impregnated in dry bulk fertilizers using tower blenders, rotary drum blenders or blending augurs or conveyors.

DO NOT impregnate Cynex 4L or tank mixes containing Cynex 4L on or in fertilizers containing Ammonium Nitrate, Potassium Nitrate, or Sodium Nitrate. Do not use on straight limestone since absorption will not be achieved. Fertilizer blends containing limestone can be impregnated when using Cynex 4L alone. Use a minimum of 200 lbs. and a maximum of 450 lbs. per acre of dry fertilizer.

1. Use equipment that will give a uniform distribution of the herbicide throughout each batch of impregnated fertilizer. Non-uniform impregnation can cause crop injury or unsatisfactory performance.

A .Cynex 4L may be used as the only herbicide for impregnation.

1. Add Cynex 4L when at least 1/2 the total fertilizer volume required is in the mixer. A minimum of 200 lbs per acre of an approved fertilizer should be used.
2. Cynex 4L can be moved from the chemical holding tank to the mixer by using an air system or a liquid pump with a hose at least 1 inch in diameter.
3. Position the spray nozzles to achieve uniform coverage of the Cynex 4L 41 on the dry fertilizer without spraying the walls of the mixer.
4. Flush the spray lines with water and spray onto the dry fertilizer.
5. Add remaining fertilizer, plus drying agents when necessary, and blend thoroughly for at least three minutes.
6. Add 2-5% of a suitable drying agent to insure a herbicide/fertilizer mixture that will spread through air spreaders. The need for a drying agent is determined by the wetness of the fertilizer batch. Wetness can change with humidity, nitrogen content, fertilizer rates and herbicide rates.

B .Cynex 4L may be used in tank mixes with other herbicides.

1. While fertilizer is blending, add the Cynex 4L first and the tank mix fertilizer last.
2. Add any necessary drying agent to insure a spreadable herbicide/ fertilizer mixture and blend thoroughly for at least three minutes.
3. Follow the other appropriate use instructions found in Section A for use with Cynex 4L tank mixes.

(For more information on Drying Agents, Application Equipment, Calibration Guide and variations of these methods, see Du Pont Bulletin on "Fertilizer Impregnation").

**CLEAN OUT:**

Equipment used to impregnate or apply fertilizer impregnated with Cynex 4L or combinations including Cynex 4L must be cleaned out by running at least 1,000 lbs. of fertilizer not impregnated with Cynex 4L through the impregnation equipment and application equipment, if the next batch of material is to be applied to a crop for which Cynex 4L or a combination herbicide is not registered.

**APPLICATION:**

Uniform application of Cynex 4L which has been impregnated in or coated on dry fertilizer is essential for satisfactory weed control and crop safety. Accurate calibration of the fertilizer applicator is necessary. Applying while turning at the ends of the fields may result in excessive application rates causing crop injury. Do not double apply across the ends or sides of the field.

Crop injury and/or poor weed control may occur where the impregnated fertilizer is not uniformly applied. Air flow or augur metered application equipment is preferred (one pass application). If other equipment is used, the recommended method of application is to apply 1/2 the recommended rate an overlap 50 percent to double apply by splitting the middles to obtain the best distribution pattern.

Apply immediately after impregnation. Impregnated fertilizer may become lumpy and difficult to spread if stored.

### RATES AND TIMING:

Use the application rates and timing shown in the appropriate sections of this label. Follow the precautions on the labels of all products used.

### CORN

#### WEEDS CONTROLLED BY Cynex 4L ALONE AND IN COMBINATION WITH OTHER HERBICIDES ON CORN

---

	<b>Grasses</b>	
) Annual bluegrass	Bullgrass	Junglerice
Annual fescues	Crabgrass	Stinkgrass
Annual (Italian)	Fall panicum	(Indian lovegrass)
ryegrass	Giant foxtail	Witchgrass
Annual sedge	Goosegrass	Yellow foxtail
Barnyardgrass(1)	Green foxtail	
Bullgrass		
	<b>Broadleaves</b>	
Annual groundcherry	Florida pusley	Ragweed (Common)
Annual morningglory	(Florida purslane)	Russian thistle
Black mustard	Hedge mustard	Shepherdspurge
Buffalobur	Jimsonweed(1)	Smallflower galinsoga
Buttercup (annual)	Kochia	Smartweed
Carpetweed	Ladysthumb	(Pennsylvania)
) Cocklebur(2)	Lambsquarters	Sunflower(2) (wild,
Common chickweed	Mayweed	annual, common)
Common groundsel	Nightshade (annual)	Tarweed cuphea
Common mallow	Pigweed(1)	(Gumweed)
Common purslane	Pineappleweed	Velvetleaf(1)
Corn spurry	Plantain	Wild buckwheat
Curly dock (seedling)	Poorjoe	Wild mustard
Fiddleneck	Prickly sida (teaweed)	Wild radish
	Prostrate knotweed	Wild turnip
	Prostrate spurge	

---

(1) Under conditions such as low temperatures, lack of soil surface moisture or other factors that may cause delay in germination of the seeds, the effectiveness of Cynex 4L may be impaired against these weeds.

(2) The degree of control will be reduced if soil moisture and temperature conditions cause deep germination of the seed.

### CORN



11/1/10

### Preemergence-Preplant Incorporated

Apply Cynex 4L treatments just before, at or after planting but before crop has emerged. Avoid removal of treated soil from seedrow prior to or during the planting operation.

Cynex 4L may also be applied early prior to planting or in a split application if pre-season weed control is desired. For split applications, do not exceed the total amount of Cynex 4L for the soil texture and organic matter shown in Table 1. If Cynex 4L is applied early, more than 15 days before planting, a split application of Cynex 4L or some other herbicide treatment may be necessary at or after planting to provide additional length of weed control. For further information see "Early Preplant" recommendations in the Conservation Tillage section of this label.

Rotary hoeing is recommended for preemergence applications which do not receive adequate rainfall or sprinkler irrigation to wet the top 2 inches of soil or depth of germinating weeds within about 10 days after application.

Cynex 4L alone or in tank mix combinations should not be incorporated more than three inches deep to keep from burying the herbicide. Single or two pass incorporation with a tool such as a field cultivator operated at 5-7 mph is acceptable. A spike-toothed harrow, deep tillage disk or rolling basket device is not recommended for incorporating Cynex 4L.

### Cynex 4L Applied Alone

Use the proper rate for the soil texture and organic matter indicated in Table 1. Any rotational crop may be planted in the fall or spring following this treatment.

**TABLE 1  
PREEMERGENCE BROADCAST APPLICATION RATES  
PER ACRE FOR Cynex 4L APPLIED ALONE ON CORN**

Quarts of Cynex 4L***						
Percent Organic Matter in Soil**						
Soil Texture Description	Less Than 1%	1%	2%	3%	4%	5% & Over
Sand, Loamy sand	Do Not Use	1.25	1.5	2.25	2.75	3.25
Sandy loam	1.25	1.75	2.0	2.5	3.0	3.5
Loam, Silt loam, Silt	1.5	2.0	2.5	3.0	3.5	4.0

Sandy clay loam, Clay loam, Silty clay loam	2.0	2.5	3.0	3.5	4.0	4.5
Sandy clay, Silty clay, Clay	2.75	3.0	3.5	4.0	4.5	4.75
Peat or Muck	*	*	*	*	*	*

\*Not Recommended

\*\*For organic matter content between those listed, adjust the rate proportionately.

\*\*\*Maximum rate limit per acre per year for all applications is 6.5 lbs. cyanazine (6.5 quarts Cynex 4L) except on highly erodible land with less than 30% plant residue cover, the rate limit is 3.0 lbs cyanazine (3.0 quarts Cynex 4L).

**Cynex 4L COMBINATIONS**

**Cynex 4L plus Atrazine**

Use Cynex 4L plus atrazine at the proper rate for soil texture and organic matter indicated in Tables 2 and 3. These tables provide rates for generally weedy conditions. The ratio of the amounts of each herbicide may be adjusted as necessary for particular weed conditions as long as the combined rate of the two products does not exceed the combined rate for the soil shown in Table 2 or 3. For grassier conditions use a ratio that contains higher levels of Cynex 4L (3:1). For fields with more broadleaves use a ratio that contains higher levels of atrazine (1:1).

**Rotational Crops:** (1) Plant only corn, peanuts, sorghum, or soybeans the year following the use of this mixture. (2) If soybeans are to be planted, injury may occur due to the carryover of atrazine. (3) If applied after June 10, do not rotate with crops other than corn or sorghum the next year or injury may occur. (4) In the high plains and intermountain areas of the West where rainfall is sparse and erratic or where irrigation is required, use only when corn or sorghum is to be planted the following year, or a crop of corn or sorghum not treated with atrazine is to precede other rotational crops. (5) Small grains may be planted 15 months following treatment. (6) All other crops may be planted 18 months after application.

For all states except Kentucky, Missouri, Tennessee and Kansas east of Highway 99 use Table 2.

In Kentucky, Missouri, Tennessee and Kansas east of Highway 99 use Table 3.

**TABLE 2  
PREEMERGENCE BROADCAST APPLICATION RATES PER ACRE FOR  
TANK-MIX COMBINATIONS OF Cynex 4L PLUS ATRAZINE 4L ON CORN  
FOR USE IN ALL STATES EXCEPT KENTUCKY, MISSOURI, TENNESSEE AND  
KANSAS EAST OF HIGHWAY 99**

Quarts of Cynex 4L **** + Quarts of Atrazine 4L***						
Percent Organic Matter in Soil**						
Soil Texture Description	Less than 1%	1%	2%	3%	4%	5% & Over
Sand, Loamy Sand	DO NOT USE	0.8 + 0.5	1.0 + 0.5	1.5 + 0.5	1.8 + 0.8	2.25 + 1.0
Sandy Loam	0.8 + 0.5	1.0 + 0.5	1.5 + 0.5	1.8 + 0.8	2.25 + 1.0	2.75 + 1.25
Loam, Silt Loam, Silt	1.0 + 0.5	1.5 + 0.5	2.0 + 0.8	2.25 + 1.0	2.75 + 1.25	3.25 + 1.25
Sandy Clay Loam, Clay Loam, Silty Clay Loam	1.5 + 0.5	2.0 + 0.8	2.25 + 1.0	2.75 + 1.25	3.25 + 1.25	3.50 + 1.25
Sandy Clay, Silty Clay, Clay	2.0 + 0.8	2.25 + 1.0	2.75 + 1.0	3.25 + 1.25	3.5 + 1.25	3.75 + 1.5
Peat or Muck	*	*	*	*	*	*

\* Not Recommended

\*\*For organic matter content between those listed, adjust the rate proportionately.

\*\*\*If Atrazine 80W is used, multiply rates shown by 1.25 to equal pounds of Atrazine 80W. If Atrazine 90% is used, multiply rates shown by 1.11 to equal pounds of Atrazine 90%.

\*\*\*\*Maximum rate limit per acre per year for all applications is 6.5 lbs. cyanazine (6.5 quarts Cynex 4L) except on highly erodible land with less than 30% plant residue cover, the rate limit is 3.0 lbs. cyanazine (3.0 quarts Cynex 4L).

**TABLE 3  
PREEMERGENCE BROADCAST APPLICATION RATES PER ACRE FOR  
TANK-MIX COMBINATIONS OF Cynex 4L PLUS ATRAZINE 4L ON CORN  
FOR USE ONLY IN ALL KENTUCKY, MISSOURI, TENNESSEE AND KANSAS  
EAST OF HIGHWAY 99**

Quarts of Cynex 4L**** + Quarts of Atrazine 4L***						
Percent Organic Matter in Soil**						
Soil Texture Description	Less Than 1%	1%	2%	3%	4%	5% & Over
Sand, Loamy Sand	DO NOT USE	0.8 + 0.5	1.0 + 0.5	1.5 + 0.6	1.8 + 0.8	2.25 + 1.0
Sandy Loam	0.8 + 0.5	1.4 + 0.7	1.6 + 0.8	1.8 + 0.9	2.2 + 1.1	2.8 + 1.3

Loam, Silt Loam, Silt	1.4 + 0.7	2.0 + 1.0	2.2 + 1.1	2.4 + 1.2	2.8 + 1.3	3.25 + 1.3
Sandy Clay Loam, Clay Loam, Silty Clay Loam	1.6 + 0.8	2.2 + 1.1	2.4 + 1.2	2.8 + 1.3	3.25 + 1.3	3.5 + 1.4
Sandy Clay, Silty Clay, Clay	2.0 + 1.0	2.4 + 1.2	2.8 + 1.3	3.25 + 1.3	3.5 + 1.4	3.75 + 1.5
Peat or Muck	*	*	*	*	*	*

\*Not Recommended

\*\*For organic matter content between those listed, adjust the rate proportionately.

\*\*\*If Atrazine 80W is used, multiply rates shown by 1.25 to equal pounds of Atrazine 80W.

If Atrazine 90% is used, multiply rates shown by 1.11 to equal pounds of Atrazine 90%.

\*\*\*\*Maximum rate limit per acre per year for all application is 6.5 lbs cyanazine (6.5 quarts Cynex 4L) except on highly erodible land with less than 30% plant residue cover, the rate limit is 3.0 lbs. cyanazine (3.0 quarts Cynex 4L).

### Cynex 4L plus LASSO 4EC

Use Cynex 4L at the proper rate for the soil texture and organic matter shown in Table 4 plus 2 quarts per acre of LASSO . (Use 2.5 quarts of LASSO on clay soils containing 5 percent organic matter and over.) Any rotational crop may be planted the fall or spring following this treatment.

### Cynex 4L plus SUTAN+ 6.7E or ERADICANE 6.7E

Use Cynex 4L at the proper rate for the soil texture and organic matter shown in Table 4 plus 1.8 quarts per acre of SUTAN+ or ERADICANE for control of many annual grasses and broadleaf weeds. (Use 2.4 quarts of SUTAN+ or ERADICANE on loam soils containing 5 percent or more organic matter, and on clay loams and clays containing 4 percent or more organic matter.) Do not use on sands and loamy sands having less than 1 percent organic matter in the light sandy soils of eastern coastal states. Do not use on corn seed stock.

Apply before planting. Incorporate the mixture immediately after application using power-driven cultivation equipment set for 2-3 inches in depth, or a tandem disc set to cut to a depth of about 4 inches while operating at 4-6 mph. For thorough mixing, disc in two directions (cross disc), and follow with a harrow, drag, or other leveling device. Prior to the second discing, readjust the disc to prevent cutting deeper than 4 inches. Cynex 4L may be applied preemergence as an overlay over previously incorporated SUTAN+ or ERADICANE. Any rotation crop may be planted in the fall or spring following these treatments.

Existing stands of quackgrass, purple and yellow nutsedge must be turned under and thoroughly chopped up prior to chemical treatments. Additional weeds controlled by SUTAN+ or ERADICANE combinations:

Grasses:

Sandbur  
Texas Panicum

Shattercane (Wild Cane)\*  
Quackgrass (ERADICANE only)  
Wild Proso Millet\* (ERADICANE only)

Perennial Weeds:

Yellow Nutsedge (Nutgrass)

Purple Nutsedge Nutgrass)

\*Suppression only - refer to SUTAN+ or ERADICANE label for appropriate supplemental cultural and tillage practices.

For fields with moderate to heavy infestations of these weeds refer to the SUTAN+ or ERADICANE label for appropriate higher rates.

**Cynex 4L plus DUAL 8E**

Use Cynex 4L at the proper rate for soil texture and organic matter shown in Table 4. Use DUAL as follows:

Soil Texture	Broadcast Rate Per Acre For DUAL
<b>Coarse</b> Sand, Loamy sand, Sandy loam	1.25 - 1.5 pints
<b>Medium</b> Loam, Silt loam, Silt	1.5 - 2.0 pints
<b>Fine</b> Sandy clay loam, Silty clay loam, Clay loam, Sandy clay, Silty clay, Clay	1.5 - 2.5 pints

The low end of the rate range should be used for lowest organic matter soils and the rate increased as organic matter increases to a point that soils containing 4 percent organic matter or more require the highest rate shown for that soil texture. Refer to the Dual label for precautions on rotational crops.

**Cynex 4L plus Atrazine plus LASSO, SUTAN+, ERADICANE, or DUAL**

Use Cynex 4L plus atrazine at the proper rate for soil texture and organic matter shown in Table 5. Use LASSO, SUTAN+, ERADICANE, or DUAL according to rates shown in "Cynex 4L Combinations" in this section of the label. Rotational Crops: Refer to Rotational Crops section of "Cynex 4L plus Atrazine" in this section of the label.

TABLE 4

**PREEMERGENCE BROADCAST APPLICATION RATES PER ACRE FOR Cynex 4L USED IN TANK-MIX COMBINATIONS WITH LASSO, SUTAN+, ERADICANE, OR DUAL ON CORN**

Quarts of Cynex 4L****						
Percent Organic Matter in Soil**						
Soil Texture Description	Less Than 1%	1%	2%	3%	4%	5% & Over
Sand, Loamy Sand	0.6***	0.75	1.25	1.5	1.75	2.0
Sandy Loam Loam, Silt Loam, Silt	0.75	1.25	1.5	1.75	2.0	2.25
Sandy Clay Loam	1.25	1.5	1.75	2.0	2.25	2.5
Clay Loam, Silty Clay Loam	1.5	1.75	2.0	2.25	2.5	2.75
Sandy Clay, Silty Clay, Clay	1.75	2.0	2.25	2.5	2.75	3.0
Peat or Muck	*	*	*	*	*	*

\*Not Recommended

\*\*For organic matter content between those listed, adjust the rate proportionately.

\*\*\*Do not use in the light sandy soils of the Atlantic Coastal Plain.

\*\*\*\*Maximum rate limit per acre per year for all applications is 6.5 lbs. cyanazine (6.5 quarts Cynex 4L) except on highly erodible land with less than 30% plant residue cover, the rate limit is 3.0 lbs. cyanazine (3.0 quarts Cynex 4L).

**TABLE 5  
PREEMERGENCE BROADCAST APPLICATION RATES PER ACRE FOR Cynex 4L PLUS ATRAZINE 4L USED IN TANK-MIX COMBINATIONS WITH LASSO, SUTAN+, ERADICANE, OR DUAL ON CORN**

Quarts of Cynex 4L + Quarts of Atrazine 4L***	
Percent Organic Matter in Soil**	

Soil Texture Description	Less Than 1%	1%	2%	3%	4%	5% & Over
Sand, Loamy Sand	0.4 + 0.2***	0.5 + 0.25	0.75 + 0.5	1.0 + 0.5	1.25 + 0.5	1.25 + 0.75
Sandy Loam	0.5 + 0.25	0.75 + 0.5	1.0 + 0.5	1.25 + 0.5	1.25 + 0.75	1.5 + 0.75
Loam, Silt Loam, Silt	0.75 + 0.5	1.0 + 0.5	1.25 + 0.5	1.25 + 0.75	1.5 + 0.75	1.75 + 0.75
Sandy Clay Loam, clay Loam, Silty Clay Loam	1.0 + 0.5	1.25 + 0.5	1.25 + 0.75	1.5 + 0.75	1.75 + 0.75	2.0 + 0.75
Sandy Clay, Silty Clay, Clay	1.25 + 0.5	1.25 + 0.75	1.5 + 0.75	1.75 + 0.75	2.0 + 0.75	2.0 + 1.0
Peat or Muck	*	*	*	*	*	*

\*Not Recommended

\*\*For organic matter content between those listed, adjust the rate proportionately.

\*\*\*Do not use in the light sandy soils of the Atlantic Coastal Plain.

\*\*\*\*If Atrazine 90% is used, multiply rates shown by 1.11 to equal quarts of Atrazine 4L. If Atrazine 80W is used, multiply rates shown by 1.25 to equal pounds of Atrazine 80W.

### CONSERVATION TILLAGE WEED CONTROL

#### Early Preplant (EPP)

Cynex 4L may be used for Early Preplant or Preemergence weed control for land going into production of corn under conservation tillage programs. Complete any planned early spring tillage prior to application. Apply herbicide treatment before weeds germinate or before weed seedlings are more than 3 inches tall. Tillage after application may reduce the effectiveness of the herbicide treatment. A nitrogen solution or complete fertilizer solution may replace all or part of the water as a carrier. The spray gallonage and spray boom design must be adequate to give thorough uniform coverage of the weed foliage. Follow label requirements of all products used in tank mix combinations.

**Cynex 4L alone or in combination with atrazine:** Apply 15 to 30 days prior to planting. Use the proper rate for soil texture and organic matter indicated in Table 1, 2 or 3. Where heavy crop residues exist, the rates shown in Tables 1, 2 or 3 should be increased by 25%.

**Cynex 4L plus PRINCEP 4L or PRINCEP CALIBER 90 or Cynex 4L plus Atrazine plus PRINCEP 4L or PRINCEP CALIBER 90:** Apply 30 days or more prior to planting. Use the proper rate of Cynex 4L or of Cynex 4L plus Atrazine for the soil texture and organic matter as shown in Table 1, 2 or 3 and add 1 quart/acre of PRINCEP 4L or 1.11 pounds/acre of PRINCEP CALIBER 90.

#### Burndown Of Existing Weeds:

**Combinations with 2,4-D:** Where broadleaf weeds are present at the time of application, add 2,4-D LV Ester at 1-1/3 - 2 pts./A. (6 lb./gal.) or 2-3 pts./A. (4 lbs./gal.) (or 2,4-D Amine at recommended rates) plus nonionic surfactant at 1 qt./100 gals. of diluted spray, or other suitable surfactant at its recommended rate.

**Combinations with Paraquat Products:** When grasses are present and/or when existing weeds exceed 3 inches in height add 1-2 pts./A. (2 lbs./gal.) or 1.3-2.7 pts./A. (1.5 lbs./gal.) of paraquat. Well established weeds 6 inches tall or taller may not be well controlled.

Apply Cynex 4L in at least 20 gal./A. of carrier by ground sprayer. (The volume of carrier and the application equipment must be adequate to give a uniform application.) Add Ortho. surfactant at 1-2 qts./100 gals. of diluted spray (or other suitable surfactant at recommended rates) where paraquat is used. Crop oil concentrate or emulsible vegetable oil are not needed where paraquat is used.

Depending upon weather conditions following the EPP application, a postemergence treatment of Cynex 4L or some other herbicide treatment may be necessary at or after planting to provide additional length of weed control. If desired, 1-1/2-2 pts./A. of DUAL 8E or 2 qts./A. of LASSO 4EC may be tank-mixed with the Cynex 4L EPP treatment or applied preemergence at planting.

**Rotational Crops:** Refer to Rotational Crops section for each treatment in the Preemergence section of the label.

### At Planting

Cynex 4L applied alone or in combination with atrazine and/or LASSO or DUAL according to the following directions will kill most existing small weeds and suppress many emerged perennial weeds when corn is planted into no-till stalk ground (corn, sorghum), stubble ground (soybean, small grains), and any minimum-till situation. This treatment then provides residual control of annual weeds as in conventional tillage.

Apply Cynex 4L alone or with other products according to the directions for those treatments in the Preemergence section of the label. Where heavy crop residues exist, the Cynex 4L rate shown in Tables 1 through 5 should be increased by 25%. Add 0.5-1.0 pt./A. of 2,4-D LV (6 lbs./gal.) or 0.75 to 1.5 pts./A. (4 lbs./gal.) (or 2,4-D Amine at recommended rates). Add the 2,4-D to the spray tank last.

Use a minimum of 15 gals./A. of carrier. Complete spray coverage of the weeds is essential for best performance. Nitrogen solutions and complete liquid fertilizers are the preferred carriers for this treatment because they aid in the burndown of existing weeds. Add nonionic surfactant at 1-2 qts. per 100 gals. of diluted spray, or other suitable surfactant at its recommended rate. If water is used as a carrier, crop oil concentrate may be used as a surfactant. Apply before weeds exceed 3 inches in height. For control of existing alfalfa add 1/3 - 1/2 pt./A. of BANVEL to the spray mixture. Apply before the alfalfa exceeds 6 inches in height.



For fields with existing sod grasses such as orchardgrass, bromegrass, rye or timothy, or when very dry conditions exist, or when existing weeds exceed 3 inches in height, add paraquat to the tank mix. Use 2 pts./A. (2 lbs./gal.) or 1.3 pts./A. (1.5 lbs./gal.) of paraquat in combination with Cynex 4L as described above in this section, except the 2,4-D may be omitted, if desired, and the gallonage should be increased to 20-40 gals./acre to give thorough coverage of the weed growth. Do not apply paraquat in suspension type liquid fertilizer.

### POSTEMERGENCE

Under warm, sunny, dry conditions of low humidity and the absence of dew formation at night, Cynex 4L may be applied preemergence on field corn only. Do not use a surfactant, crop oil, or any other adjuvant. Addition of a surfactant, EV oil, or fertilizer solution is not recommended because moderate to severe injury, including stand loss, may occur.

Do not apply this treatment under cold, wet weather conditions or to corn growing under stress caused by weather, insects, disease, etc. Yellowing of the corn, stunting or stand loss may result from this treatment, particularly if high rainfall or cold, adverse growing conditions occur after application. Do not apply postemergence on popcorn, sweet corn or corn grown for seed.

#### Cynex 4L Applied Alone

Use Cynex 4L at the proper rate for the soil texture and organic matter shown in Table 6 or 7. Use rates shown in Table 6 if Cynex 4L, CYANAZINE OR CYANAZINE/ATRAZINE herbicides have not been applied to the soil this season. Use rates shown in Table 7 if Cynex 4L, CYANAZINE OR CYANAZINE/ATRAZINE herbicides have been applied to the soil this season. This treatment may be used on peat or muck soils for burndown and suppression of existing weeds but will not provide residual control. Apply from crop emergence through the four-leaf stage of corn growth but before weeds exceed about 1 1/2 inches in height. Do not apply over the top of corn if the fifth leaf is visible. Apply in water only. Do not spray emerged corn plants with Cynex 4L in a liquid fertilizer carrier or in tank mix combinations with EC formulation herbicides.

Any rotational crop may be planted the fall or spring following this treatment.

**TABLE 6  
POSTEMERGENCE BROADCAST APPLICATION RATES PER ACRE FOR Cynex 4L  
ON FIELD CORN. NO PRIOR APPLICATION OF Cynex 4L, OR OTHER  
CYANAZINE OR CYANAZINE/ATRAZINE HERBICIDES**

Quarts of Cynex 4L
Percent Organic Matter in Soil*

Soil Texture	Less Than 1%	1%	2%	Over 2%
Sand, Loamy Sand	Do Not Use	1.1	1.5	2.0
Sandy Loam	1.1	1.5	2.0	2.0
Loam, Silt Loam, Silt	1.5	2.0	2.0	2.0
All Other Textures	2.0	2.0	2.0	2.0

\*For organic matter content between those listed, adjust the rate proportionately.

**TABLE 7**  
**POSTEMERGENCE BROADCAST APPLICATION RATES PER ACRE FOR Cynex 4L ON FIELD CORN. Cynex 4L, OR OTHER CYANAZINE OR CYANAZINE/ATRAZINE HERBICIDES USED IN PRIOR APPLICATION**

Quarts of Cynex 4L**				
Percent Organic Matter in Soil*				
Soil Texture	Less Than 1%	1%	2%	Over 2%
Sand, Loamy Sand	Do Not Use	Do Not Use	1.3	1.3
Sandy Loam	Do Not Use	Do Not Use	1.5	1.8
Loam, Silt Loam, Silt	Do Not Use	1.3	1.8	1.8
All Other Textures	Do Not Use	1.8	1.8	2.0

\*For organic matter content between those listed, adjust the rate proportionately.

\*\*Maximum rate limit per acre per year for all applications is 6.5 lbs. cyanazine (6.5 quarts Cynex 4L) except on highly erodible land with less than 30% plant residue cover, the rate limit is 3.0 cyanazine (3.0 quarts Cynex 4L).

**Cynex 4L COMBINATIONS**

) **Cynex 4L plus Atrazine**

Apply as directed in "Postemergence-Cynex 4L Applied Alone" section of this label. Use an

amount of Cynex 4L plus Atrazine equal to the rate shown in Table 6 or 7 for the proper soil texture and organic matter. To determine the amount of Cynex 4L to use, multiply the rate in Table 6 or 7 by 0.6. To determine the amount of Atrazine 4L to use, multiply the rate indicated in Table 6 or 7 by 0.4. (Multiply by 0.5 for pounds of Atrazine 80W.) (For the 2.0 qts/A. rate shown in Table 6 or 7, use 1.2 qts/A. of Cynex 4L plus 0.8 qts/A. of Atrazine 4L or 1.0 lbs/A. of Atrazine 80W.)

**Rotational crops:** See Rotational Crops section in the "Preemergence Cynex 4L plus Atrazine" section of this label.

### SWEET CORN

Cynex 4L may be applied preemergence or preplant incorporated for the control of annual grasses and broadleaved weeds in sweet corn.

**NOTE:** Cynex 4L may cause injury or stand loss on new or "super sweet" varieties of sweet corn. Consult with Agricultural Extension Agencies and sweetcorn seed suppliers about the sensitivity of new varieties to potential injury.

Apply Cynex 4L treatments just before, at or after planting but before crop has emerged. Avoid removal of treated soil from seedrow prior to or during the planting operation. Do not apply postemergence to sweet corn. Rotary hosing is recommended for preemergence application which do not receive adequate rainfall or sprinkler irrigation to wet the top 1 1/2 to 2 inches of soil within about 10 days after application. If a Cynex 4L mixture is to be incorporated, except as noted, single or two pass incorporation is acceptable. Care should be taken to incorporate the Cynex 4L mixture no deeper than the top two inches of soil.

**TABLE 8  
PREEMERGENCE BROADCAST APPLICATION RATES PER ACRE FOR  
TANK-MIX COMBINATIONS OF Cynex 4L PLUS ATRAZINE 4L ON SWEET CORN**

Quarts of Cynex 4L**** + Quarts of Atrazine 4L***						
Percent Organic Matter in Soil**						
Soil Texture Description	Less Than 1%	1%	2%	3%	4%	5% & Over
Sand, Loamy Sand	Do Not Use	0.8 + 0.4	1.1 + 0.4	1.3 + 0.7	1.5 + 0.9	2.2 + 1.1
Sandy Loam	Do Not Use	1.1 + 0.4	1.3 + 0.7	1.5 + 0.9	2.0 + 1.1	2.8 + 1.3

Loam, Silt Loam, Silt, Sandy Clay loam, Clay Loam	Do Not Use	1.3 + 0.7	1.5 + 0.9	2.0 + 1.1	2.5 + 1.3	3.2 + 1.3
Silty Clay Loam	Do Not Use	1.5 + 0.9	1.8 + 1.1	2.5 + 1.3	3.2 + 1.3	3.4 + 1.6
Sandy Clay, Silty Clay Loam	Do Not Use	1.8 + 1.1	2.8 + 1.3	3.2 + 1.3	3.4 + 1.6	3.4 + 1.8
Peat or Muck	*	*	*	*	*	*

\*Not Recommended

\*\*For organic matter content between those listed, adjust the rate proportionately.

\*\*\*If Atrazine 90% is used, multiply rates shown by 1.11 to equal lbs. of 90% Atrazine products. If Atrazine 80W is used multiply rates show by 1.25 equal lbs of Atrazine 80W.

\*\*\*\*Maximum rate limit per acre per year for all applications is 6.5 lbs. cyanazine (6.5 quarts Cynex 4L) except on highly erodible land with less than 30% plant residue cover, the rate limit is 3.0 lbs. cyanazine (3.0 quarts Cynex 4L).

### Cynex 4L PLUS ATRAZINE

Apply Cynex 4L at the proper rate for soil texture and organic matter indicated in Table 8. Table 8 provides rates for generally weedy conditions. The ratio of the amounts of each herbicide may be adjusted as necessary for particular weed conditions as long as the combined rate of the two products does not exceed the combined rate for the soil shown in Table 8. For grassier conditions use a ratio that contains higher levels of Cynex 4L (3:1). For fields with more broadleaves use a ratio that contains higher levels of atrazine (1:1).

**Rotational Crops:** (1) Plant only corn, sorghum or soybeans the year following the use of this mixture. (2) If soybeans are to be planted, injury may occur due to the carryover of Atrazine. (3) If applied after June 10, do not rotate with crops other than corn or sorghum the next year or injury may occur. (4) Small grains may be planted 15 months following treatments. (5) All other crops may be planted 18 months after application.

### Cynex 4L PLUS LASSO 4EC

Use Cynex 4L at the proper rate for the soil texture and organic matter shown in Table 9 plus 2 quarts per acre of LASSO. (Use 2.5 quarts LASSO on clay soils containing 5 percent organic matter and over.) Any rotational crop may be planted the fall or spring following this treatment.

### Cynex 4L PLUS SUTAN + 6.7E OR ERADICANE 6.7E

Use Cynex 4L at the proper rate for the soil texture and organic matter shown in Table 9 plus 1.8 quarts per acre of SUTAN+ or ERADICANE for control of many annual grasses and broadleaf weeds. (Use 2.4 quarts of SUTAN+ or ERADICANE on loam soils containing 5 percent or more organic matter, and clay loams and clays containing 4 percent or more organic matter.) Do not use on sands and loamy sands having less than 1 percent organic matter nor on the light sandy soils of eastern coastal states. Do not use on corn grown for seed.

Apply before planting. Incorporate the mixture immediately upon application using power-driven cultivation equipment set for 2-3 inch depth, or tandem disc set to cut about 4 inches deep while operating at 4-6 mph. For thorough mixing, disc in two directions (cross disc), and follow with a harrow, drag, or other leveling device. Prior to the second discing, readjust the disc to prevent cutting deeper than 4 inches. Cynex 4L may be applied preemergence as an overlay over previously incorporated SUTAN+ or ERADICANE, if desired. Any rotation crop may be planted in the fall or spring following these treatments.

Existing stands of quackgrass, purple and yellow nutsedge must be turned under and thoroughly chopped up prior to chemical treatments.

Additional weeds controlled by SUTAN+ or ERADICANE combinations:

**Grasses:**

- Sandbur, Shattercane (Wild Cane)\*
- Texas Panicum, Quackgrass (ERADICANE Only)
- Wild Proso Millet\* (ERADICANE only)

**Perennial Weeds**

- Yellow Nutsedge (Nutgrass), Purple Nutsedge (Nutgrass)

\*Suppression only - refer to SUTAN+ or ERADICANE label for appropriate supplemental cultural and tillage practices.

For fields with moderate to heavy infestations of these weeds refer to the SUTAN+ or ERADICANE labels for appropriate higher rates.

**Cynex 4L PLUS "DUAL" 8E**

Use Cynex 4L at the proper rate for soil texture and organic matter shown in Table 9. Use DUAL as follows:

Soil Texture	Broadcast Rate Per Acre for DUAL
Coarse Sand, Loamy sand, Sandy loam	1.25 - 1.5 pints
Medium	

Loam, Silt loam, Silt

1.5 - 2.0 pints

**Fine**

Sandy clay loam, Silty clay loam, Clay loam, Sandy clay, Silty clay, Clay

1.5 - 2.5 pints

The low end of the rate range should be used for lowest organic matter soils and the rate increased proportionately as the organic matter increases. Soils containing 4 percent organic matter or more require the highest rate shown for that soil texture. Refer to the DUAL label for precautions or rotational crops.

**TABLE 9  
PREEMERGENCE BROADCAST APPLICATION RATES PER ACRE FOR Cynex 4L  
USED IN TANK MIX COMBINATIONS WITH LASSO, SUTAN+, ERADICANE, OR  
DUAL ON SWEET CORN**

Quarts of Cynex 4L						
Percent Organic Matter in Soil**						
Soil Texture Description	Less than 1%	1%	2%	3%	4%	5% & Over
Sand, Loamy Sand	Do Not Use	0.8	1.2	1.4	1.6	2.0
Sandy Loam	Do Not Use	1.2	1.4	1.6	2.0	2.2
Loam, Silt Loam, Silty Loam, Sandy Clay Loam	Do Not Use	1.4	1.6	2.0	2.2	2.6
Clay Loam, Silty Clay Loam	Do Not Use	1.8	2.0	2.2	2.6	2.8
Sandy Clay, Silty Clay Loam	Do Not Use	2.0	2.2	2.6	2.8	3.0
Peat or Muck	*	*	*	*	*	*

\*Not Recommended

\*\*For organic matter content between those listed, adjust the rate proportionately.

\*\*\*Do not use in the light sandy soils of the Atlantic Coastal Plain.

**TABLE 10  
PREEMERGENCE BROADCAST APPLICATION RATES PER ACRE FOR Cynex 4L  
PLUS ATRAZINE 4L USED IN TANK-MIX COMBINATIONS WITH LASSO,  
SUTAN+,  
ERADICANE, OR DUAL ON SWEET CORN**

Quarts of Cynex 4L + Quarts of Atrazine 4L***						
Percent Organic Matter in Soil**						
Soil Texture Description	Less Than 1%	1%	2%	3%	4%	5% & Over
Sand, Loamy Sand	Do Not Use	0.6 + 0.2	0.9 + 0.4	1.0 + 0.5	1.0 + 0.6	1.4 + 0.6
Sandy Loam	Do Not Use	0.8 + 0.4	1.0 + 0.5	1.1 + 0.6	1.4 + 0.6	1.6 + 0.6
Loam, Silt Loam, Silt, Sandy Clay Loam	Do Not Use	1.0 + 0.5	1.2 + 0.6	1.4 + 0.6	1.6 + 0.6	1.8 + 0.9
Clay Loam, Silty Clay Loam	Do Not Use	1.2 + 0.6	1.4 + 0.6	1.6 + 0.6	1.8 + 0.9	2.0 + 0.9
Sandy Clay, Silty Clay Loam	Do Not Use	1.4 + 0.6	1.6 + 0.9	1.8 + 0.9	2.0 + 0.9	2.0 + 1.1
Peat or Muck	*	*	*	*	*	*

\*Not Recommended

\*\*For organic matter content between those listed, adjust the rate proportionately.

\*\*\*If a 90% Atrazine is used multiply Atrazine rates shown by 1.11 to equal lbs. of 90% Atrazine products. If Atrazine 80W is used, multiply rates shown by 1.25 to equal lbs. of Atrazine 80W.

**Cynex 4L PLUS ATRAZINE PLUS LASSO, SUTAN+, ERADICANE OR DUAL**

**NOTE:** Do not use combinations with SUTAN+ or ERADICANE in New Jersey. Use Cynex 4L plus Atrazine at the proper rate of soil texture and organic matter shown in Table 10. Use LASSO, SUTAN+, ERADICANE or DUAL according to rates shown in Cynex 4L combinations in this section of the label.

**Rotational Crops:** Refer to Rotational Crops section of Cynex 4L plus Atrazine in this section of the label.

## COTTON

### IDLE SEASON EARLY PREPLANT WEED CONTROL (CALIFORNIA ONLY)

#### Weeds Controlled

Annual Bluegrass	Rabbitsfoot Grass
Annual Ryegrass	Volunteer Small Grains
Barnyardgrass*	(suppression)
Bristly Foxtail	Wild Oat*
	Yellow Foxtail

#### Broadleaves

Annual henbit	Miners lettuce
Black nightshade	Lambsquarters
Burclover	London rocket
Cheeseweed*	Pineapple weed
Chickweed	Prickly lettuce
Fiddleneck	Shepherdspurse
Groundsel	Sowthistle
Knotweed	Wild mustard
Marestail	Wild radish

\*Under soil moisture conditions favoring deep germination, these species may not be completely controlled.

Cynex 4L may be used for burndown of small existing annual weeds and residual control of weeds during the winter and early spring season prior to planting cotton in California only. Complete any planned tillage prior to application. Apply herbicide treatment before weeds germinate or before weed seedlings are more than 3 inches tall. Tillage after application may reduce the effectiveness of the herbicide treatment.

Apply Cynex 4L at least 30 days prior to planting. Apply the proper rate for the soil texture, organic matter and time interval between application and planting indicated in the Table 11. Where existing weeds are present, add crop oil concentrate, surfactant, or emulsible vegetable oil at its recommended rate to aid in the burndown of small weeds.

Where existing weeds are greater than 3 inches in height, when very dry conditions exist or where volunteer grains are a major problem, tank-mix Cynex 4L with 1-2 pts./A. (2 lbs./gal.) or 1.3-2.7 pts./A. (1.5 lbs./gal. of paraquat. Well established weeds 6 inches tall or taller may not be well controlled.

Apply Cynex 4L in at least 20 gal./A. of carrier by ground sprayer. (The volume of carrier and the application equipment must be adequate to give a uniform application.) Add nonionic surfactant at 1-2 qts./100 gals. of diluted spray (or other suitable surfactant at recommended rates) where paraquat is used. Crop oil concentrate or emulsible vegetable oil are not needed where paraquat is used.



Cynex 4L can also be tank-mixed with TREFLAN or PROWL and incorporated for fall listed cotton beds instead of surface applied as described above. Precautions: 1). Failure to wait the recommended time interval between application and planting may result in crop injury. 2). At least one inch of rainfall or an equivalent irrigation that waters the surface of the soil after application must precede planting. 3). The use of this treatment on calcareous or caliche soil outcroppings may result in crop injury. 4). Do not graze or feed foliage from treated areas to livestock. 5). Do not apply Cynex 4L to cotton land in irrigation water.

**TABLE 11  
BROADCAST APPLICATION RATES PER ACRE OF Cynex 4L FOR IDLE SEASON  
OR EARLY PREPLANT TREATMENT ON COTTON**

Days Prior to Planting*						
30 Days		60 Days			90 Days	
Qts. of Cynex 4L**		Qts of Cynex 4L**			Qts of Cynex 4L*	
Soil Texture Description	Organic Matter Under 2%	Organic Matter Over 2%	Organic Matter Under 2%	Organic Matter Over 2%	Organic Matter Under 2%	Organic Matter Over 2%
Sands, Loamy Sands	1.5	2.0	2.5	3.0	3.0	3.5
All Other Soils	2.0	2.5	3.0	3.5	3.5	4.0

\*For time intervals between those listed, adjust rates proportionately.

\*\*Maximum rate limit per acre per year for all applications is 6.5 lbs. cyanazine (6.5 quarts Cynex 4L) except on highly erodible land with less than 30% plant residue cover, the rate limit is 3.0 lbs. cyanazine (3.0 quarts Cynex 4L).

**PREEMERGENCE**

Apply preemergence on cotton only in the States of Alabama, Arkansas, Louisiana and Mississippi.

**Weeds Controlled**

- Annual morningglory
- Prickly sida (Teaweed)
- Cocklebur
- Spurge

Cynex 4L is a selective preemergence herbicide for early season weed control in cotton. Supplemental practices (such as Cynex 4L applied directed postemergence) may be necessary to control late season weeds. Cynex 4L can be used in a tank-mix combination with Zorial Rapid 80.

Carefully match the Cynex 4L rate with the soil texture. Do not use on fields where the soil texture changes from coarse to fine. Avoid overlapping the spray pattern or overdosing the field with Cynex 4L. Application rates above those recommended for the soil texture can result in yellowing or stunting of the crop and may result in stand reduction.

While cotton exhibits tolerance to Cynex 4L, adverse growing conditions such as excessive rains, standing water or cold weather may result in stand reduction.

**DO NOT GRAZE OR FEED FOLIAGE FROM TREATED AREAS TO LIVESTOCK.**

**Cynex 4L plus ZORIAL RAPID 80.**

Apply Cynex 4L plus ZORIAL RAPID 80 at the proper rate for the soil texture shown in Table 12. The soil must contain at least 1.0 percent organic matter. Seed placement should be 1/2- 3/4 inch from the soil surface. Plant only cotton within six months after the last application of ZORIAL RAPID 80 or injury may occur.

**DIRECTED POSTEMERGENCE - LAYBY**

**Weeds Controlled**

Annual morningglory*	Palmer amaranth
Bristly starbur	Pigweed (redroot and spiny)
Cocklebur	Prickly sida (Teaweed)
Crotalaria	Sicklepod
Jimsonweed	Spurge
Lambsquarters	Tropic croton
Nightshade (annual)	Wright groundcherry

\*The degree of preemergence control from a layby treatment will be reduced if soil moisture and temperature conditions cause deep germination of the seed.

**TABLE 12  
PREEMERGENCE BROADCAST APPLICATION RATES PER ACRE FOR Cynex 4L  
+ ZORIAL RAPID 80 ON COTTON**

Soil Texture*	Quarts/Acre Cynex 4L 90DF	+	Quarts/Acre ZORIAL RAPID 80
Coarse Soils**			
Sandy loam	0.5	+	0.8
Medium Soils	0.6	+	1.3
Silt and Silt loam			
Loam, Clay loam, Sandy			

clay loam, Sandy clay      0.9                      +                      1.3

**Fine Soils**

Silty clay loam, Silty clay, Clay                      1.2                      +                      1.6

\*The soil must contain at least 1% organic matter.

\*\*Do not use on coarse soils (Sands and Loamy sands) containing more than 70% sand.

**TABLE 13  
DIRECTED POSTEMERGENCE  
APPLICATION RATES PER ACRE FOR Cynex 4L ON COTTON**

BANDED 38" ROW		
BROADCAST	12" BAND	19" BAND
0.6 - 1.0 qt.	0.2 - 0.3 qt	.0.3 - 0.5 qt.

Use the maximum rate when dry or arid conditions exist.

**TABLE 14  
LAYBY APPLICATION RATES PER ACRE FOR Cynex 4L ON COTTON**

Height of Cotton	Soil Texture		Broadcast Rates
12 inches or more	Coarse	Sandy Loam	0.8 qts
	Medium	Silt, Silt Loam, Loam, Clay Loam, Sandy ClayLoam, Sandy Clay	1.2 qts
	Fine	Silty Clay Loam, Silty Clay and Clay	1.6qts

**TABLE 15  
DIRECTED POSTEMERGENCE APPLICATION RATES PER ACRE Cynex 4L +  
MSMA ON COTTON**

Banded 38" Row			
Product	Broadcast	12" Band	19" Band

Cynex 4L + MSMA (4 lb/gal) or MSMA ( 6.6 lb/gal)	0.6 - 1.0 qt. + 4 pints or 2.4 pints	0.2 - 0.3 qt. + 1.3 pints or 0.8 pint	0.3 - 0.5 qt. + 2 pints or 1.2 pints
--	--	---	--

Cynex 4L and tank-mix combinations may be applied directed postemergence or layby to cotton and either preemergence to weeds or postemergence to weeds in all cotton growing States.

Apply before weeds are more than 2 inches tall. Apply the directed postemergence treatment after the cotton has attained the minimum height of 6 inches. For layby treatment, apply Cynex 4L after the cotton has attained the height of 12 inches or more.

The spray mixture should be directed to the soil around the base of the cotton plants. Care should be taken to prevent the spray from striking the cotton leaves as injury will occur. The use of leaf lifters or shields on application equipment is recommended to avoid spraying the cotton foliage.

Cynex 4L may be applied directed postemergence and/or layby following a preemergence application of Cynex 4L. Apply no more than two directed postemergence and one preemergence application to the same crop in any one year. If Cynex 4L is not used preemergence, apply no more than three directed postemergence applications including layby to the same crop in any one year. (In California, apply no more than two directed postemergence applications including layby.)

When applied as a layby treatment before weeds emerge, the effectiveness of Cynex 4L depends on rainfall or irrigation to move it into the soil. When irrigation water activation is used, every row must be watered and for skip row cotton all treated soil must be irrigated.

Any rotational crop may be planted the fall or spring following any of the treatments in this section providing the soil is plowed or deepdisced prior to planting the rotation crop.

Do not apply Cynex 4L to cotton in irrigation water.

**DO NOT GRAZE OR FEED FOLIAGE FROM TREATED AREAS TO LIVESTOCK.**

**Cynex 4L Applied Alone**

Apply Cynex 4L directed-postemergence at the rate shown in Table 13. Apply at layby at the rates for the soil texture indicated in Table 14. Add a nonionic agricultural surfactant suitable for use on growing cotton at the rate of 2 quarts per 100 gallons of spray mixture (or as directed by the manufacturer).

**Cynex 4L plus MSMA**

Apply a tank-mix combination of Cynex 4L plus MSMA plus surfactant after the cotton is 6 inches tall but before it reaches the bloom stage. Apply no more than two applications of this mixture before the first bloom stage. Tank-mix Cynex 4L plus MSMA at the rates indicated in Table 15. Add a nonionic surfactant at the rate of 2 qts./100 gals of spray mixture (or as directed by the manufacturer).

### GRAIN SORGHUM (MILO)

(See the **GENERAL INFORMATION** section of this label)

Tank-mix combinations of Cynex 4L plus atrazine, metolachlor (Dual), alachlor (Lasso), propazine (Milogard) or propachlor (Ramrod) may be used for selective preemergence weed control in grain sorghum. In addition, Cynex 4L may be used in tank-mix combinations with atrazine, DUAL or LASSO for the control of weeds in early spring, early preplant, 14 to 35 days or more prior to planting grain sorghum. Cynex 4L in tank mix combinations may be applied preemergence or early preplant on grain sorghum grown under conventional or conservation tillage systems.

Do not use on forage sorghums. See the appropriate sections of the label for geographic distribution restrictions.

### WEEDS CONTROLLED BY Cynex 4L IN TANK-MIX OR SEQUENTIAL COMBINATIONS WITH OTHER HERBICIDES ON GRAIN SORGHUM (MILO)

---

#### Grasses

Cheatgrass	Stinkgrass
Green foxtail	(Indian lovegrass)
Crabgrass	Volunteer Wheat (2)
Downy brome	Yellow foxtail

#### Broadleaves

Annual Morningglory	Lambsquarters	Russian thistle
Carpetweed	Pennsylvania smartweed	Shepherdspurse
Cocklebur (1)	Prostrate pigweed	Sunflower (1)
Common purslane	Prickly lettuce	Tansy and other
Horseweed (maretail)	Ragweed (Common)	mustards
Kochia	Redroot pigweed	Velvetleaf

---

(1) Under soil moisture and temperature conditions favoring deep germination or other factors that may cause delayed germination, these species may not be completely controlled.

(2) When the herbicide treatment is applied two weeks or more before planting, weed control of these species may break early if heavy rainfall occurs between application and planting.

### General Directions For Use On Grain Sorghum

Cynex 4L used in tank-mixes with the products listed above should be applied only once per crop season or in split dosage treatments. If replanting of grain sorghum is necessary, it may

be planted in soil previously treated with these mixtures. Apply Cynex 4L in these tankmix combinations before the crop has emerged.

**Precautions:** Do not make an additional application of Cynex 4L or any product containing cyanazine or crop injury may occur. Heavy rainfall between planting and crop emergence may cause crop injury or stand loss. Rainfall tends to cause excessive concentrations of herbicide in seed furrow, resulting in possible crop injury. Level deep planter marks or seed furrows before application. Do not apply to furrowplanted sorghum.

Sorghum growing under stress caused by minor element deficiency, cold, wet weather or sorghum growing on highly calcareous soil (high pH) may suffer in injury including stand reduction. Sorghum subjected to high winds, sand cutting, hail damage, or cold temperatures may be more susceptible to injury from the chemical treatment with possible stand loss. Where crop residues are pressed into the planter slot or any other factor keeps the slot from closing, crop injury or stand reduction may occur, caused by herbicides coming into direct contact with the seed from the spray or by being washed into the slot after a heavy rain.

When using DUAL on sorghum, only CONCEP II safened seed may be used. When using LASSO on sorghum, only SCREEN safened seed may be used.

Early preplant treatments will require a preemergence herbicide treatment other than Cynex 4L 9ODF or 4L at planting, and/or postemergence herbicide in the growing crop to provide required weed control, if the early preplant application is made more than 35 days prior to planting, or if weeds are present at planting time.

### WHEAT/SORGHUM/FALLOW ROTATION

For sorghum grown under a wheat/sorghum/fallow rotation, Cynex 4L plus atrazine may be applied early preplant or preemergence. If an early preplant application with Cynex 4L plus atrazine is used, it should be used in conjunction with either a residual herbicide after wheat harvest the previous year, a preemergence herbicide treatment at planting, and/or a postemergence herbicide treatment in the growing crop. The total rate of atrazine permitted in conjunction with a sorghum crop (postharvest plus early preplant plus preemergence) is limited to 3.0 lbs. active ingredient per acre. Even this or lower rates may carry over to injure rotational crops.

**Rotational Crops:** (1) Plant only corn, sorghum, or soybeans the year following the use of this mixture. (2) If soybeans are to be planted, injury may occur. (3) If applied after June 10, do not rotate with crops other than corn or sorghum the next year or injury may occur. (4) In the high plains and intermountain areas of the West where rainfall is sparse and erratic or where irrigation is required, use only when corn or sorghum is to be planted the following year, or a crop of corn or sorghum not treated with this mixture or atrazine is to precede other rotational crops. (5) Small grains may be planted 1. months following treatment. (6) All other crops may be planted 18 months after application.

### Conservation Tillage

19, 1/0

(

For grain sorghum grown under conservation tillage, any of the herbicide treatments listed above may be tank-mixed with paraquat, and/or 2,4-D where weeds and grasses are present at the time of application exceeding two inches in height.

If volunteer wheat or cheatgrass is over 2 inches tall, heavily tillered, and/or growing in a dense mat, or if the wheat stubble has been tilled (undercut or disc) or grazed and seed is buried, complete control may not be achieved. Control of volunteer and other weeds is usually better in complete no-till situations than were prior tillage "planted" the seed and allowed extensive root development.

Complete spray coverage of the weeds is essential for best performance. Apply the desired treatment in 15 to 30 gallons of spray mixture per acre by ground rig. When using paraquat in a tank-mix, apply the desired rates in 20 to 40 gallons of spray mixture per acre. Use the highest volumes where there are heavy crop residues on the soil surface. Nitrogen solutions are the preferred carriers for these treatments as they aid in the burndown of existing weeds. In addition, crop oil or a nonionic surfactant may be added to the tank-mix as they aid in the burndown of existing weeds.

(

When tank-mixing with 2,4-D to control broadleaf weeds 14-35 days prior to planting, use 1 to 3 pints per acre of 4 pounds ai per gallon, 2,4-D LV (2/3 to 2 pints of 6 pounds ai per gallon 2,4-D LV) (or 2,4-D Amine at recommended rates). Use the higher rates where overwintering weeds are present or when directed on the 2,4-D label for the control of specific hard-to-kill weed species, such as perennials.

(

**CAUTION:** Use only these 2,4-D product with properly registered labels that permit such use and application rates.

When tank-mixing with paraquat to control grass and broadleaf weeds at the time of planting, use an approved nonionic adjuvant at the rate of 1 quart per 100 gallons of dilute spray. Use 1-2 pints of paraquat (2 lbs./gal) or 1.3-2.7 pints (1.5 lbs/gal) per acre. Use the higher rate when weed growth is heavy or over 4 inches tall or when dry weather conditions prevail. Established weeds 6 inches tall or taller may not be completely controlled with paraquat.

OBSERVE ALL CAUTIONS AND LIMITATIONS ON LABELING OF ALL PRODUCTS USED IN MIXTURES.

### EARLY PREPLANT

Cynex 4L in tank-mix combinations with other herbicides may be applied Early Preplant on grain sorghum only in the states of Kansas, Nebraska and South Dakota. The tank-mixes may be applied on grain sorghum grown under conventional or conservation tillage systems. Do not use on forage sorghum.

TABLE 16  
EARLY PREPLANT BROADCAST APPLICATION RATES OF Cynex 4L PLUS  
ATRAZINE 4L ON GRAIN SORGHUM (0.8% TO 3% ORGANIC MATTER SOILS)

Quarts of Cynex 4L + Quarts of Atrazine 4L			
Days Prior to Planting (b, c)			
Soil Texture Description	14 Days	28 Days	35 Days
Sand, Loamy Sand	Do Not Use	Do Not Use	Do Not Use
Sandy Loam	1.2 + 0.6(a,d)	1.6 + 0.8	2.0 + 1.0
Loam, Silty Loam, Silt	1.5 + 0.7	2.0 + 1.0	2.3 + 1.2
Sandy Clay Loam, Clay Loam, Silty Clay Loam	2.0 + 1.0	2.3 + 1.2	2.6 + 1.4
Sandy Clay, Silty Clay, Clay	2.2 + 1.1	2.3 + 1.2	3.0 + 1.5
Peat or Muck	*	*	*
Eroded Slopes or Knobs Soils with pH greater than 8.0	*	*	*

\*Not Recommended

- (a) The first number is quarts/acre of Cynex 4L. The second Number is quarts/acre of atrazine 4L. For pounds/acre of atrazine 80W multiply by 1.25.
- (b) For intervals longer than 35 days due to delayed planting, etc., another herbicide treatment is needed before, at, or after planting.
- (c) For sandy loam soils containing less than 1.5% O.M., increase the treatment interval from 14 to 21 days. Use the 14 day interval for sandy soils containing more than 1.5%.
- (d) For rates between those listed at 14 and 35 days, adjust the rate proportionately.

**TABLE 17**  
**EARLY PREPLANT BROADCAST APPLICATION RATES PER ACRE IN QUARTS OF Cynex 4L, QUARTS OF ATRAZINE 4L AND PINTS OF DUAL 8E ON GRAIN SORGHUM (b)**

Soil Texture Description	Days Prior to Planting (c)	
	(0.8% to 3% Organic Matter)	
	14 Days**	28 Days
Dual	Qts Cynex 4L/Qts Atraz/Pts Dual	Qts Cynex 4L/qts Atraz/Pts



Sand, Loamy Sand	Do Not Use	Do Not Use	Do Not Use	1.0(a)	0.5	1.8
Sandy Loam	0.9(d)	0.4	1.35	1.1	0.5	1.8
Loam, silt Loam, Silt	1.1	0.5	1.35	1.35	0.7	1.8
Sandy clay Loam, clay Loam, Silty Clay Loam	1.35	0.7	1.6	1.7	0.8	1.8
Sandy Clay, silty clay, Clay	1.5	0.75	1.6	1.8	0.9	1.8
Peat or Muck	*	*	*	*	*	*
Eroded Slopes or Knobs Soils with pH Greater than 8.0	*	*	*	*	*	*

\*Not Recommended

(a) The first number is quarts/acre of Cynex 4L. The second number is quarts/acre of atrazine 4L. The third number is pints/acre of Dual 8E.

(b) If using Dual 8E on sorghum, Concep II safened seed should be planted.

(c) For intervals longer than 28 days due to delayed planting, etc., another herbicide treatment is needed before, at, or after planting.

(d) For sandy loam soils containing less than 1.5% O.M., increase the treatment interval from 14 to 21 days. Use the 14 day interval for sandy soils containing more than 1.5%.

\*\*For rates between those listed at 14 and 28 days, adjust the rate proportionately.

**TABLE 18**  
**EARLY PREPLANT BROADCAST APPLICATION RATES PER ACRE IN QUARTS**  
**OF Cynex 4L, QUARTS OF ATRAZINE 4L AND QUARTS OF LASSO 4EC ON**  
**GRAIN SORGHUM (b)**

Days Prior to Planting (c)

10/1/10

(0.8% to 3% Organic Matter)

Soil Texture  
Description  
Lasso

14 Days\*\*

28 Days

Qts Cynex 4L/Qts Atraz/Qts Lasso

Qts Cynex 4L/Qts Atraz/Qts

Sand, Loamy Sand	Do Not Use	Do Not Use	Do Not Use	0.9 (a)	0.45	2.25
Sandy Loam	0.7(d)	0.4	1.8	0.9	0.5	2.25
Loam, Silt Loam, Silt	0.9	0.5	1.8	1.2	0.6	2.25
Sandy Clay Loam, Clay Loam, Silty Clay Loam	1.2	0.6	2.0	1.5	0.75	2.25
Sandy Clay, Silty Clay, Clay	1.35	0.7	2.0	1.7	0.8	2.25
Peat or Muck	*	*	*	*	*	*
Eroded Slopes or Knobs Soils with pH greater than 8.0	*	*	*	*	*	*

\*Not Recommended

(a) The first number is quarts/acre of Cynex 4L. The Second number is quarts/acre of atrazine 4L. The third number is quarts /acre of Lasso 4EC.

(b) If using Lasso 4EC on sorghum, SCREEN safened seed should be planted.

(c) For intervals longer than 28 days due to delayed planting, etc., another herbicide treatment is needed before, at, or after planting.

(d) For sandy loam soils containing less than 1.5% O.M., increase the treatment interval from 14 to 21 days. Use the 14 day interval for sandy soils containing more than 1.5%

\*\*For rates between those listed at 14 and 28 days, adjust the rate proportionately.

## Cynex 4L PLUS ATRAZINE

Cynex 4L plus atrazine provides control of weeds listed on this label in the Weeds Controlled section for Cynex 4L on grain sorghum. Precautions: Crop injury can occur if the soil stays dry between application and slanting. Delay planting until at least 10 days after a soaking rain after treatment (soaks the soil to a depth of 4 inches or more). Heavy rains between planting and emergence can cause crop injury or sand loss. Rotational Crops: Refer to the General Information section of the label. Fields treated with this tank-mix may be planted only to corn or grain sorghum within 12 months after this treatment. Other crops should not be planted for 18 months following this treatment.

Use Cynex 4L plus atrazine 4L at the proper rate for soil texture and time interval indicated in Table 16.

**Split Applications:** Cynex 4L plus Atrazine 4L dosage rates in Table 16 may be applied in a split application of 80% of the dosage 21 days or more before planting and the remaining 20% at planting time. If the season has been dry without sufficient rainfall to activate the herbicide after the initial application, the remaining 20% at planting time may be omitted to reduce the chances of crop injury.

## Cynex 4L IN THREE-WAY COMBINATIONS

Cynex 4L plus atrazine in combination with DUAL or LASSO applied early preplant provides control of weeds listed on this label in the Weeds Controlled section for Cynex 4L on grain sorghum. Additional weeds controlled by one of these combinations include:

### Grasses

Barnyardgrass (1)	Stinkgrass (2)
Fall Panicum	Witchgrass (2)
Giant Foxtail	

(1) (2) Refer to Weeds Controlled section on grain sorghum for explanation.

**Rotational Crops:** If the crop treated with any of these combinations is lost, corn or grain sorghum may be replanted immediately without retreatment. Refer to the General Information section of the label for additional information regarding crop rotation.

## Cynex 4L PLUS ATRAZINE PLUS DUAL

Dual may be used on grain sorghum only with seed treated with CONCEP II seed safener.

Use Cynex 4L plus atrazine 4L plus DUAL 8E at the proper rate for soil texture and time interval indicated in Table 17.

## Cynex 4L PLUS ATRAZINE PLUS LASSO

LASSO may be used on grain sorghum only with seed treated with SCREEN seed safener.

Use Cynex 4L plus Atrazine 4L plus LASSO 4EC at the proper rate for soil texture and time interval indicated in Table 18.

## PREEMERGENCE OR SHALLOW PREPLANT INCORPORATION

Tank-mix combinations of Cynex 4L plus atrazine, metolachlor (Dual), alachlor (Lasso), or propachlor (Ramrod) may be used for selective preemergence weed control in grain sorghum. Cynex 4L and its tank mixes may be applied preemergence on grain sorghum grown under conventional or conservation tillage systems. Do not use on forage sorghum. Cynex 4L in two-way tank-mix combinations with propachlor or propazine or in three-way combinations with atrazine and propachlor, LASSO or DUAL should be applied only once per crop season or in split dosage treatments. (See the Early Preplant section.) If replanting of grain sorghum is necessary, it may be planted in soil previously treated with these mixtures. Do not make a second application of Cynex 4L or any product containing cyanazine or crop injury may occur.

Apply these tank-mix combinations at planting or after planting, but before the crop and weeds have emerged. Heavy rain immediately following application tends to cause excessive concentrations of herbicide in seed furrow and can cause crop injury and stand reduction. Do not apply to furrow-planted sorghum until furrows are leveled (plowed-in). Level deep planter marks or seed furrows before application. Sorghum growing under stress caused by minor element deficiency or cold, wet weather or sorghum growing on highly calcareous soil may suffer injury and stand loss.

### Cynex 4L plus Propachlor (RAMROD)

#### Weeds Controlled

Apply this tank-mix to grain sorghum only in States East of the Rocky Mountains.

#### Grasses

Barnyardgrass	Giant foxtail
Crabgrass	Green foxtail
Fall panicum	Yellow foxtail

#### Broadleaves

Annual morningglory	Pigweed
Charpetweed	Ragweed (Common)
Locklebur*	Smartweed
Common	purslane(Pennsylvania)
Lambsquarters	Velvetleaf*

\*Under conditions such as low temperatures, lack of soil surface moisture or other factors causing a delay in germination of the seeds, the degree of control may be impaired against these weeds.

Apply Cynex 4L plus propachlor at the proper rate for soil texture and organic matter shown in Table 19. Any rotational crop may be planted the fall or spring following this treatment.

**TABLE 19**  
**PREEMERGENCE BROADCAST-APPLICATION RATES IN QUARTS OF Cynex 4L PLUS**  
**QUARTS OF PROPACHLOR (RAMROD) 4L PER ACRE ON GRAIN SORGHUM**  
**PERCENT ORGANIC MATTER IN THE SOIL\*\***

Soil Texture Description	2% Qts Cynex 4L +	Qts Ramrod	3% Qts Cynex 4L +	Qts Ramrod

Sand, Loamy Sand	Do Not Use	Do Not Use	Do Not Use	Do Not Use
Sandy Loam	1.0(a)	2.5	1.2	3.0
Loam, Silt Loam, Silt	1.2	3.0	1.35	3.5
Sandy Clay Loam Clay Loam, Silty Clay Loam	1.35	3.5	1.6	4.0
Sandy Clay, Silty Clay, Clay	1.6	4.0	1.6	4.0
Peat or Muck	*	*	*	*
Eroded Slopers or Knobs soils with pH greater than 8.0	*	*	*	*

(a)The first number is quarts/acre of Cynex 4L. The second number is quarts/acre of RAMROD 4L.  
 \*\*For organic matter content between those listed, adjust the rate proportionately.  
 \*Not Recommended

**TABLE 20**  
**PREEMERGENCE BROADCAST APPLICATION RATES IN QUARTS OF Cynex 4L, QUARTS OF ATRAZINE 4L PLUS PINTS OF DUAL 8E PER ACRE ON GRAIN SORGHUM (a,b)**

**PERCENT ORGANIC MATTER SOIL\*\***

Soil Texture Description	Product		1%	2%	3%
Sand, Loamy Sand, Sandy Loam	Do Not Use	Do Not Use	Do Not Use	Do Not Use	Do Not Use
Loam, Silty Loam, Silt	Cynex 4L Atrazine 4L Dual 8E	Quarts	0.66 0.3 1.5	0.7 0.4 1.5	0.8 0.4 1.5
Sandy Clay Loam, Clay Loam, Silty Clay Loam	Cynex 4L Atrazine 4L Dual 8E	Quarts	0.7 0.4 1.5	0.8 0.4 1.75	0.9 0.5 2.0

Sandy Clay, Silty Clay, Clay	Cynex 4L	Quarts	0.8	0.9	0.9
	Atrazine 4L		0.4	0.5	0.5
	Dual 8E		1.5	1.75	2.0
Peat or Muck	*	*	*	*	*
Eroded Slopes or Knobs Soils with pH greater than 8.0	*	*	*	*	*

\*Not Recommended

(a)When using DUAL 8E on sorghum, only CONCEP II safened seed should be planted.

(b)To enhance weed control in areas of less than 25 inches of annual precipitation or where long dry periods are common, this treatment may require shallow incorporation with a tool such as a field cultivator operated at 5-7 mph.

\*\*For organic matter content between those listed, adjust the rate proportionately.

TABLE 21

**PREEMERGENCE OR SHALLOW PREPLANT INCORPORATION BROADCAST APPLICATION RATES IN QUARTS OF Cynex 4L, QUARTS OF ATRAZINE 4L AND QUARTS OF LASSO EC PER ACRE ON GRAIN SORGHUM (a,b)**

**PERCENT ORGANIC MATTER IN SOIL\*\***

Soil Texture Description	Product		1%	2%	3%
Sand, Loamy Sand	DO NOT USE	DO NOT USE	DO NOT USE	DO NOT USE	DO NOT USE
Sandy Loam	Cynex 4L Atrazine 4L Lasso 4EC	Quarts	none none none	0.66 0.3 2.0	0.7 .04 2.0
Loam, Silty Loam, Silt	Cynex 4L Atrazine 4L Lasso EC	Quarts	0.66 0.3 2.0	0.7 0.4 2.0	0.8 0.4 2.0
Sandy Clay Loam, Clay Loam, silty Clay Loam	Cynex 4L Atrazine 4L Lasso EC	Quarts	0.7 0.4 2.25	0.8 0.4 2.25	0.9 0.5 2.25
Sandy Clay, Silty Clay, Clay	Cynex 4L Atrazine 4L Lasso EC	Quarts	0.8 0.4 2.25	0.9 0.5 2.25	0.9 0.5 2.5

117047  
1

Peat or Muck	*	*	*	*	*
Eroded Slopes or Knobs Soils with pH greater than 8.0	*	*	*	*	*

\*Not Recommended

- (a)When using LASSO 4EC on sorghum, only SCREEN safened seed should be planted.
- (b)To enhance weed control in areas of less than 25 inches of annual precipitation or where long dry periods are common, this treatment may require shallow incorporation with a tool such as a field cultivator operated at 5-7 mph.

\*\*For organic matter content between those listed, adjust the rate proportionately.

**TABLE 22**  
**PREEMERGENCE BROADCAST APPLICATION RATES IN QUARTS OF Cynex 4L, QUARTS OF ATRAZINE 4L AND QUARTS OF PROPACHLOR (RAMROD) 4L PER ACRE ON GRAIN SORGHUM**

Percent Organic Matter in the Soil\*\*

Soil Texture Description	2%	3%
Ramrod	Qts Cynex 4L/Qts Atraz/Qts Ramrod	Qts Cynex 4L/Qts Atraz/Qts

Soil Texture Description	2% (Qts Cynex 4L/Qts Atraz/Qts Ramrod)	3% (Qts Cynex 4L/Qts Atraz/Qts)
Sand, Loamy Sand	Do Not Use	Do Not Use
Sandy Loam	0.6(a)	0.3
Loam, Silt Loam, Silt	0.7	0.4
Sandy Clay Loam, Clay Loam, Silty Clay Loam	0.8	0.4
Sandy Clay, Silty clay, Clay	0.9	0.5
Peat or Muck	*	*

oded Slopes or Knobs Soils with pH greater than 8.0	*	*	*	*	*	*
--	---	---	---	---	---	---

\*Not Recommended

(a) The first number is quarts/acre of Cynex 4L. The second number is quarts/acre of atrazine 4L. The third number is quarts/acre of Ramrod 4L.

\*\*For organic matter content between those listed, adjust the rate proportionately.

### Cynex 4L IN THREE-WAY COMBINATIONS

Cynex 4L plus atrazine plus DUAL, LASSO or RAMROD may be applied preemergence or with shallow incorporation for weed control in grain sorghum only in the states of Kansas, Nebraska and South Dakota. These treatments provide control of weeds listed on this label in the Weeds Controlled section for Cynex 4L on grain sorghum. Additional weeds controlled by these combinations include:

- Grasses
- Barnyardgrass (1)      Witchgrass (2)
- Stinkgrass (2)        Giant Foxtail
- Fall Panicum

(2) Refer to Weeds Controlled section on grain sorghum for explanation.

**Rotational Crops:** If the crop treated with this combination is lost, corn or grain sorghum may be replanted immediately without retreatment. Refer to the "General Information" section of the Cynex 4L label and to the tank mix product labels for additional information regarding crop rotation.

#### Cynex 4L Plus ATRAZINE Plus DUAL

DUAL can be used on grain sorghum only with seed treated with CONCEP II seed safener.

Use Cynex 4L plus DUAL 8E at the proper rate for soil texture and organic matter indicated in Table 20. Use only on soils having at least 1% organic matter.

#### Cynex 4L Plus ATRAZINE Plus LASSO

LASSO can be used on grain sorghum only with seed treated with SCREEN seed safener.

Use Cynex 4L plus Atrazine 4L plus LASSO 4EC at the proper rate for soil texture and organic matter indicated in Table 21. Use only on soils having at least 1% organic matter.

#### Cynex 4L Plus ATRAZINE Plus RAMROD

Use Cynex 4L plus Atrazine 4L plus RAMROD at the proper rate for soil texture and organic matter indicated in Table 22.



4914

### NOTICE OF WARRANTY

Griffin warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purposes stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Griffin. In no case shall Griffin be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer. Griffin MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

Bladex 4L, EXTRAZINE II-Trademark of E.I. Du Pont de Nemours & Co. (Inc)  
BANVEL - Trademark of Sandoz Crop Protection Corp.  
DUAL, CONCEP II - Trademarks of CIBA-Geigy Corp.  
ERADICANE , SUTAN+ - Trademarks of ICI Americas, Inc  
LASSO, RAMROD, SCREEN - Trademarks of Monsanto Co.  
PRINCER CALIBER - Trademarks of CIBA-Geigy Corp.  
PROWL - Trademark of American Cyanamid.  
TREFLAN - Trademark of Elanco Products Company.  
ZORIAL RAPID 80 - Trademark of Zoecon Corporation.