 <p>U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (7505C) 401 "M" St., S.W. Washington, D.C. 20460</p> <p>NOTICE OF PESTICIDE: <u> X </u> Registration <u> </u> Reregistration</p> <p>(under FIFRA, as amended)</p>	EPA Reg. Number: 1812-404	Date of Issuance: JUL 20 1998
	Term of Issuance: Conditional	
	Name of Pesticide Product: Chlorfos 15G	

Name and Address of Registrant (include ZIP Code):
Griffin Corporation
P.O. Box 1847
Valdosta, GA 31603-1847


Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c) (7) (A) provided that you:

1. Submit and/or cite all data or other material required for registration/reregistration of your product under FIFRA sec. 3(c) (5) or FIFRA sec. 4 when the Agency requires all registrants of similar products to submit such data.
2. Make the labeling changes listed below before you release the product for shipment:
 - a. Replace the EPA Registration No. with "1812-404".

Signature of Approving Official:  Tina E. Levine, PhD, Chief Insecticide-Rodenticide Branch Registration Division (7505C)	Date: 7/20/98
--	----------------------

Page Two
1812-404

5. Submit two (2) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records. If you have questions regarding this notice, contact Dani Daniel of my staff at 703 305-5409.

Enclosure:

CHLORFOS® 15G

INSECTICIDE

LOW ODOR FORMULA

GRANULAR

For control of various insects infesting certain field
and vegetable crops.

Active Ingredient:

Chlorpyrifos: O,O-diethyl

O-(3,5,6-trichloro-2-pyridinyl)

phosphorothioate..... 15.0%

inert Ingredients: 85.0%

Total:..... 100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la
explique a usted en detalle. (If you do not understand the label, find
someone to explain it to you in detail.)

STATEMENT OF PRACTICAL TREATMENT

Organophosphate

If swallowed: Call a physician or Poison Control Center. Drink 1 or 2
glasses of water and induce vomiting by touching back of throat with
finger. Do not induce vomiting or give anything by mouth to an
unconscious person.

In eyes: Flush with plenty of water for 5 minutes. Get medical
attention if irritation persists.

If on skin: Wash with plenty of soap and water. Get medical attention
if irritation persists.

Note to Physicians: Chlorpyrifos is a cholinesterase inhibitor. Treat
symptomatically. If exposed, plasma and red blood cell cholinesterase
tests may indicate significance of exposure (baseline data are useful).

Atropine, only by injection, is the preferable antidote. Oximes, such as
2-PAM/protopam, may be therapeutic if used early; however, use only
in conjunction with atropine. In case of severe acute poisoning, use
antidote immediately after establishing an open airway and respiration.

See side panel for additional precautionary statements.

Mfg. For:

EPA Registration No. 1812-UNO

EPA Establishment No.

Net Contents:

Griffin Corporation

1812-UNO

JUL 20 1998

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 1812-404

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS (AND DOMESTIC ANIMALS)

CAUTION

Harmful if swallowed. Causes eye irritation. Avoid contact with eyes,
skin or clothing. Avoid breathing dust.

Personal Protective Equipment

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning and maintaining
Personal Protective Equipment (PPE). If no such instructions for
washables, use detergent and hot water. Keep and wash PPE
separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or
using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash
thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to birds and wildlife, and extremely toxic to fish
and aquatic organisms. Do not apply directly to water, or to areas
where surface water is present or to intertidal areas below the mean
high water mark. Drift and runoff from treated areas may be hazardous
to aquatic organisms in adjacent aquatic sites. Cover or incorporate
spills. Do not contaminate water when disposing of equipment
washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner
inconsistent with its labeling.

Read all Directions for Use before applying.

Not for sale, distribution or use in the States of California and Arizona.

Do not apply this product in a way that will contact workers or other
persons either directly or through drift. Only protected handlers may be
in the area during application. For any requirements specific to your
state or tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the
Worker Protection Standard, 40 CFR part 170. This Standard contains
requirements for the protection of agricultural workers on farms,
forests, nurseries, and greenhouses, and handlers of agricultural
pesticides. It contains requirements for training, decontamination,
notification, and emergency assistance. It also contains specific
instructions and exceptions pertaining to the statements on this label
about personal protective equipment (PPE), and restricted-entry interval.
The requirements in this box only apply to uses of this product that are
covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the
restricted entry interval (REI) of 12 hours.

Exception: If the product is soil-injected or soil-incorporated, the
Worker Protection Standard, under certain circumstances, allows
workers to enter the treated area if there will be no contact with
anything that has been treated.

PPE required for early entry to treated areas that is permitted under the
Worker Protection Standard and that involves contact with anything
that has been treated, such as plants, soil, or water, is:

- Coveralls
- Shoes plus socks
- Waterproof gloves

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

APPROVED USES

Corn Field, Popcorn, Sweet Corn (including corn grown for seed)*

Application Rates and Pests Controlled

Pests Controlled	Banded Applications (Ounces per 1,000 Feet of Row)				Preplant Broadcast Application (lb/Acre)	Aerial Broadcast Application (lb/Acre)
	At Plant Treatments			Postplant Treatment		
	T-Band	Band	In-Furrow			
northern and western corn rootworm larvae	8	8	8	8	-	-
southern corn rootworm larvae	8	8	8	8	-	-
cutworms* *	8	8	8	-	6.75 to 13.5	-
wireworms* * *	8	8	8	-	13.5	-
Billbugs	8	8	-	-	13.5	-
seed corn maggots	8	-	8	-	-	-
grubs* * * *	8	-	8	-	13.5	-
lesser cornstalk borer	8	8	-	-	-	-
seed corn beetle	8	-	8	-	-	-
flea beetle larvae	-	-	-	-	13.5	-
symphylans	8	8	-	-	-	-
armyworms	-	-	-	6 to 8	-	-
European and southwestern corn borer						
1st Generation	-	-	-	3.5 to 8	-	5.0 to 6.5
2nd Generation	-	-	-	6 to 8	-	6.5
fire ants* *	8	-	-	-	-	-
chinch bug* *	8	-	8	-	-	-

* Chlorfos 15G insecticide is compatible with all ALS inhibitor herbicides, including Accent, Beacon and Broadstrike herbicides, applied in accordance with label recommendations. Refer to the product labels for additional precautions and directions for use.

** 8 oz rate-suppression only for in-furrow treatments.

*** For best control, apply as an in-furrow treatment. Consider using a hopper box insecticidal seed treatment with band and T-band applications. Use 12 oz/1,000 ft of row for severe infestations.

**** Use 12 oz/1,000 ft of row for severe infestations.

At Plant Band Treatment: In conventional, minimum and no-till corn, place the granules in a 6 or 7 inch wide band over the row behind the planter shoe in front of the press wheel, and incorporate into the top one inch of soil using tines or chains or other suitable equipment. For cutworm control under very dry soil conditions, best results will be obtained through proper incorporation. In soils over 10% organic matter, use the highest rate given.

At Plant T-Band Treatment: In conventional, minimum and no-till corn, for control of rootworm, cutworm, and billbug. In addition to lesser cornstalk borer, symphylans, wireworms, seed corn beetle, seed corn maggot and grubs, apply as a T-band over an open seed furrow. In conventional and minimum-till corn, incorporate into the top 1/2 to 1 inch of soil using suitable equipment. For severe wireworm and grub infestations, use 12 oz/1,000 ft of row. Chlorfos 15G has demonstrated suppression of certain soil-borne pathogens which may result in physiological and agronomic advantages to corn under environmental stress conditions when compared to corn not treated with Chlorfos 15G.

At Plant In-Furrow Treatment: Apply Chlorfos 15G at planting time as an in-furrow treatment in conventional, minimum and no-till corn. Direct the granules into the planter shoe with the seed or place the applicator tube directly behind the planter shoe so that the granules drop into the seed furrow or place the granular band applicator behind the planter shoe so that the granules fall on the soil surface and into the open seed furrow and are covered with soil. Equivalent rates of Chlorfos 15G per acre for various row spacings are given in the accompanying tables.

Postplant Treatment: To control corn rootworm larvae, apply Chlorfos 15G at cultivation time by placing the granules at the base of the plant on both sides of the row just ahead of the cultivation shovels and cover the soil. To control European and southwestern corn borer larvae, apply Chlorfos 15G in a band over the row so that the granules are directed into the whorl or use a post plant broadcast treatment. Use application rates of 3.5 to 4 oz of Chlorfos 15G per 1,000 feet of row for low to moderate first generation infestations before larvae have entered corn stalks. Use application rates of 6 to 8 oz of Chlorfos 15G per 1,000 feet of row for severe first generation infestations and all second generation infestations before larvae have entered corn stalks. Consult your state agricultural experiment station or extension service specialist for proper time to treat and local threshold information. Scouting for insect damage is strongly encouraged.

Postplant Broadcast Treatment: Use Chlorfos 15G for control of European and southwestern corn borers by uniformly broadcasting the granules over the corn plants by aerial application or by applying the granules into the corn whorls by ground application. Apply at a rate of 5.0 lb per acre for low to moderate first generation infestations or at 6.5 lb per acre for severe first generation infestations and all second generation infestations. Apply before larvae have entered corn stalks. Consult your state agricultural experiment station or extension service specialist for local threshold information. Scouting for insect damage is strongly encouraged.

Preplant Broadcast Treatment: Use equipment that will evenly distribute the granules over the treated area. Incorporate the granules into the soil using an S tine, C tine, disc cultivator, or other suitable incorporation equipment set to cut to a depth of 4 to 6 inches. Plant the crop in the usual manner as soon as possible after treatment.

Restrictions: For soil control, do not exceed the equivalent of 16 ounces of Chlorfos 15G per 1,000 feet row or 13.5 pounds of Chlorfos 15G per acre applied as a broadcast application per crop season. For foliar insect control, do not exceed the equivalent of 16 ounces of Chlorfos 15G per 1,000 feet row or 13 pounds of Chlorfos 15G per acre per crop season. Do not apply within 35 days before harvest of grain. Do not allow livestock to graze in treated areas nor harvest treated corn silage as feed for meat or dairy animals within 14 days after last treatment. Do not feed treated corn fodder to meat or dairy animals within 35 days after last treatment.

Onion (Dry Bulb)

Pest Controlled: onion maggot

Application Rate: 3.7 ounces per 1,000 feet or row at an 18-inch row spacing

At Plant In-Furrow Treatment: Apply as an at planting in-furrow treatment. In Colorado, Idaho, Washington, and Oregon, to control onion maggots in onions planted in double rows with rows spaced 2 to 4 inches apart, apply Chlorfos 15G at the rate of 3.7 oz per 1,000 feet of double row. Place the granules in a 5 to 7 inch wide band over both rows behind the planter shoe and in front of the press wheel to achieve shallow incorporation. Do not exceed 6.7 lb per acre.

Restrictions: Do not make more than one application per year.

Citrus Grove Floors

Pests Controlled: fire ants and other ant species

Application Rate: 6.7 lb/acre

Postplant Broadcast Treatment: To control foraging ants and suppress mounds, apply Chlorfos 15G with ground application equipment. Use a suitable granular applicator, such as a cyclone fertilizer spreader, that will uniformly broadcast the granules over the grove floor. Retreat as needed. Chlorfos 15G may be custom blended with granular fertilizers provided that application of the blended Chlorfos 15G plus fertilizer mixture can be applied uniformly to the grove floor. Do not apply where weed growth or other obstructions would impede uniform coverage of the grove floor.

Restrictions: Do not apply more than 67 pounds of Chlorfos 15G per acre per crop season. Do not apply last treatment within 28 days before harvest. Do not allow livestock to graze in treated areas.

Sorghum/Milo

Application Rates and Pests Controlled

Pests Controlled	Banded Applications (Ounces per 1,000 Feet of Row)	
	At Plant Treatments	
	T-Band	Band
lesser cornstalk borer	4 to 12	4 to 12
corn rootworm and cutworms	8	8
fire ants *	8	-
chinch bug *	8	-

* 8 oz. rate - suppression only

At Plant T-Band or Band Treatments: Apply in a 6 to 8 inch band over the row and incorporate into the top 1-inch of soil using suitable equipment. Equivalent rates of Chlorfos 15G per acre for various row spacings are given in Table 1. Use the lowest rate for lesser cornstalk borer control when protection is desired for 2 to 3 weeks and higher rates for longer residual activity. It is absolutely necessary to incorporate the granules, especially at lower rates.

Restrictions: Do not make more than one application per season.

Peanut

Use Chlorfos 15G to control cutworms, lesser cornstalk borer, southern corn rootworm larvae, suppress wireworms, and inhibit the growth and development of white mold (southern blight) disease caused by *Sclerotium rolfsii*. Chlorfos 15G will control only those cutworms existing in the soil from the time of application up to 30 days following application.

Application Rates and Pests Controlled

Pests Controlled	Banded Applications (Ounces/1,000 Feet of Row)		Aerial Broadcast (lb/Acre)
	At Plant Treatment	Postplant Treatment	
*Preventative Treatments: cutworms lesser cornstalk borer southern corn rootworm larvae wireworms white mold (southern blight)	7.5 to 15	7.5 to 15	-
*Rescue Treatments: lesser cornstalk borer	-	7.5 to 15	10 to 13
Burrowing bug	-	-	13.3

*At Plant Preventive Treatment: Apply Chlorfos 15G in a 6 to 12 inch band over the row behind the planter shoe and in front of the press wheel. Incorporate granules to a depth of 1-inch with tines or chains or other suitable equipment. If the 7.5 oz rate is used at planting time,

then another application of 7.5 oz per 1,000 feet of row should be made postplant to extend control.

***Postplant Preventative Treatment:** Apply Chlorfos 15G to peanuts at early flowering to pegging stage of growth in a 6 to 8 inch band over the row. For extended insect control and continued suppression of white mold (southern blight), a second application of Chlorfos 15G may be made. Best suppression of white mold (southern blight) is obtained by applying the maximum rate of 15 oz per 1,000 ft of row for each postplant treatment. Irrigation or rain following application is needed to enhance treatment effectiveness for suppression of white mold. Under conditions of heavy white mold pressure, a suitable fungicide may also be required and must be applied separately.

****Band Rescue Treatment:** Use Chlorfos 15G for the control of lesser cornstalk borer when the insect first appears, usually just prior to or at pegging. Apply in a 10 to 18 inch band over the fruiting zone.

****Broadcast Rescue Treatment:** Use Chlorfos 15G for the control of lesser cornstalk borer when the insect first appears, usually just prior to or at pegging. Uniformly apply at a rate of 10 to 13 pounds per acre by aerial broadcast application. For control of burrowing bugs, use Chlorfos 15G when adults first appear, usually at mid to late pegging. Uniformly apply at a rate of 13.3 pounds per acre by aerial broadcast application. Care must be taken to ensure proper equipment calibration and delivery.

Restrictions: Do not apply more than 15 ounces of Chlorfos 15G per application per 1,000 feet of row or 13.3 pounds of Chlorfos 15G per acre per application. Do not apply more than 30 oz per 1,000 ft of row or 26.6 pounds per acre per crop season. Do not harvest within 21 days after treatment. Do not feed peanut forage or hay to meat or dairy animals. The combined total of preplant and postplant applications of Chlorfos 4E and Chlorfos 15G must not exceed 4 pounds of active ingredient per acre per crop season.

Soybean

Application Rates and Pests Controlled

Pests Controlled	Banded Applications (Ounces per 1,000 Feet of Row)		
	At Plant Treatments		Postplant Treatment
	T-Band	Band	
lesser cornstalk borer	8	8	8
cutworms			

At Plant and Postplant Treatments: Use Chlorfos 15G insecticide to control larvae of the lesser cornstalk borer and cutworms by application at planting time or postemergence as a band (row) treatment at the rate of 8 oz per 1,000 feet of row. In the southeast apply 4 to 8 oz per 1,000 feet of row as an at-plant treatment. Equivalent rates of Chlorfos 15G per acre for various row spacing are given in the accompanying table. When applied at planting time incorporate the granules into the top 1 inch of soil by placing in a 4 to 10 inch band over the row behind the planter shoe and ahead of the press wheel. A drag chain can also be used for incorporation. For postemergence treatment when insects first appear incorporate the granules in a 4 to 10 inch band to a depth of 1/2 to 1 inch using a suitable cultivator. Apply Chlorfos 15G with equipment that will provide uniform distribution of the granules. Do not apply as an in-furrow treatment. For suppression of fire ants, use Chlorfos 15G at 8 oz per 1,000 feet of row as an at-plant T-band treatment.

Restrictions: Do not make more than one application per season.

Sugar Beet

Granular insecticides, including Chlorfos 15G, may contribute to the stress of the sugar beet plant under certain environmental conditions. This stress may reduce plant stand or interfere with normal plant development. Herbicides used preplant incorporated may interact with insecticides and enhance this stress.

Application Rates and Pests Controlled

Pests Controlled	Banded Applications (Ounces per 1,000 Feet of Row)		
	At Plant Treatments		Postplant Treatment
	T-Band	Band	
Sugar beet root maggot	-	4.5 to 9.0	6.5 to 9.0
cutworms	-	6.6 to 9.0	-
wireworms (suppression)	-	6.5 to 9.0	-

At Plant Band Treatment: To control sugar beet root maggot larvae and cutworms at planting time, place Chlorfos 15G in a band 4 to 5 inches wide behind the planter shoe, over the drill row, and in front of the press wheel. Do not apply granules in direct contact with seeds. Apply Chlorfos 15G at the rate of 4.5 to 9 ounces per 1,000 feet of row (equivalent to 6.75 to 13.5 lb per acre based on a 22 inch row spacing). When root maggot populations are expected to be low, apply Chlorfos 15G at a rate of 4.5 ounces per 1,000 feet of row (equivalent to 6.75 lb per acre based on 22 inch row spacing). If initial adult fly activity indicates higher than anticipated populations, apply Chlorfos 4E at or near the time of peak adult emergence to augment control. (Review label for Chlorfos 4E for recommended use rates, application timing, methods of application, and insecticide resistance management). Incorporate Chlorfos 15G into the top 1/2 to 1 inch of soil using suitable equipment.

Postemergence Band Treatment: For postemergence control of sugar beet maggot larvae, place Chlorfos 15G in a band 3 to 5 inches wide over the beet row (up to 2 to 4 true leaf stage of plant growth). Apply Chlorfos 15G at the rate of 6.5 to 9 oz per 1,000 feet of row (equivalent to 10 to 13.5 lb per acre based on a 22 inch row spacing). Incorporate Chlorfos 15G into the top 1/2 to 1 inch of soil using a suitable incorporation device.

Restrictions: Do not make more than one application per year.

Sunflower

Pest Controlled: cutworms

Application Rate: 8 oz per 1,000 feet of row

At Plant Band Treatment: Place the granules in a 7 inch wide band over the row behind the planter shoe in front of the press wheel and incorporate into the top 1 inch of soil using suitable equipment.

Restrictions: Do not make more than one application per season.

Sweet Potato

Pests Controlled: wireworms (*conoderus*), flea beetles (*Systema*), and sweet potato beetle

Application Rate: 13.5 lb/acre

Preplant Broadcast Treatment: Use Chlorfos 15G to reduce the feeding damage caused by populations of the listed pests. Evenly broadcast the granules over the soil surface and then incorporate the granules into the soil to a depth of 4 to 6 inches using a rotary hoe, disc cultivator, or other suitable equipment. Plant the crop in the usual manner no later than 14 days after treatment (any delay in planting will reduce the length of time that Chlorfos 15G will protect against feeding damage). Chlorfos 15G will not control false wireworm or whitefringed beetle and other grubs that attack sweet potatoes.

Restrictions: Do not make more than one application per season. Do not harvest within 125 days of treatment

Tobacco

Pests Controlled: cutworms, flea beetle larva, root maggots, and wireworms

Application Rate: 13.5 to 20.0 lb/acre

Preplant Broadcast Treatment: Apply Chlorfos 15G one week before transplanting, using equipment that will evenly distribute the granules over a treated area. Immediately following application, incorporate the granules into the soil to a depth of 2 to 4 inches using suitable equipment. The application of Chlorfos 15G will also suppress movement of imported fire ant into treated field.

Restrictions: Do not make more than one application per season.

Brassica (Cole) Crops

(Bok Choy, Broccoli, Broccoli Raab, Brussels Sprout, Cabbage, Cauliflower, Chinese Broccoli, Chinese Cabbage, Collards, Kale, Kohlrabi, Rutabagas, and Turnips)

Pest Controlled: root maggot

Application Rate: 4.6 to 9.2 ounces per 1,000 feet of row

At Plant T-Band Treatment: For direct seeded and transplanted crops, apply Chlorfos 15G as a 4 inch wide band centered over the row. This application requires a spreader or splitter on the end of the applicator drop tube. Shallow incorporation is necessary. Placement behind the planter shoe and in front of the press wheel is recommended.

Restrictions: Do not apply more than 7 1/2 pounds of Chlorfos 15G per acre to crops planted in 40 inch rows or more than 15 pounds of Chlorfos 15G per acre to crops planted in 20 inch rows (or two rows per bed). Use proportional amounts for other row spacings not to exceed 15 pounds of Chlorfos 15G per acre. Do not make more than one application per season. Do not use rutabaga tops for food or feed purposes.

Radish

Pest Controlled: root maggot

Application Rate: 3.3 ounces per 1,000 feet of row

At Plant In-Furrow Treatment: Place the granules in the seed furrow with the seed at planting time.

Restrictions: Do not apply more than 18.3 pounds of Chlorfos 15G per acre or make more than one application per season.

Table 1

Application Rates Table-Application Rates/1,000 Ft. of Row and Equivalent/Acre at Different Row Spacings

Amount of Chlorfos 15G Per 1,000 Feet of Row	Pounds of Chlorfos 15G Required Per Acre from Various Row Spacing							
	40"	38"	36"	34"	32"	30"	22"	18"
3.7 ounces	3.0	3.2	3.4	3.6	3.8	4.0	5.5	6.7
4.0 ounces	3.3	3.4	3.6	3.8	4.1	4.4	5.9	7.3
4.5 ounces	3.7	3.9	4.1	4.3	4.6	4.9	6.7	8.2
6.0 ounces	4.9	5.2	5.4	5.8	6.1	6.5	8.9	10.9
6.5 ounces	5.3	5.6	5.9	6.2	6.6	7.1	9.5	11.8
7.5 ounces	6.1	6.4	6.8	7.2	7.7	8.2	11.1	13.6
8.0 ounces	6.5	6.9	7.3	7.7	8.2	8.7	11.9	14.5
9.0 ounces	7.4	7.7	8.2	8.6	9.2	9.8	13.4	16.3
12.0 ounces	9.8	10.3	10.9	11.5	12.3	13.1	17.8	21.8
15.0 ounces	12.3	12.9	13.9	14.4	15.3	16.3	22.3	27.2
16.0 ounces	13.1	13.8	14.5	15.4	16.3	17.4	23.8	29.0

Table 2
Equipment Calibration and Calibration Settings for Different Types
of Equipment

General Instructions for Equipment Calibration

Caution: The following chart lists suggested initial gauge settings for application of Chlorfos 15G with one hopper opening per row. Be sure to check the actual application rate under your operating conditions.

1. Fill hopper.
2. Attach a plastic bag to tube opening.
3. Set your planter to the initial settings shown on chart.
4. Measure off 1,000 row feet and drive your planter the pre-measured distance at your desired speed.
5. Each bag should contain 6 to 8 ounces (wt.) of granules depending on your desired rate.
6. If the result is over or under the desired rate - adjust the settings and repeat the calibration.

CALIBRATION SETTINGS

Planter Type	Speed (mph)									
	4		5		6		7		8	
	Application Rate, oz. Per 1,000 row ft.									
	8	16	8	16	8	16	8	16	8	16
	Gauge Setting									
Gandy ¹	21.4	30.2	23.7	32.4	26.0	36.0	27.7	39.0	30.2	41.0
John Deere ¹ Max-Emerge ²	20	44	26	46	30	49	35	52	40	54
John Deere ¹ 7000 Max-Emerge (Old Nos. on Gate)	14	22	16	24	18	26	19	28	21	30
John Deere ¹ 7000 Max-Emerge (Even Nos. on Gate)	17	30	20	33	24	35	26	36	28	38
John Deere ² 71 Flexi-Planter and Older Planters	$\frac{1}{30}$	$\frac{2}{17}$	$\frac{2}{5}$	$\frac{2}{22}$	$\frac{2}{9}$	$\frac{2}{27}$	$\frac{2}{13}$	$\frac{2}{31}$	$\frac{2}{16}$	$\frac{3}{16}$
Allis Chalmers ³ 70 Series	8	13	8	13	8	13	8	13	8	13
Allis Chalmers ⁴ 78 & 79 Series	$\frac{1}{9.0}$	$\frac{3}{3.0}$	$\frac{2}{3.3}$	$\frac{3}{9.5}$	$\frac{2}{6.0}$	$\frac{4}{3.0}$	$\frac{2}{9.0}$	$\frac{5}{4.0}$	$\frac{3}{2.5}$	$\frac{6}{0}$
Noble ^{1(NEW)}	11	19	14	22	16	25	17	28	19	31
White Planter	11	19	14	22	16	25	17	28	19	31
International Harvester ⁴	$\frac{1}{9.0}$	$\frac{3}{3.0}$	$\frac{2}{3.3}$	$\frac{3}{9.5}$	$\frac{2}{6.0}$	$\frac{4}{3.0}$	$\frac{2}{9.0}$	$\frac{5}{4.0}$	$\frac{3}{2.5}$	$\frac{6}{0}$
Buffalo All-Flex ⁵ (Fleischer Mfg.)	4 7/8	10	4 7/8	10	4 7/8	10	-	-	-	-

¹ Gauge setting

² Gauge setting with range 1 & 2 - number is range,
notch

³ Gauge setting is constant regardless of speed.

⁴ Gauge setting shown with stem gates & dial settings - number shown is gate,
dial

⁵ Number of turns open on the adjustment nut.

WARRANTY STATEMENT

GRIFFIN warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of GRIFFIN. In no case shall GRIFFIN be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. The exclusive remedy of any buyer of this product for any and all losses, injuries, or damages resulting from or in any way arising from the use, handling, or application of this product, whether in contract, warranty, tort, negligence, strict liability, or otherwise, shall not exceed the purchase price paid for this product or, at Griffin Corporation's election, the replacement of this product. GRIFFIN MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESSED OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

Chlorfos is a registered trademark of Griffin Corporation