

## UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

1812-437

09/12/2000

1/6

SEP 12 2000

Judy A. Smith  
Griffin L.L.C.  
P.O. Box 1847  
Valdosta, GA 31603-1847

Dear Ms. Smith:

Subject: Supplemental Labeling - Adjust Application Rate for Various Crops  
Equus 720 Fungicide  
EPA Registration No. 1812-437  
Your Submission Received On August 14, 2000

The supplemental labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable with the following provisions:

- 1) Within the Application Directions for Soybean Determinate Varieties (Southern) for the three application program, add a statement to require a minimum interval of 14 days between applications.
- 2) Within the directions for use on Tomato, we note that the required minimum interval(s) between applications is specified under the Disease column. The minimum interval(s) must be specified in the column for Application Directions, either in addition to or in place of appearing in the Disease column.
- 3) Within the directions for use on Tree and Orchard Crops, revise the heading currently reading "Equus 720 Rate Pints/100 gallons\*" to read "Equus 720 Rate Pints/100 gallons/acre\*". In addition, add a footnote to correspond with the asterisk to read "See dilution instructions for tree and orchard crops on the main labeling."

DK: 305-7546: FB/PM 22

CONCURRENCES

SYMBOL	7505C							
SURNAME	D. KENNY							
DATE	9/12/2000							

2/6

A stamped copy is enclosed for your records. Please submit one (1) final printed copy for the referenced label, incorporating the above changes, before releasing the product for shipment.

Sincerely yours,

Daniel C. Kenny  
Product Manager (22)  
Fungicide Branch  
Registration Division (7505C)

Enclosure



ACCEPTED  
with COMMENTS  
In EPA Letter Dated

SEP 12 2000

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.  
1812-437

3/6

**SUPPLEMENTAL LABELING  
FOR USE IN ALL STATES EXCEPT CALIFORNIA**

**Equus™ 720 Fungicide**

EPA REG. NO. 1812-437

ACTIVE INGREDIENTS:	BY WEIGHT
Chlorothalonil (tetrachloroisophthalonitrile) .....	54.0%
INERT INGREDIENTS .....	46.0%
TOTAL .....	100.0%

Contains 6.0 Pounds of Active Ingredient per Gallon (720 grams per liter).

**KEEP OUT OF REACH OF CHILDREN  
WARNING-AVISO**

**Directions For Use:** It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. See package label for additional use directions.

**FIELD CROPS**

CROP	DISEASES CONTROLLED	RATE OF Equus 720 PER APPLICATION PINTS/ACRE	SEASONAL LIMITS (PINTS/ACRE/YEAR)	APPLICATION DIRECTIONS
CRAPEFRUIT	Fruit rot Lophodermium Leaf/Twig Blight	4 to 6.67	20.0	Apply at early bloom and repeat at 10- to 14-day intervals. Under severe disease conditions, use the 6.67 pints per acre rate on a 10-day schedule. DO NOT apply Equus 720 more than 3 times per season, or within 50 days before harvest. DO NOT apply to bogs when flooded or allow release of irrigation water from bogs for at least 3 days following application.  Equus 720 may be applied through sprinkler irrigation equipment. Use 300 gallons of water per acre through solid set systems only. See calibration directions preceding this section.
PAPAYA	Alternaria Fruit Spot Anthracnose Stem End Rot	3.0	9.0	Apply with ground equipment only. Use sufficient water to obtain adequate coverage of fruit and leaves. Begin treatment when conditions favor development of disease and continue treatments at 14-day intervals until weather conditions no longer favor disease development.

<b>PARSNIP</b>	Alternaria Leaf Spot Downy Mildew Anthracnose Botrytis Blight (Gray Mold) Bottom Rot (Rhizoctonia)	1.5 to 2	8.0	Apply in sufficient water to obtain adequate coverage. Make the first application at the first sign of disease or when conditions are favorable for infection. Continue applications on a 7- to 10-day schedule.
<b>PASSION FRUIT (Hawaii only)</b>	Alternaria Fruit and Leaf Spot (Passion Fruit Brown Spot)	2	10.0	Apply with ground equipment in sufficient water to obtain adequate coverage of fruit and leaves. Begin treatment when fruit spots appear (April to July) and continue treatments at 14-day intervals until weather conditions no longer favor disease development.
<b>SOYBEAN DETERMINATE VARIETIES (Southern)</b>	Anthracnose Diaporthe Pod and Stem Blight Frogeye Leaf Spot (Cercospora sojina) Purple Seed Stain Cercospora Leaf Blight (Cercospora kikuchii) Septoria Brown Spot		6.0	Apply in sufficient water to obtain complete coverage, using at least 5 gallons of water per acre for aerial application. Use the three application program in areas having a history of moderate to severe disease intensity. Equus 720 may be applied through sprinkler irrigation equipment. Follow application and calibration directions preceding this section. DO NOT exceed a total of 3 applications per season. DO NOT apply within 6 weeks of harvest.
		1.5 to 2.4		<b>Two application program:</b> Make the first application at early pod set (R3 stage, when the majority of pods are 1/8 to 3/8 inches in length) and the second at beginning of seed formation (R5) which occurs about 14 days later.
		1 to 2.0		<b>Three application program:</b> Make the first application at the beginning of flowering (R1), the second at early pod set (R3), and the third at beginning of seed formation (R5).
	Stem Canker ( <i>Diaporthe phaseolorum</i> var. <i>caulivora</i> )	1		Apply in 10 to 20 gallons of water per acre, as a band treatment, directing spray to provide coverage of entire plant. Make the application at time of emergence of the second trifoliate leaves (V2). If conditions favor stem canker disease, make a second and third application. Make all applications at 14-day intervals.

SOYBEAN INDETERMINATE VARIETIES (Northern)			6.0	Apply in sufficient water to obtain complete coverage, using at least 5 gallons of water per acre for aerial application. Use the three application program in areas having a history of moderate to severe disease intensity. Equus 720 may be applied through sprinkler irrigation equipment. Follow application and calibration directions preceding this section. DO NOT exceed total of 3 applications per season. DO NOT apply within 6 weeks of harvest.
		1.5 to 2.4		<b>Two application program:</b> Make the first application when largest pods are 1 to 1.5 inches in length and make the second application 14 days later. Equus 720 may be co-applied with Benlate® SP as a tank mix for disease control in indeterminate (northern) soybeans. Use 1 pint of Equus 720 plus 8 ounces of Benlate® SP per acre. Make the first application when pods near the top of plants are 0.5 to 1 inch in length and a second application 14 days later.
		1 to 2		<b>Three application program:</b> Make the first application 1 week after first flowering and continue applications at 14-day intervals.
TOMATO	Foliage (apply every 7 to 10 days): Early Blight Late Blight Gray Leaf Spot Gray Leaf Mold Septoria Leaf Spot Target Spot	1.35 to 2	20.1	Apply in sufficient water to obtain adequate coverage. Begin applications when dew or rain occur and disease threatens. Equus 720 may be combined in the spray tank with EPA-registered pesticide products that claim copper as the active ingredient and are labeled for control of bacterial diseases of tomatoes. Check the copper manufacturer's label for specific instructions, precautions and limitations prior to mixing with Equus 720. DO NOT use with Copper-Count N in concentrated spray suspensions.
	Fruit (apply every 7 to 14 days beginning at fruit set): Anthracnose Alternaria Fruit Rot (Black Mold) Botrytis Gray Mold Late Blight Fruit Rot Rhizoctonia Fruit Rot	2 to 2.88		Equus 720 may be applied through sprinkler irrigation equipment (solid set or portable wheel move systems only). See calibration directions preceding this section.

6/6

# **TREE AND ORCHARD CROPS**

Crop	Diseases	Equus 720 Rate Pints/Acre	Equus 720 Rate Pints/100 gallons*	Seasonal limit Pints/Acre	Application Directions
<b>FRUIT TREES</b> Apricot Cherry, Sweet Cherry, Tart Nectarine Peach Plum Prune	Leaf Curl Coryneum Blight (Shothole)	3.1 to 4.1	1.0	20.6	For best control of both diseases, apply at leaf fall in late autumn, using sufficient water and proper sprayer calibration to obtain uniform coverage. When conditions favor high disease levels, use the high rate of application and apply once or twice more in mid-to-late winter before budswell. If the leaf fall application is not practical, application of Equus 720 for control of leaf curl may be made at any time prior to budswell the following spring. Where Coryneum blight (shothole) occurs, also apply at budbreak to protect newly emerging leaves and at shuck split to prevent fruit infections.
	Brown Rot Blossom Blight Lacy (Russet) Scab (Plum/Prune)	3.1 to 4.1	1.0		Use 4.1 pints per acre on trees taller than 20 feet and 3.1 to 4.1 pints per acre on smaller trees. Make 1 application at popcorn (pink, red, or early white bud) and a second application at full bloom. If weather conditions favor disease development, make an additional application at petal fall.
	Cherry Leaf Spot Peach Nectarine Apricot Scab	3.1 to 4.1	1.0		In addition to the bloom application listed above, make 1 application at shuck split. DO NOT apply Equus 720 after shuck split and before harvest. If additional disease control is needed before harvest, use another registered fungicide. For control of cherry leaf spot after harvest, make 1 application to foliage within 7 days after fruit is removed. In orchards with a history of high leaf spot incidence, make a second application 10 to 14 days later.

## **IMPORTANT**

All applicable directions, restrictions, and precautions on the Equus 720 package label are to be followed. This labeling contains supplemental instructions and restrictions for use of this product which may not appear on the package label. Follow the instructions carefully. This labeling must be in the possession of the user at the time of pesticide application.

Expiration date: This supplemental labeling expires on December 31, 2001.

Equus is a trademark of Griffin L.L.C.

Griffin and Design are a registered trademark of Griffin Corporation

© 2000

Griffin L.L.C.

Waldosta, GA 31601

GSL 080400