


ONYX



TETROSAN

3,4D

ACCEPTED

OV-18-1971

CONCENTRATED GERMICIDE

Active Ingredients

n-Alkyl (50% C₁₂, 30% C₁₁,
17% C₁₀, 3% C₉) dimethyl
3, 4 dichlorobenzyl ammonium
chlorides 60%

Inert Ingredients 40%

Total 100%

Net Contents U.S.D.A. Reg. No. 1839-12

FOR MANUFACTURING USE ONLY

ONYX CHEMICAL COMPANY

Div. of Millmaster-Onyx Corp.

190 Warren Street

Jersey City, N. J.

DANGER

Keep out of Reach of Children. See right
panel for additional precautionary
statements.

ACCEPTED WITH COMMENTS

ONYX



Tetrosan 3, 4D is a cationic germicide.

Tetrosan 3, 4D in recommended use dilutions is non-corrosive on inanimate hard surfaces.

DO NOT MIX WITH SOAP OR ANIONIC MATERIALS.

TETROSAN

3, 4D

CONCENTRATED GERMICIDE

Active Ingredients

n-Alkyl (50% C₁₂, 30% C₁₄, 17% C₁₆, 3% C₁₈) dimethyl 3, 4 dichlorobenzyl ammonium chlorides 60%

Inert Ingredients 40%

Total 100%

Net Contents U.S.D.A. Reg. No. 1839-12

FOR MANUFACTURING USE ONLY

ONYX CHEMICAL COMPANY
Div. of Millmaster-Onyx Corp.

190 Warren Street Jersey City, N.J.

DANGER

Keep out of Reach of Children. See right panel for additional precautionary statements.

DANGER
This product is extremely toxic. It is a strong irritant to the skin, eyes, nose, and throat. It is also a powerful emetic. It should be handled with extreme care. Avoid contact with skin, eyes, nose, and throat. If contact occurs, wash immediately with large quantities of water. If swallowed, drink plenty of water. If inhaled, get fresh air immediately. If necessary, call a physician for medical advice.

FIRST AID
If in contact with skin, wash immediately with large quantities of water. If in contact with eyes, flush with water for at least 15 minutes. If in contact with nose or throat, rinse with water. If swallowed, drink plenty of water. If inhaled, get fresh air immediately. If necessary, call a physician for medical advice.

NOTE TO PHYSICIAN
To avoid the unusual danger of overdosage, the use of this product should be limited to disinfecting purposes. Measures should be taken to avoid overdosage and measures to improve breathing apparatus or mask should be used. If necessary, call a physician for medical advice. In patients, convulsions may be controlled by the cautious intravenous injection of a short-acting barbiturate drug.