

ONYX



ISOTHAN

Q-75

**CONCENTRATED FUNGICIDE
FOR MANUFACTURING USE ONLY**

Active Ingredients:

**n-Alkyl (50% C₁₂, 30% C₁₄,
17% C₁₆, 3% C₁₈) isoquinolinium
bromide 75%
Isopropanol 15%**

Inert Ingredients 10%


Total 100%

Net Contents EPA Reg. No. 1839-
ONYX CHEMICAL COMPANY
Div. of Millmaster-Onyx Corp.
190 Warren Street Jersey City, N. J.

DANGER

**Keep Out of Reach of Children. See right
panel for additional precautionary
statements.**

ONYX



Isiothan Q-75 is a concentrated fungicide.

Isiothan Q-75 in recommended use dilutions is non-corrosive on inanimate hard surfaces.

FISH & WILD-LIFE CAUTION:

Do not contaminate water by cleaning of equipment, or disposal of wastes.

Rinse empty container thoroughly with water and discard.

DO NOT MIX WITH SOAP OR ANIONIC MATERIALS.

ISIOTHAN

Q-75

CONCENTRATED FUNGICIDE
FOR MANUFACTURING USE ONLY

Active Ingredients:

n-Alkyl (50% C ₁₂ , 30% C ₁₄ , 17% C ₁₆ , 3% C ₁₈) isoquinolinium bromide	75%
Isopropanol	15%

Inert Ingredients	10%
Total 100%	

Nat Contents	EPA Reg. No. 1839-
ONYX CHEMICAL COMPANY	
Div. of Millmaster-Onyx Corp.	
190 Warren Street	Jersey City, N. J.

DANGER

Keep Out of Reach of Children. See right panel for additional precautionary statements.

DANGER

Keep Out of Reach of Children. Corrosive. Causes severe eye and skin damage. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Wash at least 15 minutes. Avoid contamination of food.

FIRST AID

In case of contact, immediately flush eye or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN:

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock as well as oxygen and measures to support breathing manually or mechanically may be needed. If persistent, convulsions may be controlled by the cautious intravenous injection of a short-acting barbituate drug.