

ONYX



BTC

ACCEPTED

9-9-72

1839-23

THE FEDERAL REGISTER
THE ADMINISTRATIVE AND
REGULATORY INFORMATION
SECTION

824

CONCENTRATED GERMICIDE

Active ingredients:

n-Alkyl (60% C₁₂, 30% C₁₄, 5%

C₁₆, 5% C₁₈) dimethyl benzyl

ammonium chlorides

50%

Inert ingredients

50%

TOTAL

100%

Net Contents:

EPA Reg. No. 1839-23

FOR MANUFACTURING USE ONLY

ONYX CHEMICAL COMPANY

Div. of Millmaster-Onyx Corp.


190 Warren St.

Jersey City, N. J.

DANGER

Keep Out of Reach of Children. See right panel for additional precautionary statements.

ONYX



BTC

824

AC
0-9-72
1939-23
ED UNL

CONCENTRATED GERMICIDE

Active ingredients:

n-Alkyl (60% C , 30% C , 5%
C , 5% C) dimethyl benzyl
ammonium chlorides 50%

Inert ingredients 50%

TOTAL 100%

Net Contents: EPA Reg. No. 1839-23

FOR MANUFACTURING USE ONLY

ONYX CHEMICAL COMPANY

Div. of Millmaster-Onyx Corp.

190 Warren St.

Jersey City, N. J.

BTC-824 is a cationic germicide.

BTC-824 in recommended use dilutions is non-corrosive to inanimate surfaces.

Rinse empty container thoroughly with water and discard.

DO NOT MIX WITH SOAP OR ANIONIC MATERIALS.

MAY 1972

DANGER

Keep Out of Reach of Children. See right panel for additional precautionary statements.

DANGER

Keep Out of Reach of Children. Corrosive. Causes severe eye and skin damage. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

FIRST AID

In case of contact, immediately flush eye or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN:

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock as well as oxygen and measures to support breathing manually or mechanically may be needed. If persistent, convulsions may be controlled by the cautious intravenous injection of a short-acting barbiturate drug.