


ONYX



ONYXIDE

200

Metal Cutting Fluid Preservative

FOR MANUFACTURING USE ONLY

Active Ingredients:

hexahydro—1,3,5—tris 78.5%
(2—hydroxyethyl)—s-triazine

Inert Ingredients 21.5%

TOTAL 100%

Net Contents

EPA Reg. No. 1839-59

ONYX CHEMICAL COMPANY

Division of Millmaster Onyx Corporation
190 Warren Street Jersey City, N.J. 07302

DANGER

Keep Out of Reach of Children. See right
panel for additional precautionary
statements.

Oct-18-1979


UNDER THE FEDERAL INSECTICIDE ACT
FOR A PESTICIDE WHICH IS REGISTERED
UNDER NO. 1839-59

Comment

ACCEPTED WITH COMMENTS

ONYX

ONYXIDE



ACCEPTED

Oct 18-1972

1839-59

1839-59

200

Metal Cutting Fluid Preservative

FOR MANUFACTURING USE ONLY

Active Ingredients:

hexahydro—1,3,5—tris 78.5%
(2—hydroxyethyl)—s-triazine

Inert Ingredients

21.5%

TOTAL 100%

Net Contents

EPA Reg. No. 1839-59

ONYX CHEMICAL COMPANY

Division of Millmaster Onyx Corporation
190 Warren Street Jersey City, N.J. 07310

DANGER

Keep Out of Reach of Children. See right panel for additional precautionary statements.

DANGER

Keep Out of Reach of Children. Corrosive. Causes severe eye and skin damage. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN:

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock as well as oxygen and measures to support breathing manually or mechanically may be needed. If persistent, convulsions may be controlled by the cautious intravenous injection of a short-acting barbiturate drug.

See Technical Bulletin for Use Direction.

JUNE, 1972