

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Veterinarian Type Disinfectant is designed specifically for use on walls, floors, operating tables, cages, crates, feeding and watering equipment and furniture in veterinarian hospitals, laboratories, kennels, pet shops, zoos and pet animal quarters.

**DISINFECTION, DEODORIZATION, AND CLEANING:**

For disinfecting, deodorization and cleaning, add 1 ounce of VTD to one gallon of water and apply solution with a mop, cloth, sponge or mechanical sprayer so as to wet all surfaces thoroughly. Allow to remain wet for 10 minutes and then let air dry.

For heavily soiled or contaminated area, a pre-cleaning step is required. Prepare a fresh solution for each use.

For feeding and water equipment, RINSE WITH POTABLE WATER after application.

**Mildewstat**- To control mold and mildew on pre-cleaned hard non-porous surfaces (such as floors, walls, table tops) add 1-oz. VTD to 1 gallon of water. Apply solution with a cloth, mop, or sponge making sure to wet all surfaces completely. Let air dry. Prepare a fresh solution for each use. Repeat application at weekly intervals or when mildew growth reappears.

**PRECAUTIONARY STATEMENTS  
HAZARD TO HUMAN AND DOMESTIC ANIMALS**

**DANGER**

Keep Out of Reach of Children. Causes severe eye and skin damage. Do not get in eyes, skin or on clothing. Harmful if swallowed. Wear goggles or face shield and rubber gloves when handling. Avoid contamination of food. Remove and wash contaminated clothing.

**STATEMENT OF PRACTICAL TREATMENT**

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before reuse. If swallowed, drink milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Call a physician.

**NOTE TO PHYSICIAN**

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

**ACCEPTED**

**JUL 01 1987**

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 1839-100



**VETERINARIAN TYPE  
DISINFECTANT**

**detergent  
virucide\***

**deodorant  
mildewstat**

**ACTIVE INGREDIENTS:**

- n-Alkyl (60XC<sub>14</sub>, 30XC<sub>16</sub>, 5XC<sub>12</sub>, 5XC<sub>18</sub>) dimethyl benzyl ammonium chlorides.....4.50Z
- n-Alkyl (68XC<sub>12</sub>, 32XC<sub>16</sub>) dimethyl ethylbenzyl ammonium chlorides.....4.50Z
- INERT INGREDIENTS:.....91.00Z
- TOTAL 100.00Z

EPA REG. No. 1839-100  
EPA EST. No. 1839-11-01

**ONLY FOR SALE TO, USE, AND STORAGE BY SERVICE PERSONS  
KEEP OUT OF REACH OF CHILDREN**

**DANGER**

See side panel for additional precautionary and practical treatment statements.

**Stepan S**

**Stepan Company  
Northfield, IL 60093**

NET CONTENTS

1071  
Pm 31  
1839-100

Veterinarian Disinfectant is designed for Veterinarian hospitals, laboratories, kennels, pet shops, zoos, and pet animal quarters. When used as directed, VTD is formulated to disinfect, deodorize and clean inanimate hard surfaces (such as walls, floors, sink tops, kennel runs, operating tables, cages, feeding and watering equipment and furniture).

In addition, Veterinarian Disinfectant deodorizes those areas which generally are hard to keep fresh smelling such as garbage storage areas, empty garbage bins and cans, and any other areas which are prone to odors caused by microorganisms.

**DISINFECTION:** When used as directed, Veterinarian Disinfectant demonstrates effective disinfection against the following:

- Pseudomonas aeruginosa*
- Escherichia coli*
- Klebsiella pneumoniae*
- Salmonella schultzei*
- Salmonella choleraesuis*
- Streptococcus faecalis*
- Shigella dysenteriae*
- Enterobacter aerogenes*
- Staphylococcus aureus*

**HOSPITAL DISINFECTIONS:** When used as directed,

Veterinarian Disinfectant is an effective disinfectant against *Pseudomonas aeruginosa* in animals hospitals and laboratories for operating tables and equipment.

**Virucidal\* Activity:** The product when used at general disinfection dilution rates on environmental inanimate hard surfaces exhibits effective virucidal activity against Canine Distemper, Pseudorabies virus, Vaccinia virus, Influenza A<sub>2</sub>, Herpes simplex and Adeno virus Type 2, and Canine Parvovirus. Efficacy tests have demonstrated that Veterinarian Disinfectant is an effective bactericide, fungicide, and virucide\* in the presence of organic soil (5% blood serum).

**Mold and Mildewstat-** Veterinarian Disinfectant prevents and controls mold and mildew and the odors they cause on hard non-porous inanimate surfaces.

**STORAGE AND DISPOSAL**

**DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE AND DISPOSAL.**

**STORAGE:** Store in a dry place no lower in temperature than 50°F or higher than 120°F.

**CONTAINER DISPOSAL** - Do not reuse empty container. Triple rinse empty container with water. Return metal drum then offer for reconditioning or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Plastic containers may be disposed of in a sanitary landfill, incinerated or, if allowed by local authorities, by burning. If burned, stay out of smoke.

**PESTICIDE DISPOSAL** - Pesticide waste are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.