

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

Stepan Company
22 West Frontage Rd.
Northfield, IL 60093

JUL 23 2001

Subject: BTC 2125M 20% Solution
EPA Registration No. 1839-155
Received Date of January 24, 2001

Attn: Tom Isaac
Sr. Regulatory Specialist

The amendment referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable.

A stamped copy of the label is enclosed for your records. Submit one (1) copy of the final printed label before this product is released for shipment.

If you have any questions concerning the comments in this letter, you should contact Zenobia Jones at (703) 308-6198.

Sincerely,



Velma Noble
Product Manager 31
Regulatory Management Branch 1
Antimicrobial Division (7510C)

CONCURRENCES

SYMBOL	7510C						
SURNAME	Z. Jones						
DATE	7-20-01						

276

BTC 2125[®]M 20% Solution

DISINFECTANT	SANITIZER
DEODORIZER	*VIRUCIDE
MILDEWSTAT (on hard inanimate surfaces)	
LAUNDRY BACTERIOSTAT	

ACTIVE INGREDIENTS

Alkyl (60% C ₁₄ , 30% C ₁₆ , 5% C ₁₂ , 5% C ₁₈) dimethyl benzyl ammonium chlorides.....	10.0%
Alkyl (68% C ₁₂ , 32% C ₁₄) dimethyl ethylbenzyl ammonium chlorides.....	10.0%
INERT INGREDIENTS	80.0%
TOTAL	100.0%

KEEP OUT OF REACH OF CHILDREN

DANGER

See side panel for additional precautionary statements.

EPA Reg. No. 1839-155
EPA Est. No. 1839-IL-01

ACCEPTED
with COMMENTS
in EPA Letter Dated:

JUL 23 2001

STEPAN COMPANY
22 West Frontage Road • Northfield, IL 60093

Under the Federal Insecticide,
Fungicide, and Rodenticide Act as
amended, for the pesticide,
registered under EPA Reg. No. **1839-155**

NET CONTENTS

(Middle panel)

Approved 6/27/2000

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

BTC 2125™M 20% Solution is designed specifically for hospitals, food processing plants, dairies, restaurants, bars, animal quarters, poultry and turkey farms, kennels and institutions where disinfection, sanitization, and deodorization is of prime importance.

When used as directed, this product is formulated to disinfect precleaned inanimate, hard surfaces such as walls, floors, sink tops, tables, chairs, and telephones.

This product deodorizes those areas that generally are hard to keep fresh smelling, such as garbage storage areas, empty garbage bins and cans, pet areas and any other areas that are prone to odors caused by micro-organisms.

In addition, this product will sanitize previously cleaned and rinsed non-porous food contact surfaces such as tanks, chopping blocks, counter tops, drinking glasses and eating utensils. This product can also be used to sanitize previously cleaned food grade eggs in shell egg and egg product processing plants.

This product is not to be used as a terminal sterilant/high-level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body.

DISINFECTION - To disinfect pre-cleaned, inanimate, hard surfaces apply this product with mop, cloth, sponge or hand-pump trigger sprayer so as to wet all surfaces thoroughly. Allow to remain wet for 10 minutes, then remove excess liquid. For disinfection, a pre-cleaning step is required. Prepare a fresh solution for each use.

DILUTION RATES:

GENERAL DISINFECTION - Add 1.5 ounces of this product per 5 gallons of water for disinfection against *Staphylococcus aureus*, *Salmonella choleraesuis*, *Listeria monocytogenes*, and *Yersinia enterocolitica*.

HOSPITAL DISINFECTION - Add 1.75 ounces of this product per 5 gallons of water for disinfection against *Pseudomonas aeruginosa*.

DISINFECTION OF POULTRY EQUIPMENT, ANIMAL QUARTERS, AND KENNELS - For disinfection of pre-cleaned poultry equipment (brooders, watering founts, feeding equipment), animal quarters and kennels, apply a solution of 1.75 ounces of this product per 5 gallons of water. Remove all poultry, animals, and feed from premises, trucks, coops, and crates. Remove all litter and droppings from floors, walls and surfaces of facilities occupied or traversed by poultry or animals. Empty all troughs, racks, and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate the surfaces with the disinfecting solution for a period of 10 minutes. Ventilate building, coops, and other closed spaces. Do not house poultry or animals or employ equipment until treatment has been absorbed, set or dried. All treated equipment that will contact feed or drinking water must be rinsed with potable water before reuse.

KILLS HIV-1 ON LAUNDRY IN COMMERCIAL, INDUSTRIAL, AND INSTITUTIONAL APPLICATIONS. This product provides virucidal activity against HIV-1 on fabrics when added to the final rinse at a rate of 3.0 ounces per 100 pounds of dry laundry.

LAUNDRY BACTERIOSTAT FOR COMMERCIAL, INDUSTRIAL, AND NON-MEDICAL INSTITUTIONAL LAUNDRY APPLICATIONS: This product provides the fabric with residual bacteriostatic activity against odor causing gram-negative and gram-positive bacteria when this product is added to the final rinse at a rate of 4.8 ounces per 100 lbs. of dry laundry.

Stepan Company
Northfield, IL 60093
EPA Reg. No. 1839-155

ACCEPTED
with COMMENTS
in EPA Letter Dated:
(Left panel)

Approved 6/27/2000

JUL 23 2001

Under the Federal Insecticide,
Fungicide, and Rodenticide Act as
amended, for the pesticide,
registered under EPA Reg. No. 1839-155

SANITIZATION - Recommended for use in restaurants, dairies, food processing plants, and bars. When used as directed this product is an effective sanitizer against *Escherichia coli* and *Staphylococcus aureus*.

Remove all gross food particles and soil from areas that are to be sanitized with a good detergent, pre-flush, pre-soak or pre-scrape treatment. Rinse with a potable water rinse.

To sanitize pre-cleaned and potable water-rinsed, non-porous food contact surfaces, prepare a 200 ppm active quaternary solution by adding 0.5 ounce of this product to 4 gallons of water, 1.25 ounces to 10 gallons of water, or 2.5 ounces to 20 gallons of water.

To sanitize immobile items such as tanks, chopping blocks and counter tops, flood the area with a 200 ppm active quaternary solution for at least 60 seconds, making sure to wet all surfaces completely. Remove, drain the use solution from the surface, and air dry. Prepare a fresh solution daily or more frequently as soil is apparent.

To sanitize mobile items such as drinking glasses and eating utensils, immerse in a 200 ppm active quaternary solution for at least 60 seconds, making sure to immerse completely. Remove, drain the use solution from the surface, and air dry. Prepare a fresh solution daily or more frequently as soil is apparent.

When used for sanitization of previously cleaned food equipment or food contact items, limit the active quaternary to 200 ppm. **NO POTABLE WATER RINSE IS REQUIRED.** This product is an effective sanitizer when diluted in water up to 750 ppm hardness (CaCO_3).

To sanitize previously cleaned food-grade eggs in shell egg and egg product processing plants, spray with a solution of 0.5 ounce of this product in 4 gallons of warm water (200 ppm active quaternary solution). The solution should be warmer than the eggs, but not to exceed 130°F. Wet eggs thoroughly and allow to drain. Eggs sanitized with this product shall be subjected to **POTABLE WATER RINSE ONLY IF THEY ARE BROKEN IMMEDIATELY** for use in the manufacture of egg products. Eggs should be dry before casing or breaking. The solution should not be reused for sanitizing eggs.

*** VIRUCIDAL ACTIVITY** - This product, when used on environmental, inanimate hard surfaces at 1.75 ounces per 5 gallons of water exhibits effective virucidal activity against Influenza A/Japan (representative of the common flu virus), Herpes Simplex Type 1 (causative agent of fever blisters), Adenovirus Type 5 (causative agent of upper respiratory infections), Vaccinia virus (representative of the pox virus group), Avian Influenza A/Turkey/Wisconsin (causative agent of an acute avian lower respiratory tract infection), Newcastle disease virus, Laryngotracheitis Virus, and HIV-1 (Human Immunodeficiency Virus Type 1).

KILLS HIV-1 ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS in health care settings (hospitals, nursing homes) or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

PERSONAL PROTECTION: When handling items soiled with blood or body fluids use disposable latex gloves, gowns, masks, or eye coverings.

CLEANING PROCEDURES: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product.

CONTACT TIME: Allow surface to remain wet for 10 minutes.

DISPOSAL OF INFECTIOUS MATERIALS: Blood and other body fluids should be autoclaved and disposed of according to local regulations for infectious waste disposal.

Stepan Company
Northfield, IL 60093
EPA Reg. No. 1839-155

ACCEPTED
with COMMENTS
in EPA Letter Dated:

(Left panel continued) JUL 23 2001

Approved 6/27/2000

Under the Federal Insecticide,
Fungicide, and Rodenticide Act as
amended, for the pesticide,
registered under EPA Reg. No. 1839-155

DEODORIZATION - To deodorize, apply this product as indicated under the heading **GENERAL DISINFECTION**.

MILDEWSTAT - To control mold and mildew on pre-cleaned, hard, non-porous surfaces add 1.5 ounces of this product per 5 gallons of water. Apply solution with a cloth, mop or sponge making sure to wet all surfaces completely. Let air dry. Prepare a fresh solution for each use. Repeat application at weekly intervals or when mildew growth appears.

EFFICACY TESTS HAVE DEMONSTRATED THAT THIS PRODUCT IS AN EFFECTIVE BACTERICIDE AND VIRUCIDE IN THE PRESENCE OF ORGANIC SOIL (5% BLOOD SERUM).

ACCEPTED
with COMMENTS
in EPA Letter Dated:

JUL 23 2001

Under the Federal Insecticide,
Fungicide, and Rodenticide Act as
amended, for the pesticide,
registered under EPA Reg. No. 1839-155

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS.**

DANGER

KEEP OUT OF REACH OF CHILDREN. **CORROSIVE.** Causes severe eye and skin damage. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food. Remove contaminated clothing and wash before reuse. Wash thoroughly with soap and water after handling.

STATEMENT OF PRACTICAL TREATMENT

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. If swallowed, promptly drink large quantities of water. Do not induce vomiting. Call a physician immediately.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact your State Water Board or Regional Office of the EPA.

Contains no phosphorus.

STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL.

STORAGE - Store in a dry place no lower in temperature than 50°F or higher than 120°F.

CONTAINER DISPOSAL - Do not reuse empty container. Triple rinse empty container with water. Return metal drum then offer for reconditioning or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers may be disposed of in a sanitary landfill, incinerated, or if allowed by local authorities, by burning. If burned stay out of smoke.

For containers 1 gallon or less: Do not reuse empty container (bottle, can, bucket). Wrap container and put in trash.

PESTICIDE DISPOSAL - Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

ACCEPTED
with COMMENTS
in EPA Letter Dated:

JUL 23 2001

Under the Federal Insecticide,
Fungicide, and Rodenticide Act as
amended, for the pesticide,
registered under EPA Reg. No. 1839-155

Stepan Company
Northfield, IL 60093
EPA Reg. No. 1839-155

(Right panel)

Approved 6/27/2000