

1839-167

9-13-2000

## UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

1/5

Stepan Company  
22 West Frontage Road  
Northfield, IL 60093

SEP 13 2000

ATTENTION: Bruce L. Greene

SUBJECT: BTC® 885 Neutral Disinfectant Cleaner-256  
EPA Registration Number: 1839-167  
Your Amendment Dated: June 7, 2000  
EPA Received Date: June 20, 2000

The amendment referred to above, submitted in connection with registration under FIFRA sec. 3(c)(7)(A), is acceptable provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. Make the following labeling change before you release the product for shipment bearing the amended labeling: Revise the second section on the left panel-1 by changing "porcelain" to read "glazed porcelain".
3. Submit two (2) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, this registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of this product bearing the amended labeling constitutes acceptance of these conditions. A stamped copy of the labeling is enclosed for your records. Should you have any questions concerning this letter, please contact Tracy Lantz at (703) 308-6415.

Sincerely,



Velma Noble

Product Manager (31)

Regulatory Management Branch I

Antimicrobials Division (7510C)

Enclosure (stamped label)

7510C:T.Lantz:9/12/00:1839-167a

## CONCURRENCES

SYMBOL	7510C						
SURNAME	J. Lantz						
DATE	9/12/00						

EPA Form 1320-1A (1/90)

Printed on Recycled Paper

OFFICIAL FILE COPY

2/5

# BTC<sup>®</sup> 885

## Neutral Disinfectant Cleaner-256

CLEANER	DISINFECTANT
DETERGENT	FUNGICIDE (against pathogenic fungi)
DEODORIZER	* VIRUCIDE
MILDEWSTAT (on hard, non-porous, inanimate surfaces)	
CITRUS CANCKER CONTROL	MUSHROOM FARM SANITATION

<b>ACTIVE INGREDIENTS</b>	
Octyl decyl dimethyl ammonium chloride .....	6.510%
Dioctyl dimethyl ammonium chloride .....	3.255%
Didecyl dimethyl ammonium chloride .....	3.255%
Alkyl (50% C <sub>14</sub> , 40% C <sub>12</sub> , 10% C <sub>16</sub> ) dimethyl benzyl ammonium chloride.....	8.680%
<b>INERT INGREDIENTS</b> .....	<u>78.300%</u>
<b>Total</b>	<b>100.000%</b>

**KEEP OUT OF REACH OF CHILDREN**  
**DANGER**

See side panels for additional precautionary statements.

EPA REG. NO. 1839-167  
EPA EST. NO. 1839-IL-01

STEPAN COMPANY  
22 West Frontage Road • Northfield, IL 60093


ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

SEP 13 2008

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act as  
amended, for the pesticide,  
registered under EPA Reg. No. **1839-167**

NET CONTENTS

(middle panel)



Amend 05/27/00

3/5

## DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

BTC® 885 Neutral Disinfectant Cleaner-256 is a phosphate-free formulation designed to provide effective cleaning, deodorizing, and disinfection specifically for hospitals, nursing homes, schools, food processing plants, food service establishments, transportation terminals, office buildings, manufacturing facilities, lodging establishments, retail business, and athletic/recreational facilities where housekeeping is of prime importance in controlling the hazard of cross-contamination. For use in federally inspected meat and poultry processing plants.

This product, when used as directed, is formulated to disinfect hard non-porous inanimate environmental surfaces such as floors, walls, metal surfaces, stainless steel surfaces, porcelain, glazed ceramic tile, plastic surfaces, vanity tops, shower stalls, bathtubs, and cabinets. For larger areas: operating rooms, patient care facilities, and restrooms, this product is designed to provide both general cleaning and disinfecting.

This product deodorizes those areas which generally are hard to keep fresh smelling, such as garbage storage areas, empty garbage bins and cans, toilet bowls, and other areas which are prone to odors caused by microorganisms.

This product is formulated to a neutral pH and will not dull high-gloss floor finishes with repeated use.

This product is not to be used as a terminal sterilant/high-level disinfectant on any surface or instrument that (1) is introduced directly into the human body either into or in contact with bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body.

**GENERAL CLEANING** – Apply product to soiled area with a mop, cloth, sponge, or hand pump trigger sprayer. Product can be diluted at ½ ounce per gallon or applied at full strength. Thoroughly wet soiled surface and scrub as necessary. Rinse surface with potable water and wipe dry with a dry paper towel or lint-free cloth.

**DISINFECTION** – To disinfect pre-cleaned, inanimate, hard, non-porous surfaces add ½ ounce of this product per gallon of water. Apply solution with a mop, cloth, sponge, or hand-pump trigger sprayer so as to wet all surfaces thoroughly. Allow to remain wet for 10 minutes, then remove excess liquid.

For heavily soiled areas, a pre-cleaning step is required. Prepare a fresh solution for each use.

**TO DISINFECT FOOD PROCESSING PREMISES:** floors, walls, and storage areas, add ½ ounce of this product per gallon of water. For heavily soiled areas, a pre-cleaning step is required. Apply solution with a mop, cloth, sponge or hand pump trigger sprayer so as to wet all surfaces thoroughly. Allow to remain wet for 10 minutes, then remove excess liquid. Before using this product, food products and packaging materials must be removed from the area or carefully protected. All surfaces which may come in contact with food must be rinsed with potable water.

**TO DISINFECT FOOD SERVICE ESTABLISHMENT FOOD CONTACT SURFACES:** countertops, appliances, tables, add ½ ounce of this product per gallon of water. For heavily soiled areas, a pre-cleaning step is required. Apply solution with a cloth, sponge or hand pump trigger sprayer so as to wet all surfaces thoroughly. Allow the surface to remain wet for 10 minutes, then remove excess liquid and rinse the surface with potable water. This product cannot be used to clean the following food contact surfaces: utensils, glassware and dishes.

**TO DISINFECT TOILET BOWLS** – Prior to disinfection, remove gross filth or soils from surfaces with bowl brush. Add ½ ounce of product to the bowl water. Brush or swab the bowl completely using a scrub brush or toilet mop making sure to get under the rim. Let stand for 10 minutes and flush.

**DEODORIZATION** – To deodorize, apply this product as indicated under the heading DISINFECTION.

**MILDEWSTAT** – To control mold and mildew (such as *Aspergillus niger*, ATCC 6275) and the odors they cause on pre-cleaned, hard, non-porous inanimate surfaces add ½ ounce of this product per gallon of water. Apply solution with a cloth, mop or sponge making sure to wet all surfaces completely. Let air dry. Prepare a fresh solution for each use. Repeat application at weekly intervals or when mildew growth appears.

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

SEP 13 2008

Stepan Company  
Northfield, IL 60093  
EPA Reg. No. 1839-167

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act as  
amended, for the pesticide,  
registered under EPA Reg. No. 1839-167

(left panel-1)

Amend 06/27/00

4/5

**BACTERICIDAL ACTIVITY** – At the ½ ounce per gallon dilution, this product demonstrates effective disinfectant activity against the organisms: *Pseudomonas aeruginosa* PRD-10, *Salmonella choleraesuis*, *Staphylococcus aureus*, *Staphylococcus aureus* (clinical isolate), *Bordetella bronchiseptica*, *Corynebacterium ammoniagenes*, *Enterobacter aerogenes*, *Enterobacter cloacae*, *Enterobacter cloacae* (clinical isolate), *Enterococcus faecalis*, *Enterococcus faecalis* (clinical isolate), *Escherichia coli*, *Escherichia coli* (clinical isolate), *Fusobacterium necrophorum*, *Klebsiella pneumoniae* subsp. *pneumoniae*, *Lactobacillus casei* subsp. *rhamnosus*, *Listeria monocytogenes*, *Pasteurella multocida*, *Proteus vulgaris*, *Proteus mirabilis* ATCC 9921, *Proteus mirabilis* ATCC 25933, *Salmonella choleraesuis* subsp. *choleraesuis* serotype paratyphi B, *Salmonella choleraesuis* subsp. *choleraesuis* serotype typhi, *Salmonella choleraesuis* subsp. *choleraesuis* serotype typhimurium, *Salmonella choleraesuis* subsp. *choleraesuis* serotype pullorum, *Serratia marcescens*, *Shigella sonnei*, *Shigella flexneri* Type 2b, *Shigella dysenteriae*, *Staphylococcus aureus* subsp. *aureus*, *Methicillin* resistant *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Staphylococcus epidermidis* (clinical isolate), *Streptococcus pyogenes* (Clinical - Flesh Eating Strain BIRD M3), *Streptococcus pyogenes* Group A, *Xanthomonas maltophilia* (clinical isolate), and *Vancomycin* resistant *Enterococcus faecalis* (ATCC 51575).

**FUNGICIDAL ACTIVITY** – At the ½ ounce per gallon dilution, this product is fungicidal against the pathogenic fungi, *Trichophyton mentagrophytes* (the fungus which causes Athlete's Foot) and *Candida albicans*. Apply solution with a cloth, sponge or hand pump trigger sprayer to hard, non-porous surfaces found in bathrooms, shower stalls, locker rooms, exercise facilities or other clean, hard non-porous surfaces commonly contacted by bare feet. Allow the surface to remain wet for 10 minutes, then remove excess liquid. Diluted product should be applied daily or more frequently with heavy facility use.

\* **VIRUCIDAL ACTIVITY** – This product when used on environmental, inanimate hard non-porous surfaces at ½ ounce per gallon of water exhibits effective virucidal activity against Human Immunodeficiency Virus Type 1 (HIV-1), Human Immunodeficiency Virus Type 2 (HIV-2), Herpes Simplex Type 1 (causative agent of fever blisters), Herpes Simplex Type 2 (genital disease), Influenza A<sub>2</sub>/Hong Kong, Vaccinia, Pseudorabies, Bovine Rhinotracheitis, Feline Leukemia, Feline Parvovirus, and Canine Distemper.

**KILLS HIV-1 and HIV-2 ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings (Hospitals, Nursing Homes) or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 or Type 2 (HIV-1 or HIV-2) (associated with AIDS).

**SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 and HIV-2 OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.**

**PERSONAL PROTECTION:** When handling items soiled with blood or body fluids, use disposable latex gloves, gowns, masks, or eye coverings.

**CLEANING PROCEDURES:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product.

**CONTACT TIME:** Allow surface to remain wet for 10 minutes.

**DISPOSAL OF INFECTIOUS MATERIALS:** Blood and other body fluids should be autoclaved and disposed of according to local regulations for infectious waste disposal.

**MUSHROOM FARM SANITATION** – This product disinfects, cleans, and deodorizes in one step. Recommended for use in mushroom farm sanitation. Apply to surfaces which do not come in contact with the crop, compost or casing. Remove gross filth and heavy soil deposits using a shovel, broom or vacuum depending on the area to be disinfected. For heavily soiled areas, a pre-cleaning step is required. Apply this product at ½ ounce per gallon of water dilution (1:256), thoroughly wetting the surfaces with a mop, sponge, cloth, spraying or soaking. Treated surfaces should remain wet for 10 minutes and let air dry. The recommended use solution is prepared fresh for each use, then discarded. If greater detergency is required, dilute this product at 1 to 2 ounces per gallon of water. **DO NOT APPLY TO MUSHROOM CROP, COMPOST OR CASING.**

**FOR FOOT DIP OF WATERPROOF FOOTWEAR** – Use this product at ½ ounce per gallon of water in foot dip tray. Shoe baths must contain at least 1 inch of freshly made solution and should be placed at the entrances to buildings. Scrape shoes and place in diluted solution for 60 seconds before entering building. Change solution in bath daily or sooner if solution appears dirty.

**BACTERICIDAL ACTIVITY** – When diluted at ½ ounce per gallon, this product demonstrates effective disinfectant activity against the following mushroom pathogens: *Pseudomonas tolaasii* (Blotch Disease) and *Pseudomonas* sp. (Mummy Disease).

**FUNGICIDAL ACTIVITY** – When diluted at ½ ounce per gallon, this product demonstrates effective fungicidal activity against the following mushroom pathogens: *Verticillium fungicola* (Verticillium Disease, dry bubble, brown spot or fungus spot), *Mycogone perniciosa* (Mycogone Disease or wet bubble), *Tricoderma viride*, *Dactylium dendroides* (Hypomyces Disease, cobweb or soft decay), and *Tricoderma harzianum*.

Stepan Company  
Northfield, IL 60093  
EPA Reg. No. 1839-167

ACCEPTED  
with Conditions  
in EPA Letter dated  
SEP 13 2000  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act as  
amended, for the pesticide,  
registered under EPA Reg. No.  
1839-167

(left panel-2)

Amend 06/27/00

5/5

**CITRUS CANCKER CONTROL** – For prevention of Citrus Canker Disease through treatment of equipment. This product, when used at a 1:108 dilution (2000 ppm active quaternary) or 1.18 ounces (35 ml) per gallon of water, is effective against *Xanthomonas axonopodis* pathovar *citri* (Citrus Canker). Treat all trucks, vehicles, and equipment thoroughly at the 1:108 dilution. Treatment can be applied to outer clothing, shoes, trucks, attached trailers, field harvesting equipment; including cargo area, wheels, tires, under carriage, hood, roof, fenders and any other part of transportation equipment that can be taken into infested areas. To prevent the spread of Citrus Canker by this artificial means of transportation, treatments are made by trigger spraying, dipping or brushing, using the maximum rate of 2000 ppm quaternary. After use, all surfaces which come in contact with food or crop must be rinsed with potable water. Clothing should be either thoroughly rinsed or laundered before reuse. Footwear should be rinse before reuse.

**EFFICACY TESTS HAVE DEMONSTRATED THAT THIS PRODUCT IS AN EFFECTIVE BACTERICIDE, FUNGICIDE, AND VIRUCIDE IN WATER UP TO 400 PPM HARDNESS (AS CaCO<sub>3</sub>) IN THE PRESENCE OF ORGANIC SOIL (5% BLOOD SERUM).**

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**  
**DANGER**

KEEP OUT OF REACH OF CHILDREN. Corrosive. Causes irreversible eye damage and skin burns. May be fatal if absorbed through the skin. Do not get in eyes, on skin or on clothing. Wear protective eyewear (goggles or face shield), clothing, and gloves when handling. Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking or using tobacco. Remove contaminated clothing and wash clothing before reuse.

**STATEMENT OF PRACTICAL TREATMENT**

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

IF IN EYES: Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention.

IF SWALLOWED: Call a doctor or get medical attention. Do not induce vomiting or give anything by mouth to an unconscious person. Drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol.

**NOTE TO PHYSICIAN**

Probable mucosal damage may contraindicate the use of gastric lavage.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

**STORAGE AND DISPOSAL**

DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL.

**PESTICIDE STORAGE** – Store in a dry place no lower in temperature than 50°F or higher than 120°F.

**CONTAINER DISPOSAL** – Do not reuse empty container. Triple rinse empty container with water. Return metal drum or offer for reconditioning or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers may be disposed of in a sanitary landfill, incinerated, or if allowed by local authorities, by burning. If burned, stay out of smoke.

For containers 1 gallon or less: Do not reuse empty container. Wrap in newspaper and discard in trash.

**PESTICIDE DISPOSAL** – Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

SEP 13 2008

(right panel)

Stepan Company  
Northfield, IL 60093  
EPA Reg. No. 1839-167

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act as  
amended, for the pesticide,  
registered under EPA Reg. No. 1839-167

Amend 06/27/07