

# Stepan S

Stepan Company  
Northfield, IL 60093

## BTC<sup>®</sup> 885 Neutral Disinfectant Cleaner-32

CLEANER  
DETERGENT  
DEODORIZER

DISINFECTANT  
\* VIRUCIDE

### ACTIVE INGREDIENTS

Octyl decyl dimethyl ammonium chloride . . . . .	0.814%
Dioctyl dimethyl ammonium chloride . . . . .	0.407%
Didecyl dimethyl ammonium chloride . . . . .	0.407%
Alkyl (50% C <sub>14</sub> , 40% C <sub>12</sub> , 10% C <sub>16</sub> ) dimethyl benzyl ammonium chloride . . . . .	1.085%
INERT INGREDIENTS . . . . .	<u>97.287%</u>

TOTAL 100.000%


EPA REG. NO. 1839-168  
EPA EST. NO. 1839-IL-01

KEEP OUT OF REACH OF CHILDREN

# DANGER

See side panels for additional precautionary statements.

NET CONTENTS  
(middle panel)



## DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

BTC® 885 Neutral Disinfectant Cleaner-32 is a phosphate-free formulation designed to provide effective cleaning, deodorizing, and disinfection specifically for hospitals, nursing homes, schools, and other institutions where housekeeping is of prime importance in controlling cross-contamination from treated surfaces.

This product, when used as directed, is formulated to disinfect inanimate hard non porous surfaces such as walls, floors, sink tops, toilet bowls, tables, chairs, telephones, and bed frames. For larger areas such as operating rooms, patient care facilities, and restrooms, this product is designed to provide both general cleaning and disinfecting.

This product deodorizes those areas which generally are hard to keep fresh smelling, such as garbage storage areas, empty garbage bins and cans, toilet bowls, and other areas which are prone to odors caused by microorganisms.

**DISINFECTION** - To disinfect inanimate, hard non porous surfaces add 4 ounces of this product per gallon of water. Apply solution with a mop, cloth, sponge, or hand-pump trigger sprayer so as to wet all surfaces thoroughly. Allow to remain wet for 10 minutes, then remove excess liquid.

For heavily soiled areas, a pre-cleaning step is required. Prepare a fresh solution for each use.

To disinfect toilet bowls, flush toilet, add 4 ounces of this product directly to the bowl water. Swab the bowl completely using a scrub brush or toilet mop, making sure to get under the rim. Let stand for 10 minutes and flush.

**EFFICACY TESTS HAVE DEMONSTRATED THAT THIS PRODUCT IS AN EFFECTIVE BACTERICIDE AND VIRUCIDE IN WATER UP TO 400 PPM HARDNESS (AS  $\text{CaCO}_3$ ) IN THE PRESENCE OF ORGANIC SOIL (5% BLOOD SERUM).**

**DEODORIZATION** - To deodorize, apply this product as indicated under the heading **DISINFECTION**.


### **BACTERICIDAL ACTIVITY -**

Disinfection: *Pseudomonas aeruginosa* PRD-10, *Salmonella choleraesuis*, *Staphylococcus aureus*, *Staphylococcus aureus* (clinical isolate), *Bordetella bronchiseptica*, *Corynebacterium ammoniagenes*, *Enterobacter aerogenes*, *Enterobacter cloacae*, *Enterobacter cloacae* (clinical isolate), *Enterococcus faecalis*, *Enterococcus faecalis* (clinical isolate), *Escherichia coli*, *Escherichia coli* (clinical isolate), *Fusobacterium necrophorum*, *Klebsiella pneumoniae* subsp. *pneumoniae*, *Lactobacillus casei* subsp. *rhannosus*, *Listeria monocytogenes*, *Pasteurella multocida*, *Proteus vulgaris*, *Proteus mirabilis* ATCC 9921, *Proteus mirabilis* ATCC 25933, *Salmonella choleraesuis* subsp. *choleraesuis* serotype *paratyphi* B, *Salmonella choleraesuis* subsp. *choleraesuis* serotype *typhi*, *Salmonella choleraesuis* subsp. *choleraesuis* serotype *typhimurium*, *Salmonella choleraesuis* subsp. *choleraesuis* serotype *pullorum*, *Serratia marcescens*, *Shigella sonnei*, *Shigella flexneri* Type 2b, *Shigella dysenteriae*, *Staphylococcus aureus* subsp. *aureus*, *Staphylococcus epidermidis*, *Staphylococcus epidermidis* (clinical isolate), *Streptococcus pyogenes* Group A, *Xanthomonas maltophilia* (clinical isolate).

\* **VIRUCIDAL ACTIVITY** - This product when used on environmental, inanimate hard non porous surfaces at 4 ounces per gallon of water exhibits effective virucidal activity against Herpes Simplex Type 1 (causative agent of fever blisters), Herpes Simplex Type 2 (genital disease), Influenza A<sub>2</sub>/hong kong, Vaccinia, Pseudorabies, Bovine Rhinotracheitis, Feline Leukemia, Feline Picornavirus, and Canine Distemper.

Stepan Company  
Northfield, IL 60093  
EPA Reg. No. 1839-168

(left panel)



**KILLS HIV-1 ON PRE-CLEANED ENVIRONMENTAL SURFACES/ OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings (Hospitals, Nursing Homes) or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS).

**SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.**

**PERSONAL PROTECTION:** When handling items soiled with blood or body fluids use disposable latex gloves, gowns, masks, or eye coverings.

**CLEANING PROCEDURES:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product.

**CONTACT TIME:** Allow surface to remain wet for 10 minutes.

**DISPOSAL OF INFECTIOUS MATERIALS:** Blood and other body fluids should be autoclaved and disposed of according to local regulations for infectious waste disposal.

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high-level disinfection.

## **PRECAUTIONARY STATEMENTS**

### **HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**DANGER. KEEP OUT OF REACH OF CHILDREN. CORROSIVE.** Causes eye and skin damage. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food. Remove contaminated clothing and wash before reuse. Wash thoroughly with soap and water after handling.

### **STATEMENT OF PRACTICAL TREATMENT**

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician.

If swallowed, drink egg whites or gelatin solution; or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

### **NOTE TO PHYSICIAN**

Probable mucosal damage may contraindicate the use of gastric lavage.

## **STORAGE AND DISPOSAL**

### **DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL**

**STORAGE** - Store in a dry place no lower in temperature than 50°F or higher than 120°F.

**CONTAINER DISPOSAL** - Do not reuse empty container. Triple rinse empty container with water. Return metal drum then offer for reconditioning or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers may be disposed of in a sanitary landfill, incinerated, or if allowed by local authorities, by burning. If burned, stay out of smoke.

**PESTICIDE DISPOSAL** - Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office of guidance.

Stepan Company  
Northfield, IL 60093  
EPA Reg. No. 1839-168

(right panel)

ACCEPTED

APR 12 1995

1839-168